



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:52 AM UTC

PDB ID : 2O19 / pdb_00002o19
Title : Structure of E. coli topoisomerase III in complex with an 8-base single stranded oligonucleotide. Frozen in glycerol at pH 5.5
Authors : Changela, A.; DiGate, R.; Mondragon, A.
Deposited on : 2006-11-28
Resolution : 2.45 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	8	12% 62% 25% 12%
1	D	8	88% 12%
2	A	659	8% 84% 11% .
2	B	659	2% 87% 9% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	7	Total	C	N	O	P	0	0	0
			138	67	26	39	6			
1	D	8	Total	C	N	O	P	0	0	0
			158	77	28	46	7			

- Molecule 2 is a protein called DNA topoisomerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	630	Total	C	N	O	S	0	0	0
			5013	3165	912	916	20			
2	B	631	Total	C	N	O	S	0	0	0
			5016	3168	913	915	20			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	654	HIS	-	expression tag	UNP P14294
A	655	HIS	-	expression tag	UNP P14294
A	656	HIS	-	expression tag	UNP P14294
A	657	HIS	-	expression tag	UNP P14294
A	658	HIS	-	expression tag	UNP P14294
A	659	HIS	-	expression tag	UNP P14294
B	654	HIS	-	expression tag	UNP P14294
B	655	HIS	-	expression tag	UNP P14294
B	656	HIS	-	expression tag	UNP P14294
B	657	HIS	-	expression tag	UNP P14294
B	658	HIS	-	expression tag	UNP P14294
B	659	HIS	-	expression tag	UNP P14294

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	5	Total	O	0	0
			5	5		
5	D	10	Total	O	0	0
			10	10		
5	A	92	Total	O	0	0
			92	92		
5	B	112	Total	O	0	0
			112	112		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'




- Molecule 1: 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'




- Molecule 2: DNA topoisomerase 3



- Molecule 2: DNA topoisomerase 3





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	102.16Å 102.16Å 451.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.05 – 2.45 29.05 – 2.45	Depositor EDS
% Data completeness (in resolution range)	98.2 (29.05-2.45) 98.1 (29.05-2.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.212 , 0.246 0.206 , 0.238	Depositor DCC
R_{free} test set	4391 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10550	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.48	0/154	1.01	0/235
1	D	0.45	0/176	1.18	0/269
2	A	0.61	1/5123 (0.0%)	0.86	0/6945
2	B	0.60	0/5127	0.85	0/6952
All	All	0.60	1/10580 (0.0%)	0.87	0/14401

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	156	VAL	CA-CB	5.17	1.56	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	292	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	138	0	80	2	0
1	D	158	0	92	1	0
2	A	5013	0	5014	38	0
2	B	5016	0	5020	30	0
3	A	2	0	0	0	0
4	B	4	0	3	0	0
5	A	92	0	0	2	0
5	B	112	0	0	0	0
5	C	5	0	0	0	0
5	D	10	0	0	2	0
All	All	10550	0	10209	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:PHE:HB2	2:B:38:VAL:HG12	1.43	0.97
1:D:4:DA:N3	5:D:196:HOH:O	2.07	0.87
2:B:579:THR:O	2:B:583:GLU:HG2	1.86	0.76
2:B:230:THR:HG22	2:B:232:ALA:H	1.55	0.70
1:C:5:DA:OP1	2:A:524:THR:HG23	1.91	0.69
2:A:467:VAL:HG12	2:A:468:ALA:H	1.58	0.68
2:A:467:VAL:HG12	2:A:468:ALA:N	2.10	0.66
2:B:4:PHE:CB	2:B:38:VAL:HG12	2.26	0.63
2:A:361:GLN:HG2	2:A:449:LEU:HD21	1.83	0.61
2:A:197:ARG:HD3	2:A:574:TRP:CZ3	2.35	0.61
2:B:15:ILE:HG23	2:B:146:ILE:HD11	1.81	0.61
1:C:2:DG:N7	2:A:178:ARG:HD2	2.16	0.60
2:A:197:ARG:HD3	2:A:574:TRP:CE3	2.35	0.60
2:A:73:TRP:H	2:A:646:GLU:HG2	1.66	0.59
2:A:100:HIS:CE1	2:A:112:VAL:HB	2.40	0.56
2:B:586:CYS:HB2	2:B:590:ASP:OD2	2.04	0.56
2:A:187:ALA:HB2	2:A:611:VAL:HG13	1.87	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:259:ARG:N	2:A:260:PRO:HD2	2.21	0.55
2:B:250:GLN:HA	2:B:255:ARG:O	2.09	0.53
2:A:467:VAL:CG1	2:A:468:ALA:H	2.22	0.53
2:A:520:ASP:HB3	2:A:525:GLU:OE2	2.09	0.52
2:A:176:MET:CE	2:A:603:ILE:HD13	2.39	0.52
2:A:150:ARG:HD2	5:A:875:HOH:O	2.10	0.51
2:B:177:THR:HG23	2:B:194:SER:HA	1.93	0.50
2:A:276:VAL:HG21	2:A:467:VAL:HG11	1.93	0.50
2:B:507:VAL:O	2:B:513:LYS:NZ	2.43	0.50
2:B:515:ILE:HD13	2:B:576:SER:HB2	1.94	0.49
2:A:646:GLU:HB2	2:A:649:SER:HB3	1.94	0.49
2:B:100:HIS:NE2	2:B:112:VAL:HB	2.28	0.48
2:A:177:THR:HG23	2:A:194:SER:HA	1.95	0.48
2:B:230:THR:HG21	2:B:234:GLU:OE1	2.14	0.48
2:A:507:VAL:HG23	2:A:513:LYS:HG3	1.96	0.47
2:B:230:THR:HB	2:B:234:GLU:H	1.79	0.47
2:A:291:PRO:HD3	2:A:413:GLN:CD	2.39	0.47
2:A:467:VAL:CG1	2:A:468:ALA:N	2.77	0.47
2:B:100:HIS:CE1	2:B:112:VAL:HB	2.50	0.47
2:A:176:MET:HA	2:A:176:MET:HE2	1.96	0.46
2:A:299:ILE:HD11	2:A:498:SER:HA	1.96	0.46
2:A:137:LEU:HD12	2:A:321:GLU:HA	1.98	0.46
2:A:151:SER:O	2:A:154:GLU:HG2	2.15	0.46
2:B:649:SER:HA	2:B:652:ILE:HD12	1.99	0.45
2:B:2:ARG:HG2	2:B:97:GLU:HB2	1.98	0.45
2:A:98:ILE:HB	2:A:129:VAL:HG22	1.98	0.45
2:B:585:GLN:O	2:B:586:CYS:HB3	2.17	0.45
2:A:228:ILE:CG2	2:A:229:VAL:N	2.79	0.45
2:A:75:LEU:HG	2:A:647:VAL:HG21	1.98	0.45
2:B:400:ALA:O	2:B:404:ASN:HB2	2.16	0.44
2:A:247:GLU:N	2:A:248:PRO:HD2	2.33	0.43
2:A:46:LEU:HD23	2:A:77:PRO:HA	1.99	0.43
2:B:3:LEU:HD11	2:B:39:THR:HG22	2.00	0.43
2:B:89:LYS:HD3	2:B:119:LEU:HD22	2.00	0.43
2:A:365:ASP:OD2	2:A:365:ASP:C	2.62	0.42
2:A:365:ASP:HB3	2:A:368:ILE:HD12	2.00	0.42
2:A:72:LYS:HB3	2:A:72:LYS:HE2	1.73	0.42
2:A:208:ARG:NH2	2:A:567:ARG:HE	2.17	0.42
2:A:369:ARG:HG3	2:A:373:TRP:CD1	2.54	0.42
2:B:389:ARG:HH21	2:B:393:ILE:HD11	1.85	0.42
5:D:54:HOH:O	2:B:48:GLN:HG3	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:122:ALA:HA	2:B:123:PRO:HD3	1.88	0.42
2:B:230:THR:HG23	2:B:231:PRO:HD2	2.02	0.42
2:B:605:GLN:HG3	2:B:608:ARG:HH22	1.85	0.42
2:B:276:VAL:HA	2:B:427:LEU:HD23	2.02	0.41
2:A:495:THR:HB	5:A:886:HOH:O	2.19	0.41
2:B:106:ARG:HG3	2:B:162:ALA:HB2	2.03	0.41
2:B:544:LYS:O	2:B:544:LYS:HG3	2.19	0.41
2:B:55:ASP:OD1	2:B:57:ARG:HB2	2.21	0.41
2:B:354:HIS:HE1	2:B:389:ARG:O	2.04	0.41
2:A:7:GLU:H	2:A:11:LEU:HD23	1.86	0.40
2:A:187:ALA:HB3	2:A:614:PHE:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	626/659 (95%)	611 (98%)	15 (2%)	0	100	100
2	B	627/659 (95%)	609 (97%)	18 (3%)	0	100	100
All	All	1253/1318 (95%)	1220 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	535/555 (96%)	516 (96%)	19 (4%)	31	45
2	B	535/555 (96%)	522 (98%)	13 (2%)	43	58
All	All	1070/1110 (96%)	1038 (97%)	32 (3%)	36	51

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	124	GLU
2	A	126	ARG
2	A	237	THR
2	A	257	LEU
2	A	258	HIS
2	A	264	HIS
2	A	363	VAL
2	A	391	SER
2	A	454	GLU
2	A	456	ASP
2	A	457	GLU
2	A	460	ASP
2	A	498	SER
2	A	586	CYS
2	A	605	GLN
2	A	611	VAL
2	A	612	ARG
2	A	647	VAL
2	A	654	HIS
2	B	2	ARG
2	B	18	VAL
2	B	121	LEU
2	B	148	ARG
2	B	244	GLU
2	B	253	GLU
2	B	452	SER
2	B	462	THR
2	B	473	LEU
2	B	544	LYS
2	B	601	GLN
2	B	604	ASP
2	B	658	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	50	GLN
2	A	76	GLN
2	A	93	HIS
2	A	135	ASN
2	A	199	GLN
2	A	280	ASN
2	A	323	HIS
2	A	593	GLN
2	A	655	HIS
2	B	227	HIS
2	B	241	GLN
2	B	354	HIS
2	B	370	ASN
2	B	413	GLN
2	B	585	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACY	B	803	-	3,3,3	0.88	0	3,3,3	0.99	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	7/8 (87%)	-0.55	1 (14%) 6 4	21, 23, 46, 75	0
1	D	8/8 (100%)	-1.12	0 100 100	20, 26, 44, 56	0
2	A	630/659 (95%)	0.19	56 (8%) 15 13	19, 42, 115, 173	0
2	B	631/659 (95%)	-0.17	15 (2%) 59 62	16, 39, 80, 114	0
All	All	1276/1334 (95%)	0.00	72 (5%) 30 27	16, 40, 89, 173	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	654	HIS	6.9
2	A	656	HIS	6.8
2	B	658	HIS	6.5
2	A	658	HIS	6.1
2	A	277	THR	5.5
2	A	647	VAL	5.4
2	A	275	ILE	5.1
2	B	656	HIS	5.1
2	A	651	ALA	5.1
2	A	652	ILE	5.0
2	A	650	GLY	4.8
2	B	654	HIS	4.7
2	A	232	ALA	4.7
2	A	257	LEU	4.6
2	A	245	ALA	4.6
2	A	264	HIS	4.6
2	A	653	ALA	4.4
2	B	648	GLY	4.2
2	A	243	SER	4.1
2	A	655	HIS	4.1
2	A	274	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	276	VAL	3.9
2	A	231	PRO	3.9
2	A	234	GLU	3.8
2	A	254	GLY	3.7
2	A	240	TRP	3.6
2	A	657	HIS	3.6
2	A	258	HIS	3.5
2	A	218	ALA	3.5
2	A	646	GLU	3.5
2	A	453	LYS	3.4
2	A	455	ARG	3.4
2	A	252	GLU	3.4
2	A	253	GLU	3.3
2	A	230	THR	3.3
2	B	655	HIS	3.2
2	A	452	SER	3.1
2	B	619	ALA	3.1
2	A	648	GLY	2.9
2	B	618	VAL	2.9
2	B	258	HIS	2.9
2	A	229	VAL	2.8
2	B	657	HIS	2.8
2	A	248	PRO	2.7
2	A	649	SER	2.7
2	A	256	LEU	2.7
2	A	544	LYS	2.6
2	A	249	TYR	2.6
2	A	456	ASP	2.6
2	B	652	ILE	2.5
2	A	217	VAL	2.5
1	C	7	DT	2.5
2	A	462	THR	2.5
2	A	241	GLN	2.4
2	A	244	GLU	2.4
2	A	432	GLY	2.3
2	B	35	GLY	2.3
2	A	224	VAL	2.3
2	B	126	ARG	2.3
2	A	260	PRO	2.3
2	A	457	GLU	2.3
2	A	271	GLY	2.2
2	B	620	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	653	ALA	2.2
2	A	486	GLN	2.2
2	A	238	ALA	2.1
2	A	255	ARG	2.1
2	A	233	ASP	2.1
2	A	220	ASP	2.1
2	A	251	ASP	2.1
2	A	124	GLU	2.0
2	B	649	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACY	B	803	4/4	0.74	0.18	36,37,38,38	0
3	CL	A	801	1/1	0.99	0.02	21,21,21,21	0
3	CL	A	800	1/1	0.99	0.03	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.