



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:59 AM UTC

PDB ID : 6O1Z / pdb_00006o1z
Title : Structure of pCW3 conjugation coupling protein TcpA hexagonal crystal form
Authors : Traore, D.A.K.; Ahktar, N.; Torres, V.T.; Adams, V.; Coulibaly, F.; Panjkar, S.; Caradoc-Davies, T.T.; Rood, J.I.; Whisstock, J.C.
Deposited on : 2019-02-22
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	71% 25% ..

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA translocase coupling protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	0
			2854	1824	482	541	7			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	SER	-	expression tag	UNP Q1PLI0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA translocase coupling protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, α , β , γ	108.50Å 108.50Å 148.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.98 – 3.10 46.98 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.98-3.10) 99.7 (46.98-3.10)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 3.12Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.197 , 0.275 0.210 , 0.279	Depositor DCC
R_{free} test set	492 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtriage
Anisotropy	0.689	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 102.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2854	wwPDB-VP
Average B, all atoms (Å ²)	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	0/2897	1.45	23/3895 (0.6%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	347	ASP	CA-CB-CG	6.68	119.28	112.60
1	A	367	SER	N-CA-C	6.04	118.36	111.11
1	A	376	ILE	N-CA-C	6.04	116.62	108.17
1	A	112	TYR	CA-C-N	5.78	129.49	120.82
1	A	112	TYR	C-N-CA	5.78	129.49	120.82
1	A	355	ARG	CA-C-N	5.56	128.08	120.46
1	A	355	ARG	C-N-CA	5.56	128.08	120.46
1	A	392	ASN	CA-CB-CG	5.54	118.14	112.60
1	A	463	GLU	CA-C-N	5.45	131.94	121.54
1	A	463	GLU	C-N-CA	5.45	131.94	121.54
1	A	382	ASP	CA-CB-CG	5.44	118.04	112.60
1	A	460	ASP	CA-CB-CG	5.25	117.85	112.60
1	A	299	ASN	CA-C-N	5.24	127.56	120.44
1	A	299	ASN	C-N-CA	5.24	127.56	120.44
1	A	298	GLU	CA-C-N	5.21	127.27	120.28
1	A	298	GLU	C-N-CA	5.21	127.27	120.28
1	A	366	LEU	CA-C-N	5.15	127.49	120.54
1	A	366	LEU	C-N-CA	5.15	127.49	120.54
1	A	309	SER	N-CA-C	-5.14	106.27	112.54
1	A	126	SER	CA-C-N	5.11	128.03	120.82
1	A	126	SER	C-N-CA	5.11	128.03	120.82
1	A	116	PHE	CA-C-N	5.04	127.29	120.38
1	A	116	PHE	C-N-CA	5.04	127.29	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2854	0	2940	29	0
All	All	2854	0	2940	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ARG:O	1:A:117:GLU:HB2	1.96	0.65
1:A:273:LEU:HA	1:A:276:GLU:HG3	1.84	0.60
1:A:233:LEU:HB2	1:A:394:VAL:HG11	1.87	0.57
1:A:330:ILE:HD11	1:A:375:CYS:SG	2.44	0.57
1:A:378:VAL:HG21	1:A:386:ILE:HD11	1.87	0.57
1:A:409:GLU:O	1:A:413:SER:HA	2.05	0.56
1:A:211:VAL:HA	1:A:224:ILE:O	2.06	0.56
1:A:167:LEU:HB2	1:A:169:ILE:HG12	1.88	0.55
1:A:397:ARG:HB2	1:A:430:LYS:HB3	1.89	0.54
1:A:449:PHE:CE2	1:A:451:PRO:HG3	2.45	0.51
1:A:149:SER:HB2	1:A:151:ILE:HD12	1.93	0.50
1:A:164:GLN:HE21	1:A:170:SER:HA	1.78	0.49
1:A:336:LEU:HG	1:A:340:MET:HG3	1.94	0.49
1:A:458:ARG:O	1:A:462:ILE:HG23	2.13	0.48
1:A:365:ARG:HG3	1:A:393:ASN:ND2	2.29	0.48
1:A:362:THR:O	1:A:366:LEU:HG	2.15	0.47
1:A:331:VAL:HB	1:A:374:LEU:HD23	1.96	0.47
1:A:391:LYS:HE2	1:A:397:ARG:HH12	1.81	0.46
1:A:232:ILE:HD13	1:A:249:ILE:HD13	1.99	0.44
1:A:124:ALA:HB3	1:A:150:MET:HE3	2.00	0.43
1:A:246:LEU:HD22	1:A:375:CYS:HB3	2.00	0.43
1:A:214:VAL:HG12	1:A:249:ILE:HD11	2.01	0.43
1:A:407:ALA:HA	1:A:410:ILE:HD12	2.00	0.43
1:A:421:PRO:HG2	1:A:426:ARG:HG2	2.00	0.42
1:A:152:PRO:HB2	1:A:154:GLU:HG2	2.02	0.42
1:A:428:LEU:HD23	1:A:437:GLN:HA	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:GLN:HE21	1:A:171:ILE:H	1.68	0.41
1:A:444:ASP:HB3	1:A:447:LYS:HG2	2.03	0.41
1:A:123:GLY:HA2	1:A:151:ILE:HG13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/365 (98%)	323 (90%)	30 (8%)	5 (1%)	9	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	343	THR
1	A	368	ARG
1	A	464	ASP
1	A	395	PRO
1	A	192	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/317 (98%)	275 (88%)	37 (12%)	5	21

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	113	ARG
1	A	117	GLU
1	A	127	LYS
1	A	135	ILE
1	A	154	GLU
1	A	167	LEU
1	A	183	VAL
1	A	184	ILE
1	A	194	ILE
1	A	197	TYR
1	A	206	GLU
1	A	237	GLU
1	A	240	SER
1	A	244	VAL
1	A	245	ILE
1	A	274	GLU
1	A	284	GLU
1	A	286	ASP
1	A	296	VAL
1	A	304	LYS
1	A	307	ARG
1	A	311	SER
1	A	327	LYS
1	A	331	VAL
1	A	332	VAL
1	A	339	LEU
1	A	348	GLU
1	A	355	ARG
1	A	378	VAL
1	A	389	GLN
1	A	391	LYS
1	A	398	ILE
1	A	399	CYS
1	A	405	SER
1	A	434	ASP
1	A	456	LYS
1	A	462	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	164	GLN
1	A	216	GLN
1	A	221	ASN
1	A	379	GLN
1	A	392	ASN
1	A	393	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	360/365 (98%)	-0.10	2 (0%)	85 70	66, 115, 154, 181	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	318	ASN	2.3
1	A	322	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.