



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 12:31 AM UTC

PDB ID : 1O2G / pdb_00001o2g
Title : Elaborate Manifold of Short Hydrogen Bond Arrays Mediating Binding of Active Site-Directed Serine Protease Inhibitors
Authors : Katz, B.A.; Elrod, K.; Verner, E.; Mackman, R.L.; Luong, C.; Shrader, W.; Sendzik, M.; Spencer, J.R.; Sprengeler, P.A.; Kolesnikov, A.; Tai, W.F.; Hui, H.; Breitenbucher, G.; Allen, D.; Janc, J.
Deposited on : 2003-03-06
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	36	25% 53% 25% 19% .
2	H	259	% 68% 25% . .
3	I	11	18% 45% 45% 9%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5815 atoms, of which 3041 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	36	Total	C	H	N	O	S	0	0	0
			565	177	278	48	61	1			

- Molecule 2 is a protein called THROMBIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	251	Total	C	H	N	O	S	0	5	0
			4089	1310	2042	359	364	14			


- Molecule 3 is a protein called ACETYL HIRUDIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	I	11	Total	C	H	N	O	S	0	0	0
			188	64	84	12	27	1			

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Na	0	0
			1	1		

- Molecule 5 is 3-{5-[AMINO(IMINIO)METHYL]-1H-INDOL-2-YL}-1,1'-BIPHENYL-2-OLATE (CCD ID: 696) (formula: C₂₁H₁₇N₃O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	H	1	Total	C	H	N	O	0	0
			42	21	17	3	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	50	Total	H	O	0	0
			150	100	50		
6	H	250	Total	H	O	0	1
			756	504	252		
6	I	8	Total	H	O	0	0
			24	16	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: THROMBIN



• Molecule 2: THROMBIN



• Molecule 3: ACETYL HIRUDIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.36Å 72.06Å 72.96Å 90.00° 101.77° 90.00°	Depositor
Resolution (Å)	7.00 – 1.58 7.00 – 1.58	Depositor EDS
% Data completeness (in resolution range)	87.3 (7.00-1.58) 86.3 (7.00-1.58)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.59 (at 1.58Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.219 , 0.250 0.213 , 0.208	Depositor DCC
R_{free} test set	4349 reflections (10.17%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.216	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5815	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TYS, 696

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	1.82	3/290 (1.0%)	1.92	6/384 (1.6%)
2	H	1.78	21/2123 (1.0%)	1.74	47/2866 (1.6%)
3	I	1.62	0/88	1.98	6/115 (5.2%)
All	All	1.78	24/2501 (1.0%)	1.77	59/3365 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2
2	H	0	7
All	All	0	9

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	230	HIS	CG-ND1	-8.94	1.28	1.38
2	H	71	HIS	CG-ND1	-8.45	1.28	1.38
2	H	119	HIS	CG-ND1	-8.17	1.29	1.38
2	H	19	GLY	N-CA	7.72	1.50	1.44
2	H	91	HIS	CD2-NE2	-7.38	1.29	1.37
2	H	57	HIS	CG-ND1	-7.09	1.30	1.38
2	H	29	TRP	NE1-CE2	-6.23	1.30	1.37
2	H	91	HIS	CG-ND1	-6.19	1.31	1.38
2	H	230	HIS	CD2-NE2	-5.89	1.31	1.37
2	H	241	VAL	CA-CB	5.88	1.61	1.54
2	H	57	HIS	CD2-NE2	-5.75	1.31	1.37
2	H	231	VAL	CA-CB	5.74	1.61	1.54
2	H	237	TRP	NE1-CE2	-5.72	1.31	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	207	TRP	NE1-CE2	-5.71	1.31	1.37
2	H	71	HIS	CD2-NE2	-5.54	1.31	1.37
2	H	48	SER	CA-CB	5.52	1.62	1.53
2	H	60(D)	TRP	NE1-CE2	-5.52	1.31	1.37
1	L	14(G)	LEU	CB-CG	5.48	1.64	1.53
1	L	1(B)	ALA	N-CA	5.35	1.53	1.46
2	H	119	HIS	CD2-NE2	-5.12	1.32	1.37
2	H	211	GLY	N-CA	5.10	1.50	1.45
2	H	141	TRP	NE1-CE2	-5.06	1.31	1.37
2	H	51	TRP	NE1-CE2	-5.04	1.31	1.37
1	L	1(F)	GLY	N-CA	5.01	1.52	1.45

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	1(E)	SER	N-CA-C	-16.35	75.98	110.80
3	I	56	PHE	CA-CB-CG	-9.86	103.94	113.80
2	H	232	PHE	CA-CB-CG	-9.30	104.50	113.80
2	H	199	PHE	N-CA-C	-8.48	94.59	108.41
2	H	227	PHE	CA-CB-CG	-8.16	105.64	113.80
2	H	181	PHE	CA-CB-CG	8.14	121.94	113.80
1	L	14(A)	LYS	N-CA-C	7.88	121.99	112.38
2	H	205	ASN	CA-CB-CG	-7.67	104.94	112.60
2	H	57	HIS	CA-CB-CG	7.50	121.30	113.80
2	H	123	LEU	N-CA-C	-7.43	100.06	110.31
1	L	1(D)	GLY	N-CA-C	7.43	130.79	113.18
2	H	33	LEU	N-CA-C	-7.41	96.65	108.73
2	H	190	ALA	N-CA-C	-7.38	98.92	110.36
2	H	230	HIS	CE1-NE2-CD2	-7.14	101.86	109.00
2	H	71	HIS	CE1-NE2-CD2	-6.99	102.01	109.00
2	H	119	HIS	CE1-NE2-CD2	-6.97	102.03	109.00
2	H	91	HIS	ND1-CG-CD2	6.87	112.97	106.10
2	H	97(A)	GLU	N-CA-C	6.71	121.32	112.13
2	H	244	GLN	N-CA-C	-6.63	105.31	113.41
2	H	154	VAL	N-CA-CB	-6.58	101.42	112.47
2	H	86	GLU	N-CA-C	-6.57	102.70	111.96
2	H	43	GLY	N-CA-C	-6.52	101.36	112.51
2	H	78	ASN	OD1-CG-ND2	-6.40	116.20	122.60
2	H	153	SER	N-CA-C	-6.21	105.84	113.41
2	H	173	ARG	N-CA-C	-6.16	105.04	113.30
2	H	64	LEU	N-CA-C	6.14	119.55	109.85
2	H	181	PHE	N-CA-C	-6.12	99.76	109.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	40	LEU	N-CA-C	-6.07	100.85	109.96
2	H	76	TYR	N-CA-C	-6.06	98.66	108.41
2	H	91	HIS	CA-CB-CG	-6.05	107.75	113.80
2	H	74	THR	N-CA-C	5.94	120.69	113.38
1	L	14(L)	ASP	CA-CB-CG	5.94	118.54	112.60
2	H	186(D)	LYS	N-CA-C	-5.91	100.36	109.52
2	H	60(G)	ASN	CA-CB-CG	-5.79	106.81	112.60
2	H	19	GLY	N-CA-C	-5.74	104.89	112.81
1	L	14(K)	ILE	N-CA-C	5.72	121.23	109.34
2	H	71	HIS	CA-CB-CG	-5.71	108.09	113.80
3	I	65	GLN	OE1-CD-NE2	-5.67	116.93	122.60
1	L	14(J)	TYR	N-CA-C	-5.64	102.84	110.68
2	H	125	ASP	N-CA-C	-5.61	100.98	109.85
2	H	23	GLU	N-CA-C	-5.56	101.80	110.42
2	H	162	ILE	N-CA-C	-5.56	101.88	109.55
2	H	205	ASN	N-CA-C	5.53	119.61	112.86
2	H	173	ARG	CB-CA-C	-5.50	100.88	110.17
2	H	60(G)	ASN	OD1-CG-ND2	-5.49	117.11	122.60
3	I	64	LEU	CA-C-N	5.46	131.52	121.70
3	I	64	LEU	C-N-CA	5.46	131.52	121.70
2	H	219	GLY	N-CA-C	-5.43	102.98	112.53
3	I	64	LEU	O-C-N	5.42	131.66	123.00
2	H	94	TYR	N-CA-C	-5.36	101.92	109.96
2	H	187	ARG	NE-CZ-NH1	-5.33	116.17	121.50
2	H	204(A)	PHE	CA-CB-CG	-5.31	108.49	113.80
2	H	60(B)	PRO	N-CA-C	5.29	117.16	110.70
2	H	136	GLY	N-CA-C	-5.29	103.27	112.06
3	I	57	GLU	N-CA-C	-5.25	101.97	110.32
2	H	119	HIS	CA-CB-CG	-5.14	108.66	113.80
2	H	38	GLN	OE1-CD-NE2	-5.13	117.47	122.60
2	H	79	ILE	N-CA-CB	-5.12	107.44	111.64
2	H	237	TRP	N-CA-C	-5.04	105.87	111.36

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	101	ARG	Sidechain
2	H	126	ARG	Sidechain
2	H	173	ARG	Sidechain
2	H	187	ARG	Sidechain
2	H	233	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	H	50	ARG	Sidechain
2	H	67	ARG	Sidechain
1	L	14(D)	ARG	Sidechain
1	L	15	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	287	278	278	7	0
2	H	2047	2042	2024	21	2
3	I	104	84	81	1	0
4	H	1	0	0	0	0
5	H	25	17	17	4	0
6	H	252	504	0	2	2
6	I	8	16	0	0	0
6	L	50	100	0	0	0
All	All	2774	3041	2400	24	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:195:SER:HG	5:H:410:696:HN3	1.24	0.81
2:H:236:LYS:HE2	6:H:631:HOH:O	1.87	0.73
1:L:1(G):PHE:CE1	2:H:239:GLN:HA	2.30	0.67
1:L:1(C):GLU:HB3	1:L:1:CYS:HB3	1.80	0.62
3:I:58:GLU:H	3:I:58:GLU:CD	2.06	0.62
2:H:195:SER:OG	5:H:410:696:O6'	2.14	0.59
2:H:195:SER:OG	5:H:410:696:N3	2.29	0.57
2:H:169:LYS:NZ	6:H:446:HOH:O	2.39	0.55
2:H:192:GLU:HG3	5:H:410:696:C2'	2.37	0.55
1:L:1(G):PHE:CD1	2:H:239:GLN:HG2	2.41	0.54
2:H:204(B):ASN:C	2:H:204(B):ASN:HD22	2.19	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:1(G):PHE:CE1	2:H:242:ILE:HB	2.50	0.47
2:H:187:ARG:NH2	2:H:221:ASP:O	2.49	0.46
2:H:85:LEU:HD13	2:H:106:MET:HE2	1.98	0.45
2:H:17:VAL:O	2:H:188:GLY:HA2	2.16	0.45
2:H:136:GLY:HA3	2:H:199:PHE:CZ	2.53	0.44
2:H:22:ALA:HB2	2:H:157[A]:VAL:CG2	2.48	0.44
1:L:1(G):PHE:CZ	2:H:242:ILE:CG2	3.01	0.44
2:H:35:ARG:O	2:H:38:GLN:HA	2.19	0.43
1:L:14(A):LYS:HB2	1:L:14(A):LYS:NZ	2.35	0.42
2:H:204(B):ASN:C	2:H:204(B):ASN:ND2	2.77	0.41
2:H:80:GLU:C	2:H:81:LYS:HD2	2.45	0.41
1:L:1(H):THR:O	2:H:123:LEU:HD12	2.21	0.41
2:H:29:TRP:CG	2:H:121:VAL:HB	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:77(A):ARG:HE	6:H:451:HOH:H2[2_555]	1.35	0.25
2:H:172:THR:O	6:H:542:HOH:H1[4_546]	1.48	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	34/36 (94%)	27 (79%)	4 (12%)	3 (9%)	0	0
2	H	252/259 (97%)	245 (97%)	7 (3%)	0	100	100
3	I	8/11 (73%)	6 (75%)	2 (25%)	0	100	100
All	All	294/306 (96%)	278 (95%)	13 (4%)	3 (1%)	12	2

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	1(B)	ALA
1	L	1(E)	SER
1	L	14(K)	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	31/31 (100%)	24 (77%)	7 (23%)	1	0
2	H	224/225 (100%)	218 (97%)	6 (3%)	39	10
3	I	10/10 (100%)	9 (90%)	1 (10%)	7	0
All	All	265/266 (100%)	251 (95%)	14 (5%)	20	2

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1(H)	THR
1	L	1(G)	PHE
1	L	1(E)	SER
1	L	1(C)	GLU
1	L	1(A)	ASP
1	L	1	CYS
1	L	6	LEU
2	H	46	LEU
2	H	50	ARG
2	H	65	LEU
2	H	154	VAL
2	H	205	ASN
2	H	241	VAL
3	I	65	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	204(B)	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	239	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TYS	I	63	3	15,16,17	0.92	1 (6%)	15,22,24	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	I	63	3	-	0/10/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	63	TYS	OH-CZ	-2.52	1.38	1.42

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	696	H	410	-	28,28,28	2.22	8 (28%)	36,40,40	2.50	7 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	696	H	410	-	-	0/12/12/12	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	410	696	C4-N3	-5.33	1.29	1.38
5	H	410	696	C1'-C8	-5.03	1.38	1.47
5	H	410	696	C1-C7	-3.93	1.40	1.47
5	H	410	696	C5'-C1B	-3.90	1.42	1.49
5	H	410	696	C8-N3	-3.64	1.34	1.38
5	H	410	696	CN4-C8	3.36	1.43	1.37
5	H	410	696	C2-C1	2.74	1.43	1.39
5	H	410	696	C6-C5	-2.41	1.36	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	410	696	C5-CN4-C8	-10.07	96.70	107.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	410	696	C6-C5-C4	6.40	123.08	119.13
5	H	410	696	C4-C5-CN4	5.05	110.94	106.80
5	H	410	696	CN4-C8-N3	4.60	114.58	108.65
5	H	410	696	C3-C4-C5	-2.32	120.03	122.13
5	H	410	696	C6-C5-CN4	-2.13	125.37	132.70
5	H	410	696	C1'-C8-CN4	-2.12	120.88	128.60

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	410	696	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	36/36 (100%)	0.70	9 (25%) 2 2	1, 20, 35, 82	12 (33%)
2	H	251/259 (96%)	-0.34	2 (0%) 82 87	10, 22, 39, 53	29 (11%)
3	I	10/11 (90%)	0.93	2 (20%) 3 4	15, 39, 42, 46	5 (50%)
All	All	297/306 (97%)	-0.17	13 (4%) 39 51	1, 22, 41, 82	46 (15%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	14(L)	ASP	6.2
1	L	14(M)	GLY	5.9
1	L	1(F)	GLY	5.5
1	L	1(G)	PHE	5.0
1	L	1(B)	ALA	3.1
1	L	15	ARG	3.1
1	L	1(D)	GLY	3.0
2	H	246	GLY	3.0
2	H	247	GLU	2.8
1	L	1(H)	THR	2.2
1	L	1(E)	SER	2.2
3	I	61	GLU	2.2
3	I	64	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TYS	I	63	16/17	0.96	0.06	29,35,46,47	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NA	H	409	1/1	0.93	0.06	29,29,29,29	0
5	696	H	410	25/25	0.94	0.05	15,25,27,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.