



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:31 PM UTC

PDB ID : 2O26 / pdb_00002o26
Title : Structure of a class III RTK signaling assembly
Authors : Liu, H.; Chen, X.; Focia, P.J.; He, X.
Deposited on : 2006-11-29
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	145	59%31%8%..
1	B	145	53%34%8%..
1	E	145	51%37%6%..
1	F	145	56%32%7%..
2	U	290	47%40%7%.5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	W	290	
2	X	290	
2	Y	290	
3	C	3	
3	G	3	
3	I	3	
3	K	3	
4	D	3	
4	H	3	
4	J	3	
4	L	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	L	1	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15104 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kit ligand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	S	0	0	0
			1140	722	189	220	9			
1	B	139	Total	C	N	O	S	0	0	0
			1090	692	174	215	9			
1	E	139	Total	C	N	O	S	0	0	0
			1090	692	174	215	9			
1	F	139	Total	C	N	O	S	0	0	0
			1090	692	174	215	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	HIS	-	expression tag	UNP P20826
A	143	HIS	-	expression tag	UNP P20826
A	144	HIS	-	expression tag	UNP P20826
A	145	HIS	-	expression tag	UNP P20826
A	146	HIS	-	expression tag	UNP P20826
A	147	HIS	-	expression tag	UNP P20826
B	142	HIS	-	expression tag	UNP P20826
B	143	HIS	-	expression tag	UNP P20826
B	144	HIS	-	expression tag	UNP P20826
B	145	HIS	-	expression tag	UNP P20826
B	146	HIS	-	expression tag	UNP P20826
B	147	HIS	-	expression tag	UNP P20826
E	142	HIS	-	expression tag	UNP P20826
E	143	HIS	-	expression tag	UNP P20826
E	144	HIS	-	expression tag	UNP P20826
E	145	HIS	-	expression tag	UNP P20826
E	146	HIS	-	expression tag	UNP P20826
E	147	HIS	-	expression tag	UNP P20826
F	142	HIS	-	expression tag	UNP P20826
F	143	HIS	-	expression tag	UNP P20826
F	144	HIS	-	expression tag	UNP P20826

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	145	HIS	-	expression tag	UNP P20826
F	146	HIS	-	expression tag	UNP P20826
F	147	HIS	-	expression tag	UNP P20826


- Molecule 2 is a protein called Mast/stem cell growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	275	Total	C	N	O	S	0	0	0
			2164	1369	372	411	12			
2	Y	275	Total	C	N	O	S	0	0	0
			2164	1369	372	411	12			
2	U	275	Total	C	N	O	S	0	0	0
			2164	1369	372	411	12			
2	W	275	Total	C	N	O	S	0	0	0
			2164	1369	372	411	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
U	146	GLN	ASN	engineered mutation	UNP P05532
W	146	GLN	ASN	engineered mutation	UNP P05532
X	146	GLN	ASN	engineered mutation	UNP P05532
Y	146	GLN	ASN	engineered mutation	UNP P05532

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O		0	0	0
			38	22	2	14				
3	G	3	Total	C	N	O		0	0	0
			38	22	2	14				
3	I	3	Total	C	N	O		0	0	0
			38	22	2	14				
3	K	3	Total	C	N	O		0	0	0
			38	22	2	14				

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-

beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	H	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	J	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	L	3	Total	C	N	O	0	0	0
			39	22	2	15			


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	171	Total	O	0	0
			171	171		
5	B	144	Total	O	0	0
			144	144		
5	E	152	Total	O	0	0
			152	152		
5	F	143	Total	O	0	0
			143	143		
5	X	273	Total	O	0	0
			273	273		
5	Y	310	Total	O	0	0
			310	310		
5	U	270	Total	O	0	0
			270	270		
5	W	267	Total	O	0	0
			267	267		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Kit ligand




• Molecule 1: Kit ligand



• Molecule 1: Kit ligand







• Molecule 1: Kit ligand







• Molecule 2: Mast/stem cell growth factor receptor



• Molecule 2: Mast/stem cell growth factor receptor




• Molecule 2: Mast/stem cell growth factor receptor





- Molecule 2: Mast/stem cell growth factor receptor



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.85Å 200.15Å 82.02Å 90.00° 91.42° 90.00°	Depositor
Resolution (Å)	19.93 – 2.50 19.93 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.7 (19.93-2.50) 96.6 (19.93-2.50)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.237 , 0.270 0.242 , 0.274	Depositor DCC
R_{free} test set	4160 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	65.2	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.23 , 61.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15104	wwPDB-VP
Average B, all atoms (Å ²)	98.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, FUL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.50	0/1164	1.04	8/1579 (0.5%)
1	B	0.60	1/1109 (0.1%)	1.00	7/1504 (0.5%)
1	E	0.49	0/1109	1.00	7/1504 (0.5%)
1	F	0.49	0/1109	1.03	10/1504 (0.7%)
2	U	0.46	0/2214	1.02	8/3011 (0.3%)
2	W	0.41	0/2214	0.93	8/3011 (0.3%)
2	X	0.44	0/2214	1.00	10/3011 (0.3%)
2	Y	0.42	0/2214	0.95	6/3011 (0.2%)
All	All	0.47	1/13347 (0.0%)	0.99	64/18135 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	133	SER	CA-CB	8.29	1.64	1.53

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	U	138	PRO	CA-N-CD	-15.13	90.81	112.00
2	X	41	PRO	CA-N-CD	-12.80	94.08	112.00
2	X	43	GLN	OE1-CD-NE2	-9.79	112.81	122.60
1	A	102	PRO	CA-N-CD	-9.22	99.09	112.00
1	B	117	SER	N-CA-C	-8.74	101.83	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1140	0	1121	69	0
1	B	1090	0	1086	74	0
1	E	1090	0	1086	69	0
1	F	1090	0	1086	60	0
2	U	2164	0	2140	146	0
2	W	2164	0	2140	121	0
2	X	2164	0	2140	150	0
2	Y	2164	0	2140	132	0
3	C	38	0	34	2	0
3	G	38	0	34	2	0
3	I	38	0	34	3	0
3	K	38	0	34	0	0
4	D	39	0	34	1	0
4	H	39	0	34	2	0
4	J	39	0	34	1	0
4	L	39	0	34	3	0
5	A	171	0	0	25	0
5	B	144	0	0	32	0
5	E	152	0	0	24	0
5	F	143	0	0	24	0
5	U	270	0	0	38	0
5	W	267	0	0	35	0
5	X	273	0	0	50	0
5	Y	310	0	0	48	0
All	All	15104	0	13211	807	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 807 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:184:CYS:HB2	5:W:321:HOH:O	1.50	1.08
2:U:251:ASN:HB2	2:U:252:PRO:HD3	1.36	1.06
2:Y:251:ASN:HB2	2:Y:252:PRO:HD3	1.34	1.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:251:ASN:HB2	2:W:252:PRO:HD3	1.37	1.06
2:X:251:ASN:HB2	2:X:252:PRO:HD3	1.35	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/145 (98%)	124 (87%)	13 (9%)	5 (4%)	3	4
1	B	137/145 (94%)	122 (89%)	10 (7%)	5 (4%)	2	3
1	E	137/145 (94%)	124 (90%)	9 (7%)	4 (3%)	3	5
1	F	137/145 (94%)	121 (88%)	12 (9%)	4 (3%)	3	5
2	U	273/290 (94%)	252 (92%)	15 (6%)	6 (2%)	5	9
2	W	273/290 (94%)	255 (93%)	12 (4%)	6 (2%)	5	9
2	X	273/290 (94%)	253 (93%)	15 (6%)	5 (2%)	6	12
2	Y	273/290 (94%)	249 (91%)	16 (6%)	8 (3%)	3	5
All	All	1645/1740 (94%)	1500 (91%)	102 (6%)	43 (3%)	4	7

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	102	PRO
1	E	102	PRO
1	F	102	PRO
2	X	73	ASN
2	Y	73	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/136 (99%)	123 (91%)	12 (9%)	9	20
1	B	130/136 (96%)	120 (92%)	10 (8%)	12	25
1	E	130/136 (96%)	120 (92%)	10 (8%)	12	25
1	F	130/136 (96%)	122 (94%)	8 (6%)	16	34
2	U	247/259 (95%)	228 (92%)	19 (8%)	12	25
2	W	247/259 (95%)	230 (93%)	17 (7%)	14	30
2	X	247/259 (95%)	225 (91%)	22 (9%)	9	20
2	Y	247/259 (95%)	227 (92%)	20 (8%)	11	23
All	All	1513/1580 (96%)	1395 (92%)	118 (8%)	11	24

5 of 118 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	X	236	ILE
2	W	150	ILE
2	Y	150	ILE
2	W	139	LEU
2	U	276	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
2	X	297	ASN
2	W	266	HIS
2	Y	259	GLN
2	W	259	GLN
2	W	81	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	2,3	14,14,15	0.61	0	17,19,21	1.03	1 (5%)
3	NAG	C	2	3	14,14,15	0.57	0	17,19,21	0.68	0
3	FUL	C	3	3	10,10,11	0.46	0	14,14,16	0.45	0
4	NAG	D	1	2,4	14,14,15	0.73	0	17,19,21	1.38	2 (11%)
4	NAG	D	2	4	14,14,15	0.71	0	17,19,21	0.77	0
4	MAN	D	3	4	11,11,12	0.65	0	15,15,17	0.61	0
3	NAG	G	1	2,3	14,14,15	0.73	0	17,19,21	1.17	1 (5%)
3	NAG	G	2	3	14,14,15	0.64	0	17,19,21	0.60	0
3	FUL	G	3	3	10,10,11	0.49	0	14,14,16	0.45	0
4	NAG	H	1	2,4	14,14,15	0.81	0	17,19,21	1.64	3 (17%)
4	NAG	H	2	4	14,14,15	0.65	0	17,19,21	0.95	2 (11%)
4	MAN	H	3	4	11,11,12	0.59	0	15,15,17	0.51	0
3	NAG	I	1	2,3	14,14,15	0.77	0	17,19,21	1.11	1 (5%)
3	NAG	I	2	3	14,14,15	0.61	0	17,19,21	0.73	1 (5%)
3	FUL	I	3	3	10,10,11	0.51	0	14,14,16	0.54	0
4	NAG	J	1	2,4	14,14,15	0.79	0	17,19,21	1.35	2 (11%)
4	NAG	J	2	4	14,14,15	0.73	0	17,19,21	0.78	0
4	MAN	J	3	4	11,11,12	0.65	0	15,15,17	0.68	0
3	NAG	K	1	2,3	14,14,15	0.60	0	17,19,21	0.92	0
3	NAG	K	2	3	14,14,15	0.55	0	17,19,21	0.57	0
3	FUL	K	3	3	10,10,11	0.48	0	14,14,16	0.33	0
4	NAG	L	1	2,4	14,14,15	0.83	0	17,19,21	0.57	0
4	NAG	L	2	4	14,14,15	0.65	0	17,19,21	0.82	1 (5%)
4	MAN	L	3	4	11,11,12	0.69	0	15,15,17	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	2,3	-	6/6/23/26	0/1/1/1
3	NAG	C	2	3	-	3/6/23/26	0/1/1/1
3	FUL	C	3	3	-	-	0/1/1/1
4	NAG	D	1	2,4	-	5/6/23/26	0/1/1/1
4	NAG	D	2	4	-	4/6/23/26	0/1/1/1
4	MAN	D	3	4	-	2/2/19/22	0/1/1/1
3	NAG	G	1	2,3	-	4/6/23/26	0/1/1/1
3	NAG	G	2	3	-	6/6/23/26	0/1/1/1
3	FUL	G	3	3	-	-	0/1/1/1
4	NAG	H	1	2,4	-	5/6/23/26	0/1/1/1
4	NAG	H	2	4	-	5/6/23/26	0/1/1/1
4	MAN	H	3	4	-	0/2/19/22	0/1/1/1
3	NAG	I	1	2,3	-	4/6/23/26	0/1/1/1
3	NAG	I	2	3	-	5/6/23/26	0/1/1/1
3	FUL	I	3	3	-	-	0/1/1/1
4	NAG	J	1	2,4	-	4/6/23/26	0/1/1/1
4	NAG	J	2	4	-	4/6/23/26	0/1/1/1
4	MAN	J	3	4	-	1/2/19/22	0/1/1/1
3	NAG	K	1	2,3	-	2/6/23/26	0/1/1/1
3	NAG	K	2	3	-	3/6/23/26	0/1/1/1
3	FUL	K	3	3	-	-	0/1/1/1
4	NAG	L	1	2,4	1/1/5/7	4/6/23/26	0/1/1/1
4	NAG	L	2	4	-	4/6/23/26	0/1/1/1
4	MAN	L	3	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1	NAG	C4-C3-C2	4.95	118.27	111.02
4	J	1	NAG	C4-C3-C2	3.83	116.63	111.02
4	D	1	NAG	C4-C3-C2	3.43	116.05	111.02
4	H	1	NAG	C3-C4-C5	3.05	115.77	110.23
4	J	1	NAG	C3-C4-C5	2.91	115.50	110.23

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	L	1	NAG	C1

5 of 73 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C1-C2-N2-C7
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	2	NAG	C3-C2-N2-C7
3	C	2	NAG	C8-C7-N2-C2


There are no ring outliers.


12 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	2	0
3	I	2	NAG	3	0
4	J	1	NAG	1	0
4	H	2	NAG	1	0
4	L	1	NAG	2	0
4	L	2	NAG	1	0
4	D	2	NAG	1	0
4	L	3	MAN	1	0
3	C	1	NAG	2	0
3	I	3	FUL	1	0
3	C	2	NAG	1	0
4	H	1	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	144/145 (99%)	-0.28	1 (0%) 84 81	42, 68, 170, 188	0
1	B	139/145 (95%)	-0.07	1 (0%) 84 81	39, 76, 187, 194	0
1	E	139/145 (95%)	-0.11	1 (0%) 84 81	39, 69, 184, 200	0
1	F	139/145 (95%)	-0.14	0 100 100	45, 86, 185, 197	0
2	U	275/290 (94%)	0.02	2 (0%) 84 81	44, 85, 168, 187	0
2	W	275/290 (94%)	0.09	2 (0%) 84 81	59, 104, 183, 193	0
2	X	275/290 (94%)	-0.05	1 (0%) 88 86	49, 90, 162, 189	0
2	Y	275/290 (94%)	-0.03	3 (1%) 78 75	46, 81, 180, 191	0
All	All	1661/1740 (95%)	-0.04	11 (0%) 84 81	39, 87, 177, 200	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Y	40	HIS	3.5
2	Y	248	LEU	3.0
2	Y	232	VAL	2.5
1	E	35	GLY	2.5
2	U	291	PHE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	1	14/15	-	-	111,124,133,143	0
3	NAG	C	2	14/15	-	-	156,163,170,171	0
3	FUL	C	3	10/11	-	-	112,119,127,129	0
3	NAG	G	1	14/15	-	-	156,169,179,192	0
3	NAG	G	2	14/15	-	-	185,191,194,196	0
3	FUL	G	3	10/11	-	-	154,171,175,177	0
3	NAG	K	2	14/15	0.52	0.09	185,191,194,196	0
4	NAG	J	2	14/15	0.54	0.10	179,186,194,197	0
4	NAG	L	2	14/15	0.59	0.09	187,195,198,198	0
4	MAN	J	3	11/12	0.65	0.07	191,196,198,198	0
4	MAN	L	3	11/12	0.68	0.08	190,196,198,198	0
3	NAG	K	1	14/15	0.71	0.08	145,156,169,178	0
4	NAG	D	1	14/15	-	-	112,143,156,171	0
4	NAG	D	2	14/15	-	-	183,193,195,196	0
4	MAN	D	3	11/12	-	-	193,197,198,198	0
4	NAG	H	1	14/15	-	-	137,150,160,170	0
4	NAG	H	2	14/15	-	-	180,190,198,198	0
4	MAN	H	3	11/12	-	-	161,171,175,177	0
3	NAG	I	2	14/15	0.75	0.09	170,189,193,194	0
3	FUL	K	3	10/11	0.80	0.12	160,171,172,173	0
3	FUL	I	3	10/11	0.85	0.10	124,131,139,141	0
4	NAG	J	1	14/15	0.86	0.09	151,162,169,171	0
4	NAG	L	1	14/15	0.86	0.07	164,180,187,190	0
3	NAG	I	1	14/15	0.87	0.09	97,118,131,148	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.