



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 11:46 PM UTC

PDB ID : 3O2Q / pdb_00003o2q
Title : Crystal structure of the human symplekin-Ssu72-CTD phosphopeptide complex
Authors : Tong, L.; Xiang, K.
Deposited on : 2010-07-22
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	
1	D	351	
2	B	214	
2	E	214	
3	F	6	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Symplekin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	0	0
			2506	1596	433	464	13			
1	D	311	Total	C	N	O	S	0	0	0
			2461	1569	422	457	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	expression tag	UNP Q92797
A	11	GLY	-	expression tag	UNP Q92797
A	12	SER	-	expression tag	UNP Q92797
A	13	SER	-	expression tag	UNP Q92797
A	14	HIS	-	expression tag	UNP Q92797
A	15	HIS	-	expression tag	UNP Q92797
A	16	HIS	-	expression tag	UNP Q92797
A	17	HIS	-	expression tag	UNP Q92797
A	18	HIS	-	expression tag	UNP Q92797
A	19	HIS	-	expression tag	UNP Q92797
A	20	SER	-	expression tag	UNP Q92797
A	21	SER	-	expression tag	UNP Q92797
A	22	GLY	-	expression tag	UNP Q92797
A	23	LEU	-	expression tag	UNP Q92797
A	24	VAL	-	expression tag	UNP Q92797
A	25	PRO	-	expression tag	UNP Q92797
A	26	ARG	-	expression tag	UNP Q92797
A	27	GLY	-	expression tag	UNP Q92797
A	28	SER	-	expression tag	UNP Q92797
A	29	HIS	-	expression tag	UNP Q92797
D	10	MET	-	expression tag	UNP Q92797
D	11	GLY	-	expression tag	UNP Q92797
D	12	SER	-	expression tag	UNP Q92797
D	13	SER	-	expression tag	UNP Q92797
D	14	HIS	-	expression tag	UNP Q92797

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	15	HIS	-	expression tag	UNP Q92797
D	16	HIS	-	expression tag	UNP Q92797
D	17	HIS	-	expression tag	UNP Q92797
D	18	HIS	-	expression tag	UNP Q92797
D	19	HIS	-	expression tag	UNP Q92797
D	20	SER	-	expression tag	UNP Q92797
D	21	SER	-	expression tag	UNP Q92797
D	22	GLY	-	expression tag	UNP Q92797
D	23	LEU	-	expression tag	UNP Q92797
D	24	VAL	-	expression tag	UNP Q92797
D	25	PRO	-	expression tag	UNP Q92797
D	26	ARG	-	expression tag	UNP Q92797
D	27	GLY	-	expression tag	UNP Q92797
D	28	SER	-	expression tag	UNP Q92797
D	29	HIS	-	expression tag	UNP Q92797

- Molecule 2 is a protein called RNA polymerase II subunit A C-terminal domain phosphatase SSU72.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	189	Total	C	N	O	S	0	0	0
			1546	960	272	303	11			
2	E	192	Total	C	N	O	S	0	0	0
			1565	971	275	308	11			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	expression tag	UNP Q9NP77
B	-18	GLY	-	expression tag	UNP Q9NP77
B	-17	SER	-	expression tag	UNP Q9NP77
B	-16	SER	-	expression tag	UNP Q9NP77
B	-15	HIS	-	expression tag	UNP Q9NP77
B	-14	HIS	-	expression tag	UNP Q9NP77
B	-13	HIS	-	expression tag	UNP Q9NP77
B	-12	HIS	-	expression tag	UNP Q9NP77
B	-11	HIS	-	expression tag	UNP Q9NP77
B	-10	HIS	-	expression tag	UNP Q9NP77
B	-9	SER	-	expression tag	UNP Q9NP77
B	-8	SER	-	expression tag	UNP Q9NP77
B	-7	GLY	-	expression tag	UNP Q9NP77
B	-6	LEU	-	expression tag	UNP Q9NP77

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	VAL	-	expression tag	UNP Q9NP77
B	-4	PRO	-	expression tag	UNP Q9NP77
B	-3	ARG	-	expression tag	UNP Q9NP77
B	-2	GLY	-	expression tag	UNP Q9NP77
B	-1	SER	-	expression tag	UNP Q9NP77
B	0	HIS	-	expression tag	UNP Q9NP77
B	12	SER	CYS	engineered mutation	UNP Q9NP77
E	-19	MET	-	expression tag	UNP Q9NP77
E	-18	GLY	-	expression tag	UNP Q9NP77
E	-17	SER	-	expression tag	UNP Q9NP77
E	-16	SER	-	expression tag	UNP Q9NP77
E	-15	HIS	-	expression tag	UNP Q9NP77
E	-14	HIS	-	expression tag	UNP Q9NP77
E	-13	HIS	-	expression tag	UNP Q9NP77
E	-12	HIS	-	expression tag	UNP Q9NP77
E	-11	HIS	-	expression tag	UNP Q9NP77
E	-10	HIS	-	expression tag	UNP Q9NP77
E	-9	SER	-	expression tag	UNP Q9NP77
E	-8	SER	-	expression tag	UNP Q9NP77
E	-7	GLY	-	expression tag	UNP Q9NP77
E	-6	LEU	-	expression tag	UNP Q9NP77
E	-5	VAL	-	expression tag	UNP Q9NP77
E	-4	PRO	-	expression tag	UNP Q9NP77
E	-3	ARG	-	expression tag	UNP Q9NP77
E	-2	GLY	-	expression tag	UNP Q9NP77
E	-1	SER	-	expression tag	UNP Q9NP77
E	0	HIS	-	expression tag	UNP Q9NP77
E	12	SER	CYS	engineered mutation	UNP Q9NP77

- Molecule 3 is a protein called RNA polymerase II CTD Serine-5 phosphopeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	6	Total	C	N	O	P	0	0	0
			49	29	6	13	1			

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	85	Total	O	0	0
			85	85		
5	B	44	Total	O	0	0
			44	44		
5	D	75	Total	O	0	0
			75	75		
5	E	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Symplekin



K92	K93	R94	P95	E96	Q99	M100	C101	K102	D103	T110	Y116	E121	T130	C131	Q132	H136	V139	D143	H144	H145	Q160	Q163	H164	T166	E166	D167	M170	E171	I172	D173	E174	L175	E180	T186	F187	L188	H189	Y194								
Met	Gly	Ser	Ser	His	His	His	His	His	Ser	Ser	Gly	Leu	Val	Pro	Arg	Gly	Ser	His	Met	Pro	S3	S4	P5	L6	S14	M15	I25	R29	R34	T38	C39	T40	H41	L44	P45	M64	L68	L69	R70	K71	D72	Y76	N79	G80	I81	H84

- Chain F:  67% 17% 17%

P3	T4	S5	P6	S7	Y8
----	----	----	----	----	----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.05Å 97.58Å 105.01Å 90.00° 98.66° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40 30.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-2.40) 99.4 (30.00-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.184 , 0.242 0.183 , 0.241	Depositor DCC
R_{free} test set	2661 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8382	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	0/2544	1.05	1/3449 (0.0%)
1	D	0.91	0/2497	1.06	7/3385 (0.2%)
2	B	0.95	0/1574	1.08	4/2124 (0.2%)
2	E	0.96	0/1594	1.05	0/2152
3	F	0.70	0/40	0.82	0/52
All	All	0.95	0/8249	1.06	12/11162 (0.1%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	92	LYS	CA-C-N	-8.99	110.56	119.64
2	B	92	LYS	C-N-CA	-8.99	110.56	119.64
1	A	245	HIS	CB-CA-C	8.29	122.44	109.42
2	B	94	ARG	N-CA-C	6.02	115.23	108.25
2	B	139	VAL	CB-CA-C	-5.99	101.58	110.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2506	0	2610	44	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2461	0	2566	30	0
2	B	1546	0	1493	44	0
2	E	1565	0	1510	43	0
3	F	49	0	38	3	0
4	B	5	0	0	0	0
5	A	85	0	0	6	0
5	B	44	0	0	3	0
5	D	75	0	0	0	0
5	E	46	0	0	2	0
All	All	8382	0	8217	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60:THR:HG22	2:B:63:GLN:H	1.21	1.05
2:E:165:THR:HG21	2:E:175:LEU:HD11	1.37	1.02
1:A:339:MET:HB3	1:A:340:PRO:HD3	1.36	1.00
2:B:135:HIS:HD2	2:B:189:HIS:HE1	1.03	0.92
2:B:99:GLN:HE21	2:B:99:GLN:H	1.15	0.92

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/351 (90%)	309 (98%)	5 (2%)	1 (0%)	36	50
1	D	309/351 (88%)	305 (99%)	4 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	187/214 (87%)	181 (97%)	6 (3%)	0	100	100
2	E	190/214 (89%)	185 (97%)	3 (2%)	2 (1%)	11	18
3	F	3/6 (50%)	3 (100%)	0	0	100	100
All	All	1004/1136 (88%)	983 (98%)	18 (2%)	3 (0%)	36	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	MET
2	E	5	PRO
2	E	72	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/315 (90%)	265 (94%)	18 (6%)	16	28
1	D	278/315 (88%)	260 (94%)	18 (6%)	15	27
2	B	177/199 (89%)	162 (92%)	15 (8%)	10	16
2	E	180/199 (90%)	166 (92%)	14 (8%)	11	20
3	F	5/5 (100%)	5 (100%)	0	100	100
All	All	923/1033 (89%)	858 (93%)	65 (7%)	14	24

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	101	CYS
2	E	103	ASP
2	B	70	ARG
2	B	68	LEU
2	E	121	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34

such sidechains are listed below:

Mol	Chain	Res	Type
2	E	124	ASN
2	E	128	GLN
2	E	170	ASN
2	B	164	HIS
2	B	135	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SEP	F	5	3	8,9,10	1.55	1 (12%)	7,12,14	1.67	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SEP	F	5	3	-	4/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	5	SEP	P-O1P	3.53	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	5	SEP	O3P-P-OG	3.15	114.88	106.67
3	F	5	SEP	OG-CB-CA	2.90	110.96	108.14

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	5	SEP	CB-OG-P-O1P
3	F	5	SEP	CB-OG-P-O2P
3	F	5	SEP	CB-OG-P-O3P
3	F	5	SEP	N-CA-CB-OG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	5	SEP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	B	195	-	4,4,4	0.91	0	6,6,6	0.90	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/351 (90%)	-0.49	1 (0%) 90 88	18, 32, 52, 66	0
1	D	311/351 (88%)	-0.38	1 (0%) 90 88	18, 35, 63, 77	0
2	B	189/214 (88%)	-0.42	0 100 100	18, 34, 54, 68	0
2	E	192/214 (89%)	-0.37	1 (0%) 87 85	21, 35, 53, 61	0
3	F	5/6 (83%)	0.55	0 100 100	41, 43, 47, 55	0
All	All	1014/1136 (89%)	-0.42	3 (0%) 90 88	18, 34, 57, 77	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	30	MET	2.9
2	E	5	PRO	2.3
1	A	340	PRO	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SEP	F	5	10/11	0.98	0.06	28,36,39,40	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PO4	B	195	5/5	0.99	0.04	24,27,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.