



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:32 PM UTC

PDB ID : 3O2V / pdb_00003o2v
Title : Crystal structure of 1E9 PheL89Ser/LeuH47Trp/MetH100bPhe, an engineered Diels-Alderase Fab with modified specificity and catalytic activity
Authors : Verdino, P.; Wilson, I.A.
Deposited on : 2010-07-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	219	
2	H	227	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TRS	H	239	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chimeric antibody Fab 1E9, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	218	Total	C	N	O	S	0	9	0
			1739	1099	291	344	5			

- Molecule 2 is a protein called Chimeric antibody Fab 1E9, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	219	Total	C	N	O	S	0	3	0
			1676	1069	274	326	7			

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	L	1	Total	C	N	O	0	0
			8	4	1	3		
3	L	1	Total	C	N	O	0	0
			8	4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	L	1	Total	C	N	O	0	0
			8	4	1	3		
3	L	1	Total	C	N	O	0	0
			8	4	1	3		
3	L	1	Total	C	N	O	0	0
			8	4	1	3		
3	L	1	Total	C	N	O	0	0
			8	4	1	3		
3	H	1	Total	C	N	O	0	0
			8	4	1	3		
3	H	1	Total	C	N	O	0	0
			8	4	1	3		
3	H	1	Total	C	N	O	0	0
			8	4	1	3		
3	H	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			13	6	7		
4	H	1	Total	C	O	0	0
			13	6	7		
4	H	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	77	Total 80	O 80	0	3
6	H	85	Total 86	O 86	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chimeric antibody Fab 1E9, light chain



- Molecule 2: Chimeric antibody Fab 1E9, heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.73Å 128.73Å 92.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-2.30) 99.4 (50.00-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.219 0.195 , 0.222	Depositor DCC
R_{free} test set	1973 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.473	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3765	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, SO4, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.10	1/1804 (0.1%)	0.94	2/2444 (0.1%)
2	H	1.07	0/1726	1.00	2/2351 (0.1%)
All	All	1.08	1/3530 (0.0%)	0.97	4/4795 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	27(E)	SER	C-O	6.23	1.31	1.24

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	83	LEU	CA-CB-CG	5.47	135.45	116.30
1	L	66	GLY	CA-C-O	-5.41	118.42	122.37
2	H	125	ALA	CA-C-N	5.14	126.26	119.84
2	H	125	ALA	C-N-CA	5.14	126.26	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1739	0	1706	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1676	0	1644	13	0
3	H	32	0	48	7	0
3	L	48	0	72	7	0
4	H	26	0	10	2	0
4	L	13	0	5	0	0
5	H	30	0	0	1	0
5	L	35	0	0	0	0
6	H	86	0	0	0	0
6	L	80	0	0	1	0
All	All	3765	0	3485	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:54:ARG:NH2	3:L:216:TRS:HN1	2.00	0.60
1:L:66:GLY:HA2	3:L:215:TRS:N	2.17	0.58
1:L:54:ARG:HH21	3:L:216:TRS:HN1	1.50	0.58
2:H:40:ALA:HB3	2:H:43:LYS:HD3	1.86	0.57
1:L:66:GLY:HA2	3:L:215:TRS:HN3	1.71	0.55
2:H:118:GLY:H	3:H:237:TRS:HN1	1.54	0.55
2:H:17:THR:H	3:H:239:TRS:HN2	1.56	0.53
1:L:54:ARG:NH2	3:L:216:TRS:N	2.57	0.52
2:H:187:LEU:C	2:H:187:LEU:HD12	2.34	0.52
2:H:17:THR:H	3:H:239:TRS:HN1	1.59	0.51
4:H:241:FLC:OG1	4:H:241:FLC:OHB	2.28	0.51
1:L:142:ARG:NH2	1:L:163[B]:VAL:HG11	2.26	0.51
1:L:46:PHE:CZ	1:L:49:TYR:HB3	2.47	0.49
2:H:74:SER:H	3:H:236:TRS:HN3	1.61	0.47
2:H:8:GLY:N	4:H:240:FLC:OHB	2.45	0.47
1:L:142:ARG:CZ	1:L:163[B]:VAL:HG11	2.45	0.47
3:L:216:TRS:O3	3:L:216:TRS:O1	2.33	0.46
1:L:83:LEU:HD12	6:L:271:HOH:O	2.16	0.46
1:L:33:LEU:C	1:L:33:LEU:HD13	2.41	0.45
1:L:159:SER:HA	1:L:178:THR:O	2.17	0.45
1:L:158:ASN:OD1	1:L:158:ASN:N	2.50	0.44
2:H:179[B]:GLN:HG2	5:H:246:SO4:O3	2.16	0.44
1:L:145:LYS:HB3	1:L:197:THR:HB	1.99	0.44
2:H:17:THR:N	3:H:239:TRS:N	2.61	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:6:GLN:HA	2:H:21:SER:O	2.18	0.43
2:H:17:THR:N	3:H:239:TRS:HN2	2.15	0.43
1:L:120:PRO:HD3	1:L:132:VAL:HG22	2.00	0.42
2:H:26:GLY:HA2	3:H:238:TRS:HN3	1.85	0.42
1:L:54:ARG:CZ	3:L:216:TRS:N	2.83	0.41
1:L:79:GLU:OE1	1:L:81[A]:GLU:OE2	2.38	0.41
2:H:20:ILE:HD13	2:H:20:ILE:HG21	1.83	0.41
1:L:39:LYS:HB3	1:L:40:PRO:HD2	2.03	0.41
1:L:40:PRO:HB3	1:L:165:GLU:HG3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	225/219 (103%)	218 (97%)	7 (3%)	0	100	100
2	H	220/227 (97%)	213 (97%)	7 (3%)	0	100	100
All	All	445/446 (100%)	431 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	203/195 (104%)	195 (96%)	8 (4%)	28	43
2	H	187/192 (97%)	183 (98%)	4 (2%)	47	66
All	All	390/387 (101%)	378 (97%)	12 (3%)	37	52

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	22	SER
1	L	63	SER
1	L	81[A]	GLU
1	L	81[B]	GLU
1	L	129	THR
1	L	136	LEU
1	L	190	LYS
1	L	213	GLU
2	H	117	LYS
2	H	131	THR
2	H	153	THR
2	H	222	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	147	GLN
2	H	81	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	L	216	-	7,7,7	0.43	0	9,9,9	0.92	0
5	SO4	L	222	-	4,4,4	0.31	0	6,6,6	0.37	0
5	SO4	H	242	-	4,4,4	0.24	0	6,6,6	0.88	0
5	SO4	L	224	-	4,4,4	0.23	0	6,6,6	0.66	0
3	TRS	L	219	-	7,7,7	0.49	0	9,9,9	0.56	0
3	TRS	L	217	-	7,7,7	0.50	0	9,9,9	0.63	0
5	SO4	L	228	-	4,4,4	0.23	0	6,6,6	0.25	0
3	TRS	H	236	-	7,7,7	0.50	0	9,9,9	1.28	2 (22%)
3	TRS	H	239	-	7,7,7	0.62	0	9,9,9	1.37	2 (22%)
4	FLC	H	240	-	12,12,12	1.00	0	17,17,17	2.39	9 (52%)
3	TRS	L	215	-	7,7,7	0.35	0	9,9,9	1.18	1 (11%)
3	TRS	L	220	-	7,7,7	0.64	0	9,9,9	1.19	0
5	SO4	H	243	-	4,4,4	0.36	0	6,6,6	0.54	0
3	TRS	H	237	-	7,7,7	0.59	0	9,9,9	0.90	0
5	SO4	L	227	-	4,4,4	0.32	0	6,6,6	0.19	0
3	TRS	H	238	-	7,7,7	0.44	0	9,9,9	1.34	2 (22%)
4	FLC	H	241	-	12,12,12	1.07	0	17,17,17	1.95	4 (23%)
5	SO4	H	246	-	4,4,4	0.27	0	6,6,6	0.47	0
5	SO4	H	247	-	4,4,4	0.28	0	6,6,6	0.23	0
5	SO4	L	226	-	4,4,4	0.34	0	6,6,6	0.18	0
5	SO4	H	245	-	4,4,4	0.31	0	6,6,6	0.17	0
3	TRS	L	218	-	7,7,7	0.64	0	9,9,9	0.73	0
4	FLC	L	221	-	12,12,12	1.08	0	17,17,17	1.40	1 (5%)
5	SO4	L	223	-	4,4,4	0.27	0	6,6,6	0.11	0
5	SO4	L	225	-	4,4,4	0.23	0	6,6,6	0.32	0
5	SO4	H	244	-	4,4,4	0.22	0	6,6,6	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	L	216	-	-	5/9/9/9	-
4	FLC	L	221	-	-	8/16/16/16	-
3	TRS	H	236	-	-	5/9/9/9	-
3	TRS	H	239	-	-	8/9/9/9	-
4	FLC	H	240	-	-	7/16/16/16	-
3	TRS	L	215	-	-	7/9/9/9	-
3	TRS	L	220	-	-	5/9/9/9	-
3	TRS	L	219	-	-	7/9/9/9	-
3	TRS	H	238	-	-	3/9/9/9	-
3	TRS	H	237	-	-	4/9/9/9	-
3	TRS	L	217	-	-	6/9/9/9	-
3	TRS	L	218	-	-	6/9/9/9	-
4	FLC	H	241	-	-	10/16/16/16	-

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	241	FLC	OHB-CB-CBC	4.60	115.48	108.96
4	L	221	FLC	OB2-CBC-CB	4.54	121.85	113.14
4	H	240	FLC	CB-CA-CAC	-4.20	102.43	113.92
4	H	240	FLC	OB2-CBC-CB	4.07	120.94	113.14
4	H	241	FLC	OB2-CBC-CB	3.72	120.28	113.14
4	H	240	FLC	OHB-CB-CA	-3.39	101.66	109.38
4	H	240	FLC	OB1-CBC-CB	-3.03	116.22	122.09
4	H	240	FLC	CG-CB-CA	3.03	117.09	109.31
3	H	239	TRS	C2-C-N	-2.95	100.65	108.17
4	H	241	FLC	OHB-CB-CA	-2.61	103.43	109.38
4	H	240	FLC	OG1-CGC-CG	-2.55	115.73	122.95
3	H	236	TRS	C3-C-N	2.54	114.64	108.17
4	H	241	FLC	CA-CB-CBC	-2.52	104.47	110.03
3	H	236	TRS	C1-C-N	-2.39	102.08	108.17
4	H	240	FLC	CG-CB-CBC	-2.38	104.77	110.03
4	H	240	FLC	OG2-CGC-CG	2.30	121.62	114.35
3	L	215	TRS	C3-C-N	-2.28	102.34	108.17
3	H	238	TRS	C1-C-N	-2.24	102.45	108.17
3	H	239	TRS	O2-C2-C	-2.20	104.74	110.88
4	H	240	FLC	CB-CG-CGC	2.04	119.50	113.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	238	TRS	O1-C1-C	-2.04	105.19	110.88

There are no chirality outliers.

All (81) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	216	TRS	C1-C-C2-O2
3	L	216	TRS	C3-C-C2-O2
3	L	216	TRS	N-C-C2-O2
3	L	217	TRS	C1-C-C2-O2
3	L	217	TRS	C3-C-C2-O2
3	L	217	TRS	N-C-C2-O2
3	L	218	TRS	C2-C-C1-O1
3	L	218	TRS	C3-C-C1-O1
3	L	218	TRS	N-C-C1-O1
3	L	218	TRS	C1-C-C2-O2
3	L	219	TRS	C1-C-C2-O2
3	L	219	TRS	C1-C-C3-O3
3	L	219	TRS	C2-C-C3-O3
3	L	219	TRS	N-C-C3-O3
3	L	220	TRS	C1-C-C2-O2
3	L	220	TRS	C3-C-C2-O2
3	H	236	TRS	C1-C-C2-O2
3	H	236	TRS	C3-C-C2-O2
3	H	236	TRS	N-C-C2-O2
3	H	236	TRS	N-C-C3-O3
3	H	237	TRS	C1-C-C2-O2
3	H	238	TRS	C3-C-C2-O2
3	H	238	TRS	N-C-C2-O2
3	H	239	TRS	C1-C-C2-O2
4	L	221	FLC	CAC-CA-CB-CBC
4	L	221	FLC	CAC-CA-CB-CG
4	L	221	FLC	CAC-CA-CB-OHB
4	L	221	FLC	CG-CB-CBC-OB1
4	L	221	FLC	CG-CB-CBC-OB2
4	L	221	FLC	OHB-CB-CBC-OB1
4	L	221	FLC	OHB-CB-CBC-OB2
4	H	240	FLC	CAC-CA-CB-OHB
4	H	240	FLC	CA-CB-CBC-OB1
4	H	240	FLC	CA-CB-CBC-OB2
4	H	240	FLC	OHB-CB-CBC-OB1
4	H	240	FLC	OHB-CB-CBC-OB2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	H	241	FLC	CAC-CA-CB-CBC
4	H	241	FLC	CAC-CA-CB-OHB
4	H	241	FLC	CAC-CA-CB-CG
4	H	240	FLC	CAC-CA-CB-CBC
4	H	240	FLC	CAC-CA-CB-CG
4	H	241	FLC	CBC-CB-CG-CGC
3	L	215	TRS	C3-C-C1-O1
3	L	218	TRS	C3-C-C2-O2
3	L	219	TRS	C3-C-C2-O2
3	H	237	TRS	C3-C-C2-O2
3	H	239	TRS	C3-C-C2-O2
4	H	241	FLC	CA-CB-CG-CGC
4	H	241	FLC	OHB-CB-CG-CGC
3	L	215	TRS	N-C-C1-O1
3	L	215	TRS	C1-C-C2-O2
3	L	215	TRS	N-C-C2-O2
3	L	217	TRS	C2-C-C3-O3
3	L	217	TRS	N-C-C3-O3
3	L	220	TRS	C2-C-C1-O1
3	L	220	TRS	N-C-C1-O1
3	L	220	TRS	N-C-C2-O2
3	H	238	TRS	C1-C-C2-O2
3	H	239	TRS	C1-C-C3-O3
3	H	239	TRS	N-C-C3-O3
4	L	221	FLC	CA-CB-CBC-OB2
4	H	241	FLC	CA-CB-CBC-OB1
4	H	241	FLC	CA-CB-CBC-OB2
4	H	241	FLC	CG-CB-CBC-OB2
3	L	215	TRS	C2-C-C1-O1
3	L	216	TRS	C3-C-C1-O1
3	L	217	TRS	C1-C-C3-O3
3	H	236	TRS	C2-C-C3-O3
3	H	237	TRS	C2-C-C1-O1
3	H	239	TRS	C2-C-C3-O3
4	H	241	FLC	CG-CB-CBC-OB1
3	L	215	TRS	C3-C-C2-O2
3	L	215	TRS	C2-C-C3-O3
3	L	216	TRS	N-C-C1-O1
3	L	218	TRS	N-C-C2-O2
3	L	219	TRS	C3-C-C1-O1
3	L	219	TRS	N-C-C2-O2
3	H	237	TRS	N-C-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	H	239	TRS	N-C-C2-O2
3	H	239	TRS	C3-C-C1-O1
3	H	239	TRS	N-C-C1-O1

There are no ring outliers.

9 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	216	TRS	5	0
3	H	236	TRS	1	0
3	H	239	TRS	4	0
4	H	240	FLC	1	0
3	L	215	TRS	2	0
3	H	237	TRS	1	0
3	H	238	TRS	1	0
4	H	241	FLC	1	0
5	H	246	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	L	218/219 (99%)	0.12	8 (3%)	45	48	19, 49, 72, 94	9 (4%)
2	H	219/227 (96%)	0.15	13 (5%)	28	30	24, 46, 84, 128	3 (1%)
All	All	437/446 (97%)	0.14	21 (4%)	35	37	19, 47, 81, 128	12 (2%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	131	THR	6.4
2	H	128	SER	5.8
2	H	130	SER	4.9
2	H	127	SER	4.8
2	H	129	LYS	4.6
2	H	133	GLY	4.0
2	H	227	PRO	3.6
2	H	53	TYR	3.3
2	H	132	SER	3.2
1	L	213	GLU	3.1
1	L	24[A]	ARG	2.7
1	L	152	ASN	2.5
2	H	202	GLY	2.4
1	L	83	LEU	2.4
2	H	205	THR	2.3
2	H	136	GLY	2.3
1	L	156	SER	2.3
1	L	95	PHE	2.3
1	L	1	GLU	2.2
1	L	158	ASN	2.1
2	H	1	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TRS	L	220	8/8	0.55	0.24	82,84,85,85	0
4	FLC	L	221	13/13	0.58	0.20	113,116,117,117	0
3	TRS	L	215	8/8	0.61	0.19	84,84,85,85	0
3	TRS	L	218	8/8	0.61	0.20	82,84,85,85	0
5	SO4	L	223	5/5	0.63	0.14	138,139,139,139	0
5	SO4	H	247	5/5	0.69	0.13	125,125,126,126	0
5	SO4	L	227	5/5	0.71	0.18	123,124,124,125	0
4	FLC	H	241	13/13	0.72	0.17	78,83,85,85	0
5	SO4	H	244	5/5	0.72	0.14	121,122,122,123	0
4	FLC	H	240	13/13	0.72	0.20	97,100,101,101	0
3	TRS	H	239	8/8	0.73	0.21	84,85,85,87	0
3	TRS	L	219	8/8	0.74	0.21	100,100,100,100	0
5	SO4	L	222	5/5	0.74	0.13	103,104,104,104	0
5	SO4	L	226	5/5	0.76	0.17	114,114,115,116	0
5	SO4	L	228	5/5	0.78	0.13	127,127,128,128	0
5	SO4	H	245	5/5	0.79	0.19	134,134,135,136	0
3	TRS	H	238	8/8	0.79	0.25	92,93,94,95	0
3	TRS	L	217	8/8	0.80	0.18	84,84,85,86	0
3	TRS	H	236	8/8	0.82	0.19	73,76,77,79	0
5	SO4	L	224	5/5	0.85	0.10	86,86,88,88	0
3	TRS	H	237	8/8	0.86	0.24	88,88,89,90	0
5	SO4	L	225	5/5	0.87	0.11	93,94,95,96	0
3	TRS	L	216	8/8	0.89	0.13	82,83,83,83	0
5	SO4	H	246	5/5	0.90	0.11	75,76,78,78	0
5	SO4	H	242	5/5	0.95	0.09	44,46,49,51	5
5	SO4	H	243	5/5	0.96	0.11	53,55,59,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.