



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 09:12 AM UTC

PDB ID : 3O2X / pdb_00003o2x
Title : MMP-13 in complex with selective tetrazole core inhibitor
Authors : Kiefer, J.R.; Barta, T.E.; Becker, D.P.; Mathis, K.J.; Williams, J.
Deposited on : 2010-07-22
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	9% 90% 8% .
1	B	164	10% 90% 7% .
1	C	164	10% 92% 7% .
1	D	164	9% 95% . .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6167 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Collagenase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	2	0
			1314	848	213	248	5			
1	B	164	Total	C	N	O	S	0	0	0
			1300	841	211	243	5			
1	C	164	Total	C	N	O	S	0	2	0
			1314	850	213	246	5			
1	D	164	Total	C	N	O	S	0	0	0
			1300	841	211	243	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1104	ALA	-	expression tag	UNP P45452
B	2104	ALA	-	expression tag	UNP P45452
C	1104	ALA	-	expression tag	UNP P45452
D	2104	ALA	-	expression tag	UNP P45452

- Molecule 2 is N-hydroxy-1-(2-methoxyethyl)-4-{[4-(3-{5-[4-(trifluoromethoxy)phenyl]-2H-tetrazol-2-yl}propoxy)phenyl]sulfonyl}piperidine-4-carboxamide (CCD ID: 3O2) (formula: C₂₆H₃₁F₃N₆O₇S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 43	C 26	F 3	N 6	O 7	S 1	0	0
2	B	1	Total 43	C 26	F 3	N 6	O 7	S 1	0	0
2	C	1	Total 43	C 26	F 3	N 6	O 7	S 1	0	0
2	D	1	Total 43	C 26	F 3	N 6	O 7	S 1	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		
3	D	2	Total	Zn	0	0
			2	2		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		
4	C	2	Total	Ca	0	0
			2	2		
4	D	2	Total	Ca	0	0
			2	2		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	2	3	1		
6	B	1	Total	C	N	O	S	0	0
			12	6	2	3	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	173	Total	O	0	0
			173	173		
7	B	151	Total	O	0	0
			151	151		
7	C	180	Total	O	0	0
			180	180		
7	D	178	Total	O	0	0
			178	178		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Collagenase 3




- Molecule 1: Collagenase 3



- Molecule 1: Collagenase 3



- Molecule 1: Collagenase 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.86Å 95.26Å 120.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.72 – 1.90 19.72 – 1.90	Depositor EDS
% Data completeness (in resolution range)	84.1 (19.72-1.90) 84.0 (19.72-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.177 , 0.210 0.192 , 0.225	Depositor DCC
R_{free} test set	2879 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6167	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8373e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN, 3O2, EPE, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1360	0.63	0/1848
1	B	0.45	0/1346	0.68	0/1829
1	C	0.44	0/1360	0.64	0/1848
1	D	0.41	0/1346	0.63	0/1829
All	All	0.43	0/5412	0.64	0/7354

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	2247	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1314	0	1218	11	0
1	B	1300	0	1211	16	0
1	C	1314	0	1225	16	0
1	D	1300	0	1211	9	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0
2	C	43	0	30	1	0
2	D	43	0	30	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	A	15	0	0	0	0
5	B	15	0	0	0	0
5	C	5	0	0	0	0
5	D	10	0	0	0	0
6	A	12	0	12	0	0
6	B	12	0	12	0	0
7	A	173	0	0	0	0
7	B	151	0	0	1	0
7	C	180	0	0	1	0
7	D	178	0	0	2	0
All	All	6167	0	5009	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1124:ASN:HD21	1:C:1167:PHE:H	1.06	0.95
1:D:2124:ASN:HD21	1:D:2167:PHE:H	1.07	0.95
1:A:1218:LEU:HD22	1:A:1246:TYR:HB2	1.56	0.88
1:B:2109:ARG:HB3	1:B:2110:THR:HA	1.63	0.81
1:B:2109:ARG:CB	1:B:2110:THR:HA	2.15	0.75

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/164 (100%)	155 (94%)	7 (4%)	2 (1%)	10	4
1	B	162/164 (99%)	152 (94%)	10 (6%)	0	100	100
1	C	164/164 (100%)	156 (95%)	7 (4%)	1 (1%)	21	13
1	D	162/164 (99%)	154 (95%)	7 (4%)	1 (1%)	21	13
All	All	652/656 (99%)	617 (95%)	31 (5%)	4 (1%)	21	13

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	2249	LYS
1	A	1194	ASN
1	A	1248	GLY
1	C	1109	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/139 (101%)	135 (96%)	5 (4%)	31	23
1	B	138/139 (99%)	133 (96%)	5 (4%)	31	23
1	C	140/139 (101%)	137 (98%)	3 (2%)	47	44
1	D	138/139 (99%)	137 (99%)	1 (1%)	76	78
All	All	556/556 (100%)	542 (98%)	14 (2%)	43	37

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2136	LYS
1	B	2247	THR
1	D	2124	ASN
1	C	1136	LYS
1	C	1182	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1131	HIS
1	C	1251	HIS
1	D	2157	HIS
1	D	2124	ASN
1	C	1124	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 16 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	C	1	-	4,4,4	0.22	0	6,6,6	0.18	0
5	SO4	A	9	-	4,4,4	0.23	0	6,6,6	0.06	0
5	SO4	D	5	-	4,4,4	0.23	0	6,6,6	0.14	0
6	EPE	A	400	-	12,12,15	0.93	1 (8%)	15,16,20	1.48	2 (13%)
2	3O2	C	1801	3	44,46,46	0.93	1 (2%)	57,66,66	1.34	9 (15%)
5	SO4	B	3	-	4,4,4	0.24	0	6,6,6	0.10	0
5	SO4	B	4	-	4,4,4	0.24	0	6,6,6	0.17	0
5	SO4	B	6	-	4,4,4	0.24	0	6,6,6	0.10	0
6	EPE	B	400	-	12,12,15	0.96	1 (8%)	15,16,20	1.47	1 (6%)
2	3O2	B	1801	3	44,46,46	0.94	1 (2%)	57,66,66	1.37	10 (17%)
5	SO4	D	8	-	4,4,4	0.24	0	6,6,6	0.11	0
2	3O2	A	1801	3	44,46,46	0.95	1 (2%)	57,66,66	1.45	11 (19%)
2	3O2	D	1801	3	44,46,46	0.96	2 (4%)	57,66,66	1.25	9 (15%)
5	SO4	A	7	-	4,4,4	0.23	0	6,6,6	0.09	0
5	SO4	A	2	-	4,4,4	0.22	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EPE	A	400	-	-	1/6/14/19	0/1/1/1
2	3O2	C	1801	3	-	9/39/55/55	0/4/4/4
6	EPE	B	400	-	-	1/6/14/19	0/1/1/1
2	3O2	B	1801	3	-	5/39/55/55	0/4/4/4
2	3O2	A	1801	3	-	9/39/55/55	0/4/4/4
2	3O2	D	1801	3	-	8/39/55/55	0/4/4/4

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	400	EPE	C10-S	2.84	1.81	1.77
6	A	400	EPE	C10-S	2.83	1.81	1.77
2	B	1801	3O2	O7-C26	2.29	1.43	1.31
2	D	1801	3O2	O7-C26	2.20	1.43	1.31
2	A	1801	3O2	O7-C26	2.17	1.43	1.31

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1801	3O2	C16-O4-C11	-3.94	107.69	117.93
2	A	1801	3O2	N6-N3-N4	3.84	117.27	114.01
2	A	1801	3O2	C16-O4-C11	-3.62	108.53	117.93
2	B	1801	3O2	C4-C3-N1	-3.53	107.55	111.16
2	C	1801	3O2	N6-N3-N4	3.42	116.91	114.01

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1801	3O2	C4-C1-S1-O3
2	B	1801	3O2	C16-C17-C18-N3
2	C	1801	3O2	C4-C1-S1-O3
2	D	1801	3O2	C4-C1-S1-O3
6	A	400	EPE	C10-C9-N1-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1801	3O2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 3O2 C 1801


The figure displays four panels of structural data for Ligand 3O2 C 1801. The top-left panel shows bond lengths with green lines and magenta highlights for specific bonds. The top-right panel shows bond angles with green lines and magenta highlights for specific angles. The bottom-left panel shows torsions with green lines and magenta highlights for specific torsions. The bottom-right panel shows rings with green lines and magenta highlights for specific rings.


Bond lengths


Bond angles

Torsions

Rings







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9
1	A	164/164 (100%)	0.31	15 (9%)	15	4, 9, 20, 25	2 (1%)
1	B	164/164 (100%)	0.52	16 (9%)	13	4, 9, 22, 27	0
1	C	164/164 (100%)	0.43	17 (10%)	11	4, 9, 17, 24	2 (1%)
1	D	164/164 (100%)	0.46	14 (8%)	16	5, 9, 18, 27	0
All	All	656/656 (100%)	0.43	62 (9%)	14	4, 9, 20, 27	4 (0%)

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2248	GLY	10.0
1	D	2246	TYR	9.9
1	B	2247	THR	9.9
1	D	2249	LYS	9.9
1	B	2249	LYS	8.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EPE	A	400	12/15	0.87	0.14	29,29,30,30	0
2	SO2	B	1801	43/43	0.88	0.14	19,28,50,51	0
5	SO4	A	7	5/5	0.89	0.12	45,46,46,46	0
6	EPE	B	400	12/15	0.89	0.15	37,37,37,37	0
5	SO4	B	4	5/5	0.90	0.13	39,39,39,40	0
5	SO4	B	6	5/5	0.90	0.10	40,41,41,41	0
2	SO2	A	1801	43/43	0.91	0.09	11,15,20,24	0
5	SO4	A	9	5/5	0.92	0.16	47,48,48,48	0
5	SO4	D	8	5/5	0.92	0.12	33,33,34,34	0
5	SO4	C	1	5/5	0.93	0.13	41,41,41,42	0
2	SO2	D	1801	43/43	0.93	0.11	15,22,30,31	0
5	SO4	A	2	5/5	0.93	0.12	47,47,48,48	0
2	SO2	C	1801	43/43	0.93	0.09	12,17,22,26	0
5	SO4	D	5	5/5	0.95	0.09	36,36,36,36	0
5	SO4	B	3	5/5	0.95	0.13	45,45,46,46	0
4	CA	B	2996	1/1	0.97	0.18	34,34,34,34	0
4	CA	A	1996	1/1	0.97	0.15	37,37,37,37	0
4	CA	C	1997	1/1	0.98	0.03	8,8,8,8	0
4	CA	B	2997	1/1	0.98	0.05	11,11,11,11	0
4	CA	C	1996	1/1	0.98	0.20	35,35,35,35	0
3	ZN	C	1999	1/1	0.99	0.04	9,9,9,9	0
3	ZN	D	2998	1/1	0.99	0.10	12,12,12,12	0
3	ZN	D	2999	1/1	0.99	0.11	11,11,11,11	0
4	CA	D	2996	1/1	0.99	0.18	37,37,37,37	0
4	CA	D	2997	1/1	0.99	0.02	7,7,7,7	0
3	ZN	A	1999	1/1	0.99	0.04	8,8,8,8	0
4	CA	A	1997	1/1	0.99	0.03	6,6,6,6	0
3	ZN	B	2999	1/1	0.99	0.11	14,14,14,14	0
3	ZN	A	1998	1/1	1.00	0.06	8,8,8,8	0
3	ZN	C	1998	1/1	1.00	0.06	8,8,8,8	0
3	ZN	B	2998	1/1	1.00	0.11	13,13,13,13	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 3O2 B 1801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 3O2 A 1801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3O2 D 1801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 3O2 C 1801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.