



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:08 PM UTC

PDB ID : 4O23 / pdb_00004o23
Title : Crystal structure of mono-zinc form of succinyl diaminopimelate desuccinylase from *Neisseria meningitidis* MC58
Authors : Nocek, B.; Holz, R.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2013-12-16
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	4% 95% ••
1	B	384	32% 91% 5% •

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinyl-diaminopimelate desuccinylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	376	Total	C	N	O	S	0	0	0
			2859	1798	501	551	9			
1	B	372	Total	C	N	O	S	0	0	0
			2805	1763	488	545	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9JYL2
A	-1	ASN	-	expression tag	UNP Q9JYL2
A	0	ALA	-	expression tag	UNP Q9JYL2
B	-2	SER	-	expression tag	UNP Q9JYL2
B	-1	ASN	-	expression tag	UNP Q9JYL2
B	0	ALA	-	expression tag	UNP Q9JYL2

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	238	Total	O	0	0
			238	238		
4	B	142	Total	O	0	0
			142	142		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinyl-diaminopimelate desuccinylase



- Molecule 1: Succinyl-diaminopimelate desuccinylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.32Å 111.39Å 132.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.25 – 2.09 39.25 – 2.09	Depositor EDS
% Data completeness (in resolution range)	89.0 (39.25-2.09) 88.8 (39.25-2.09)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 2.08Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.211 , 0.241 0.217 , 0.247	Depositor DCC
R_{free} test set	2197 reflections (4.45%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6057	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2916	0.62	0/3962
1	B	0.34	0/2862	0.64	1/3896 (0.0%)
All	All	0.33	0/5778	0.63	1/7858 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	355	ASN	N-CA-C	5.14	116.11	108.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2859	0	2816	6	0
1	B	2805	0	2728	8	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	B	10	0	0	0	0
4	A	238	0	0	1	0
4	B	142	0	0	2	0
All	All	6057	0	5544	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:THR:N	4:A:678:HOH:O	2.37	0.56
1:B:312:GLU:OE1	1:B:374:ARG:NH2	2.40	0.54
1:B:327:SER:OG	1:B:341:GLU:OE1	2.23	0.52
1:B:313:THR:OG1	4:B:613:HOH:O	2.19	0.51
1:B:10:ALA:O	1:B:13:LEU:N	2.48	0.46
1:A:361:ILE:HB	1:A:362:PRO:HD3	1.98	0.46
1:A:3:GLU:OE2	1:A:5:GLN:N	2.48	0.46
1:B:143:THR:OG1	1:B:328:ASP:OD2	2.31	0.46
1:B:313:THR:HG22	1:B:363:LYS:HG3	1.99	0.44
1:A:327:SER:OG	1:A:341:GLU:OE2	2.31	0.44
1:A:141:ASP:HA	1:A:145:LYS:HD2	2.00	0.43
1:B:147:VAL:HG13	1:B:335:ILE:HD12	2.00	0.42
1:A:255:LYS:NZ	4:B:617:HOH:O	2.54	0.41
1:B:7:LEU:HD21	1:B:89:GLU:OE2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/384 (97%)	365 (98%)	9 (2%)	0	100	100
1	B	370/384 (96%)	360 (97%)	10 (3%)	0	100	100
All	All	744/768 (97%)	725 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/313 (96%)	301 (100%)	1 (0%)	86	91
1	B	295/313 (94%)	287 (97%)	8 (3%)	39	45
All	All	597/626 (95%)	588 (98%)	9 (2%)	57	65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	287	GLN
1	B	16	ARG
1	B	32	GLU
1	B	169	ASP
1	B	335	ILE
1	B	357	ARG
1	B	359	ASN
1	B	361	ILE
1	B	369	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	HIS
1	A	207	HIS
1	A	350	HIS
1	B	27	GLN
1	B	207	HIS
1	B	239	ASN
1	B	271	GLN
1	B	274	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	404	-	4,4,4	0.24	0	6,6,6	0.06	0
3	SO4	B	403	-	4,4,4	0.24	0	6,6,6	0.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	376/384 (97%)	0.31	14 (3%)	45 47	15, 30, 51, 72	0
1	B	372/384 (96%)	1.48	124 (33%)	1 1	16, 57, 93, 118	0
All	All	748/768 (97%)	0.89	138 (18%)	3 3	15, 37, 84, 118	0

All (138) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	316	ILE	5.6
1	B	37	ILE	5.4
1	B	88	ALA	4.9
1	B	318	ALA	4.9
1	B	95	TYR	4.9
1	B	60	ALA	4.9
1	B	348	THR	4.8
1	B	92	GLY	4.7
1	B	375	LEU	4.5
1	B	311	ALA	4.3
1	B	376	LEU	4.3
1	B	168	VAL	4.3
1	B	314	CYS	4.2
1	B	367	VAL	4.1
1	B	362	PRO	4.0
1	B	117	VAL	4.0
1	B	347	ALA	3.9
1	B	138	ASP	3.9
1	B	118	ALA	3.9
1	B	85	PHE	3.8
1	B	94	LEU	3.8
1	B	335	ILE	3.8
1	B	368	TYR	3.8
1	B	366	ALA	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	7	LEU	3.7
1	B	171	LEU	3.6
1	B	87	PRO	3.6
1	B	310	ILE	3.6
1	B	306	ALA	3.6
1	B	90	ARG	3.6
1	B	317	GLU	3.6
1	B	80	TRP	3.6
1	B	361	ILE	3.6
1	B	25	ASP	3.5
1	B	139	ALA	3.4
1	B	9	LEU	3.4
1	A	168	VAL	3.4
1	B	315	GLY	3.4
1	B	352	ILE	3.4
1	B	121	PRO	3.3
1	B	308	ALA	3.3
1	B	305	VAL	3.3
1	B	159	TYR	3.2
1	B	300	GLY	3.2
1	B	83	PRO	3.2
1	B	147	VAL	3.2
1	B	116	PHE	3.1
1	A	377	ALA	3.1
1	B	93	ARG	3.1
1	B	374	ARG	3.1
1	B	302	LEU	3.1
1	B	364	LEU	3.1
1	B	62	VAL	3.0
1	B	156	LEU	3.0
1	B	349	ILE	3.0
1	B	123	HIS	3.0
1	B	373	ALA	2.9
1	B	371	ILE	2.9
1	B	89	GLU	2.9
1	A	154	ASP	2.8
1	B	175	ILE	2.8
1	B	61	PRO	2.8
1	B	359	ASN	2.8
1	B	58	THR	2.8
1	B	5	GLN	2.8
1	B	307	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	360	ASP	2.8
1	B	108	CYS	2.8
1	B	356	VAL	2.8
1	B	365	SER	2.7
1	B	169	ASP	2.7
1	B	149	VAL	2.7
1	A	271	GLN	2.7
1	B	309	ALA	2.6
1	B	227	TYR	2.6
1	B	77	VAL	2.6
1	A	136	GLU	2.6
1	B	76	PRO	2.6
1	A	169	ASP	2.6
1	B	59	LYS	2.6
1	B	363	LYS	2.6
1	B	112	ALA	2.6
1	B	345	SER	2.6
1	B	353	ASN	2.5
1	B	351	GLN	2.5
1	B	358	LEU	2.5
1	B	301	LYS	2.5
1	A	22	ASP	2.5
1	B	8	GLU	2.4
1	B	372	LEU	2.4
1	B	26	CYS	2.4
1	B	79	LYS	2.4
1	B	313	THR	2.4
1	B	357	ARG	2.4
1	B	128	ALA	2.4
1	B	299	ALA	2.4
1	B	14	ILE	2.4
1	B	72	VAL	2.4
1	A	290	CYS	2.4
1	A	78	GLU	2.3
1	B	38	GLY	2.3
1	B	170	LYS	2.3
1	A	77	VAL	2.3
1	B	354	GLU	2.3
1	B	148	ASP	2.3
1	B	86	GLU	2.3
1	B	137	GLY	2.3
1	B	126	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	110	VAL	2.3
1	B	334	ALA	2.3
1	B	29	LEU	2.3
1	B	124	GLN	2.2
1	B	166	THR	2.2
1	B	304	ASP	2.2
1	B	163	GLY	2.2
1	B	370	GLY	2.2
1	A	267	ALA	2.2
1	B	312	GLU	2.2
1	B	17	PRO	2.2
1	B	152	ALA	2.2
1	B	140	LEU	2.2
1	B	19	VAL	2.1
1	B	326	THR	2.1
1	B	96	GLY	2.1
1	B	167	ALA	2.1
1	A	138	ASP	2.1
1	B	22	ASP	2.1
1	B	84	PRO	2.1
1	A	191	GLY	2.1
1	B	172	GLY	2.1
1	B	39	PHE	2.1
1	A	23	ASP	2.0
1	B	35	HIS	2.0
1	B	120	HIS	2.0
1	B	105	SER	2.0
1	B	342	LEU	2.0
1	B	285	ASP	2.0
1	B	224	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	403	5/5	0.78	0.16	57,57,57,57	0
3	SO4	B	404	5/5	0.84	0.14	76,76,76,77	0
2	ZN	B	402	1/1	0.90	0.08	34,34,34,34	1
2	ZN	B	401	1/1	0.98	0.04	41,41,41,41	1
2	ZN	A	401	1/1	1.00	0.01	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.