



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:47 AM UTC

PDB ID : 5O23 / pdb_00005o23
Title : Crystal structure of WNK3 kinase domain in a monophosphorylated apo state
Authors : Pinkas, D.M.; Bufton, J.C.; Kupinska, K.; Wang, D.; Sethi, R.; Newman, J.A.; Chalk, R.; Berridge, G.; Burgess-Brown, N.A.; von Delft, F.; Arrow-smith, C.H.; Edwards, A.M.; Bountra, C.; Bullock, A.; Structural Genomics Consortium (SGC)
Deposited on : 2017-05-19
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	7% 87% 8% 5%
1	B	285	13% 91% 7% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase WNK3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	P	S	0	2	0
			2147	1369	363	398	1	16			
1	B	279	Total	C	N	O	P	S	0	1	0
			2219	1416	372	412	1	18			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	130	SER	-	expression tag	UNP Q9BYP7
A	131	MET	-	expression tag	UNP Q9BYP7
B	130	SER	-	expression tag	UNP Q9BYP7
B	131	MET	-	expression tag	UNP Q9BYP7

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	A	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	101	Total 101	O 101	0	0
4	B	123	Total 123	O 123	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase WNK3



- Molecule 1: Serine/threonine-protein kinase WNK3



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	124.02Å 124.02Å 107.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	75.93 – 2.25 75.93 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.93-2.25) 99.9 (75.93-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.25Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.205 , 0.239 (Not available) , 0.234	Depositor DCC
R_{free} test set	2140 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4727	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/2182	0.26	0/2947
1	B	0.09	0/2254	0.29	0/3040
All	All	0.09	0/4436	0.28	0/5987

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2147	0	2115	13	0
1	B	2219	0	2192	13	0
2	A	10	0	0	0	0
2	B	15	0	0	2	0
3	A	48	0	72	4	0
3	B	64	0	96	4	0
4	A	101	0	0	0	0
4	B	123	0	0	1	0
All	All	4727	0	4475	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:HIS:HD1	3:A:509:EDO:HO1	1.28	0.79
1:B:131:MET:HE1	1:B:163:LYS:HG2	1.88	0.56
1:A:299:THR:HB	3:A:512:EDO:H12	1.89	0.54
1:B:359:ARG:NH2	2:B:501:SO4:O4	2.38	0.54
1:B:194:GLU:OE1	3:B:507:EDO:O2	2.29	0.49
1:B:208:ARG:HB3	1:B:228:GLU:HB2	1.96	0.48
1:A:139:ALA:HB3	1:A:148:LYS:HB3	1.96	0.48
1:B:153:LEU:HD21	1:B:163:LYS:HB2	1.97	0.47
1:B:313:PRO:HD2	3:B:511:EDO:H21	1.96	0.46
1:B:238:TYR:HB2	1:B:283:ILE:HD11	1.97	0.46
1:A:327:GLU:H	1:A:327:GLU:CD	2.24	0.46
1:B:190:ARG:NH2	4:B:601:HOH:O	2.21	0.46
1:A:275:ASP:HA	3:A:513:EDO:H11	1.98	0.45
1:B:338:MET:HA	1:B:341:MET:HE3	1.99	0.45
1:B:276:LEU:H	3:B:506:EDO:H12	1.82	0.44
1:B:327:GLU:H	1:B:327:GLU:CD	2.26	0.44
1:B:137:ALA:HB1	1:B:147:LEU:HG	2.00	0.44
1:A:389:ASN:OD1	1:A:391[B]:SER:OG	2.35	0.43
1:B:389:ASN:OD1	1:B:391:SER:OG	2.35	0.43
1:A:364:GLY:HA2	3:A:506:EDO:H21	2.00	0.43
2:B:503:SO4:O1	3:B:510:EDO:O2	2.35	0.43
1:A:375:ASP:HB3	1:A:378:VAL:HB	2.02	0.42
1:A:284:THR:HG21	1:A:291:LYS:HE3	2.02	0.41
1:A:383:GLU:O	1:A:387:ARG:HG3	2.20	0.41
1:A:145:ARG:HG3	1:A:168:GLU:HG3	2.03	0.41
1:A:236:LYS:HE2	1:A:240:LYS:HD2	2.03	0.40
1:A:194:GLU:OE1	1:A:302:ARG:NH2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/285 (94%)	262 (97%)	7 (3%)	0	100	100
1	B	277/285 (97%)	268 (97%)	9 (3%)	0	100	100
All	All	546/570 (96%)	530 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/250 (92%)	227 (99%)	2 (1%)	70	79
1	B	235/250 (94%)	233 (99%)	2 (1%)	70	79
All	All	464/500 (93%)	460 (99%)	4 (1%)	70	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	136	LYS
1	A	302	ARG
1	B	131	MET
1	B	394	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	401	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	B	304	1	8,9,10	1.62	1 (12%)	7,12,14	1.45	1 (14%)
1	SEP	A	304	1	8,9,10	1.62	1 (12%)	7,12,14	1.35	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	304	1	-	5/6/8/10	-
1	SEP	A	304	1	-	3/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	304	SEP	P-O1P	3.55	1.61	1.50
1	B	304	SEP	P-O1P	3.53	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	304	SEP	OG-CB-CA	3.28	111.33	108.14
1	A	304	SEP	OG-CB-CA	2.96	111.03	108.14

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	304	SEP	CB-OG-P-O2P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	304	SEP	CB-OG-P-O3P
1	B	304	SEP	CB-OG-P-O2P
1	B	304	SEP	CB-OG-P-O3P
1	B	304	SEP	CB-OG-P-O1P
1	B	304	SEP	CA-CB-OG-P
1	B	304	SEP	N-CA-CB-OG
1	A	304	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

33 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	510	-	3,3,3	0.43	0	2,2,2	0.39	0
3	EDO	B	505	-	3,3,3	0.42	0	2,2,2	0.36	0
2	SO4	B	501	-	4,4,4	0.24	0	6,6,6	0.08	0
3	EDO	B	511	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	A	505	-	3,3,3	0.42	0	2,2,2	0.39	0
3	EDO	B	507	-	3,3,3	0.43	0	2,2,2	0.40	0
3	EDO	B	518	-	3,3,3	0.42	0	2,2,2	0.40	0
3	EDO	B	515	-	3,3,3	0.43	0	2,2,2	0.39	0
2	SO4	B	503	-	4,4,4	0.23	0	6,6,6	0.08	0
3	EDO	B	514	-	3,3,3	0.43	0	2,2,2	0.39	0
3	EDO	B	510	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	A	514	-	3,3,3	0.42	0	2,2,2	0.38	0
3	EDO	B	508	-	3,3,3	0.43	0	2,2,2	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	B	516	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	B	517	-	3,3,3	0.41	0	2,2,2	0.43	0
3	EDO	A	506	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	B	512	-	3,3,3	0.42	0	2,2,2	0.38	0
3	EDO	B	506	-	3,3,3	0.43	0	2,2,2	0.39	0
2	SO4	A	502	-	4,4,4	0.24	0	6,6,6	0.08	0
3	EDO	A	508	-	3,3,3	0.42	0	2,2,2	0.42	0
3	EDO	A	509	-	3,3,3	0.41	0	2,2,2	0.47	0
3	EDO	A	507	-	3,3,3	0.43	0	2,2,2	0.35	0
3	EDO	B	519	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	B	504	-	3,3,3	0.42	0	2,2,2	0.38	0
3	EDO	B	509	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	A	503	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	A	512	-	3,3,3	0.43	0	2,2,2	0.38	0
3	EDO	B	513	-	3,3,3	0.42	0	2,2,2	0.39	0
3	EDO	A	504	-	3,3,3	0.42	0	2,2,2	0.39	0
3	EDO	A	511	-	3,3,3	0.42	0	2,2,2	0.37	0
2	SO4	B	502	-	4,4,4	0.23	0	6,6,6	0.08	0
3	EDO	A	513	-	3,3,3	0.43	0	2,2,2	0.38	0
2	SO4	A	501	-	4,4,4	0.23	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	510	-	-	0/1/1/1	-
3	EDO	B	505	-	-	0/1/1/1	-
3	EDO	B	511	-	-	0/1/1/1	-
3	EDO	B	507	-	-	0/1/1/1	-
3	EDO	A	505	-	-	0/1/1/1	-
3	EDO	B	518	-	-	0/1/1/1	-
3	EDO	B	515	-	-	0/1/1/1	-
3	EDO	B	514	-	-	0/1/1/1	-
3	EDO	B	510	-	-	0/1/1/1	-
3	EDO	A	514	-	-	0/1/1/1	-
3	EDO	B	508	-	-	1/1/1/1	-
3	EDO	B	516	-	-	0/1/1/1	-
3	EDO	B	517	-	-	0/1/1/1	-
3	EDO	A	506	-	-	0/1/1/1	-
3	EDO	B	512	-	-	0/1/1/1	-
3	EDO	B	506	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	508	-	-	0/1/1/1	-
3	EDO	A	509	-	-	0/1/1/1	-
3	EDO	A	507	-	-	0/1/1/1	-
3	EDO	B	519	-	-	0/1/1/1	-
3	EDO	B	504	-	-	0/1/1/1	-
3	EDO	B	509	-	-	0/1/1/1	-
3	EDO	A	503	-	-	0/1/1/1	-
3	EDO	A	512	-	-	0/1/1/1	-
3	EDO	B	513	-	-	0/1/1/1	-
3	EDO	A	504	-	-	0/1/1/1	-
3	EDO	A	511	-	-	0/1/1/1	-
3	EDO	A	513	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	508	EDO	O1-C1-C2-O2

There are no ring outliers.

10 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	SO4	1	0
3	B	511	EDO	1	0
3	B	507	EDO	1	0
2	B	503	SO4	1	0
3	B	510	EDO	1	0
3	A	506	EDO	1	0
3	B	506	EDO	1	0
3	A	509	EDO	1	0
3	A	512	EDO	1	0
3	A	513	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/285 (94%)	0.66	20 (7%) 20 19	30, 59, 90, 117	2 (0%)
1	B	278/285 (97%)	0.78	36 (12%) 7 7	28, 56, 95, 116	1 (0%)
All	All	547/570 (95%)	0.72	56 (10%) 12 11	28, 57, 92, 117	3 (0%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	286	PRO	4.7
1	A	285	GLY	4.5
1	A	217	LEU	4.3
1	B	406	ALA	4.1
1	B	216	ILE	4.1
1	B	170	TRP	4.1
1	B	408	ASP	3.8
1	B	407	GLU	3.8
1	A	216	ILE	3.7
1	B	346	TYR	3.7
1	B	219	GLY	3.6
1	B	242	PHE	3.6
1	B	249	VAL	3.4
1	B	130	SER	3.4
1	B	218	LYS	3.4
1	A	136	LYS	3.2
1	B	268	THR	3.1
1	B	287	THR	3.1
1	A	286	PRO	3.0
1	B	239	LEU	2.9
1	A	220	LYS	2.9
1	A	284	THR	2.8
1	B	283	ILE	2.8
1	B	285	GLY	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	244	VAL	2.7
1	B	290	VAL	2.7
1	A	221	LYS	2.6
1	A	391[A]	SER	2.6
1	B	200	GLY	2.5
1	B	284	THR	2.4
1	B	171	VAL	2.4
1	B	135	MET	2.4
1	B	238	TYR	2.4
1	B	397	ARG	2.4
1	A	368	ALA	2.4
1	B	279	ASP	2.3
1	A	403	ALA	2.3
1	B	245	MET	2.3
1	B	217	LEU	2.3
1	A	364	GLY	2.3
1	A	352	GLN	2.3
1	B	201	LEU	2.2
1	A	268	THR	2.2
1	B	244	VAL	2.2
1	B	289	SER	2.2
1	B	288	GLY	2.2
1	B	255[A]	ARG	2.2
1	A	225	LEU	2.1
1	A	218	LYS	2.1
1	A	219	GLY	2.1
1	B	233	GLY	2.1
1	A	287	THR	2.1
1	B	190	ARG	2.1
1	A	346	TYR	2.0
1	B	276	LEU	2.0
1	B	247	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	304	10/11	0.89	0.13	38,54,92,101	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	SEP	B	304	10/11	0.90	0.11	44,51,91,94	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	511	4/4	0.63	0.21	66,76,78,95	0
3	EDO	B	504	4/4	0.71	0.30	113,119,124,126	0
3	EDO	A	509	4/4	0.75	0.22	56,64,67,70	0
3	EDO	A	510	4/4	0.76	0.23	74,75,85,92	0
3	EDO	B	519	4/4	0.78	0.18	82,85,85,91	0
3	EDO	B	512	4/4	0.79	0.24	83,88,88,103	0
2	SO4	B	502	5/5	0.79	0.09	87,100,117,127	0
3	EDO	B	516	4/4	0.80	0.23	71,72,77,86	0
3	EDO	B	514	4/4	0.83	0.24	62,64,69,78	0
3	EDO	A	505	4/4	0.84	0.22	52,63,66,81	0
3	EDO	B	510	4/4	0.84	0.21	76,78,78,78	0
2	SO4	B	501	5/5	0.84	0.12	83,96,107,124	0
3	EDO	B	517	4/4	0.85	0.22	68,75,86,87	0
3	EDO	B	518	4/4	0.86	0.28	53,61,62,82	0
3	EDO	A	506	4/4	0.87	0.19	68,75,86,94	0
3	EDO	B	505	4/4	0.87	0.14	57,57,74,83	0
3	EDO	B	507	4/4	0.87	0.23	54,77,78,90	0
3	EDO	B	515	4/4	0.87	0.23	64,67,79,79	0
3	EDO	B	511	4/4	0.88	0.19	71,78,82,96	0
3	EDO	B	508	4/4	0.88	0.22	68,69,71,75	0
3	EDO	B	506	4/4	0.89	0.20	46,61,64,82	0
3	EDO	A	508	4/4	0.90	0.18	59,60,68,76	0
3	EDO	A	514	4/4	0.90	0.21	64,70,79,90	0
2	SO4	A	501	5/5	0.91	0.08	71,72,100,114	0
3	EDO	A	507	4/4	0.91	0.15	52,53,62,67	0
3	EDO	B	509	4/4	0.91	0.15	66,68,81,88	0
3	EDO	A	512	4/4	0.92	0.16	52,62,63,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	513	4/4	0.92	0.15	71,72,73,79	0
3	EDO	A	503	4/4	0.92	0.14	58,65,65,71	0
3	EDO	A	504	4/4	0.94	0.12	59,62,65,75	0
2	SO4	A	502	5/5	0.94	0.09	40,68,77,85	0
3	EDO	B	513	4/4	0.95	0.16	42,50,52,60	0
2	SO4	B	503	5/5	0.96	0.10	50,55,72,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.