



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:46 AM UTC

PDB ID : 1O37 / pdb_00001o37
Title : Elaborate Manifold of Short Hydrogen Bond Arrays Mediating Binding of Active Site-Directed Serine Protease Inhibitors
Authors : Katz, B.A.; Elrod, K.; Verner, E.; Mackman, R.L.; Luong, C.; Shrader, W.D.; Sendzik, M.; Spencer, J.R.; Sprengeler, P.A.; Kolesnikov, A.; Tai, V.W.; Hui, H.C.; Breitenbucher, J.G.; Allen, D.; Janc, J.W.
Deposited on : 2003-03-06
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	17% 69% 30% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3810 atoms, of which 1966 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-TRYPSIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	223	Total	C	H	N	O	S	0	16	0
			3360	1047	1678	284	337	14			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 5-(2-AMINOETHYL)-3-{5-[AMINO(IMINIO)METHYL]-1H-BENZIMIDAZO L-2-YL}-1,1'-BIPHENYL-2-OLATE (CCD ID: 653) (formula: C₂₂H₂₂N₅O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			50	22	22	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	132	Total 399	H 266	O 133	0	2

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-TRYPSIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.09Å 62.18Å 70.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.45 7.00 – 1.45	Depositor EDS
% Data completeness (in resolution range)	84.2 (7.00-1.45) 83.2 (7.00-1.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.40 (at 1.45Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.225 , 0.241 0.228 , 0.220	Depositor DCC
R_{free} test set	4294 reflections (10.18%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 67.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3810	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, 653

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.06	36/1773 (2.0%)	1.81	39/2399 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	118	VAL	CA-CB	8.42	1.65	1.54
1	A	73	ILE	CA-CB	7.44	1.64	1.54
1	A	91	HIS	CD2-NE2	-7.37	1.29	1.37
1	A	176	ILE	CA-CB	7.31	1.62	1.54
1	A	40	HIS	CB-CG	7.27	1.60	1.50
1	A	54	SER	CA-CB	7.10	1.64	1.54
1	A	76	VAL	CA-C	7.09	1.61	1.52
1	A	64	GLN	CA-C	7.05	1.60	1.52
1	A	180	MET	CA-C	7.03	1.61	1.52
1	A	82	PHE	N-CA	6.92	1.54	1.46
1	A	40	HIS	CD2-NE2	-6.88	1.30	1.37
1	A	76	VAL	CA-CB	6.82	1.62	1.54
1	A	19	GLY	N-CA	6.71	1.51	1.44
1	A	148	GLY	N-CA	6.39	1.50	1.45
1	A	57	HIS	CG-ND1	-6.34	1.31	1.38
1	A	91	HIS	CA-C	6.18	1.60	1.53
1	A	33	LEU	N-CA	6.17	1.53	1.46
1	A	87[A]	LYS	N-CA	5.91	1.53	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	87[B]	LYS	N-CA	5.91	1.53	1.45
1	A	200	VAL	CA-CB	5.59	1.60	1.54
1	A	42	CYS	N-CA	5.46	1.52	1.45
1	A	73	ILE	N-CA	5.45	1.53	1.46
1	A	53[A]	VAL	CA-CB	5.45	1.61	1.54
1	A	53[B]	VAL	CA-CB	5.45	1.61	1.54
1	A	145	LYS	N-CA	5.38	1.52	1.46
1	A	91	HIS	CG-ND1	-5.38	1.32	1.38
1	A	240	GLN	CA-CB	5.37	1.61	1.53
1	A	78	GLY	CA-C	5.32	1.59	1.52
1	A	231	VAL	CA-CB	5.25	1.60	1.54
1	A	57	HIS	CD2-NE2	-5.24	1.32	1.37
1	A	179	ASN	N-CA	5.23	1.52	1.46
1	A	234	TYR	CA-CB	5.21	1.61	1.53
1	A	99	LEU	CA-CB	5.21	1.61	1.53
1	A	89	ILE	CA-CB	5.13	1.60	1.53
1	A	167	SER	CA-C	5.12	1.59	1.52
1	A	40	HIS	N-CA	5.08	1.52	1.46

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76	VAL	N-CA-C	-8.09	96.93	108.58
1	A	210	GLN	OE1-CD-NE2	-8.06	114.53	122.60
1	A	199	VAL	N-CA-C	-7.72	94.27	107.24
1	A	148	GLY	CA-C-O	-7.29	117.05	122.37
1	A	190	SER	N-CA-C	-7.07	99.40	110.36
1	A	22	CYS	N-CA-C	6.96	118.87	111.28
1	A	79	ASN	CA-CB-CG	-6.82	105.78	112.60
1	A	57	HIS	CA-CB-CG	6.73	120.53	113.80
1	A	153	ASP	N-CA-C	-6.68	104.77	113.12
1	A	231	VAL	N-CA-C	6.67	118.22	110.62
1	A	71	ASP	CA-CB-CG	-6.37	106.23	112.60
1	A	75	VAL	CA-C-O	-6.28	113.70	120.36
1	A	37	SER	N-CA-C	-6.19	103.01	110.44
1	A	162	ILE	N-CA-C	-5.91	101.17	109.21
1	A	148	GLY	N-CA-C	-5.90	106.12	112.08
1	A	237	TRP	NE1-CE2-CZ2	5.77	138.76	130.10
1	A	133	GLY	N-CA-C	-5.76	107.73	115.21
1	A	95	ASN	OD1-CG-ND2	-5.72	116.88	122.60
1	A	86[A]	SER	N-CA-C	-5.72	104.43	111.40
1	A	86[B]	SER	N-CA-C	-5.72	104.43	111.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	181	PHE	CA-CB-CG	5.55	119.35	113.80
1	A	118	VAL	N-CA-C	-5.48	99.61	107.78
1	A	79	ASN	CB-CA-C	-5.45	104.43	109.83
1	A	235	VAL	N-CA-C	5.44	115.64	110.42
1	A	141	TRP	CG-CD1-NE1	-5.43	103.14	110.20
1	A	113[A]	SER	N-CA-C	-5.41	99.91	108.73
1	A	113[B]	SER	N-CA-C	-5.41	99.91	108.73
1	A	237	TRP	NE1-CE2-CD2	-5.25	100.57	107.40
1	A	154	VAL	O-C-N	5.22	128.63	122.36
1	A	34	ASN	N-CA-C	5.22	117.41	108.90
1	A	211	GLY	O-C-N	5.22	128.45	123.65
1	A	215	TRP	NE1-CE2-CZ2	5.21	137.92	130.10
1	A	50	GLN	CA-CB-CG	-5.15	103.80	114.10
1	A	33	LEU	N-CA-C	-5.14	100.35	108.73
1	A	115	ASN	OD1-CG-ND2	-5.13	117.47	122.60
1	A	215	TRP	CG-CD1-NE1	-5.10	103.57	110.20
1	A	169	LYS	CA-CB-CG	-5.06	103.98	114.10
1	A	100	ASN	N-CA-C	-5.02	102.85	110.28
1	A	141	TRP	CD1-NE1-CE2	5.02	117.93	108.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1682	1678	1643	15	0
2	A	1	0	0	0	0
3	A	28	22	22	3	0
4	A	133	266	0	2	0
All	All	1844	1966	1665	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195[A]:SER:HB2	4:A:383[A]:HOH:O	1.90	0.72
1:A:104:MET:HE3	1:A:106:ILE:HD11	1.85	0.58
1:A:31:VAL:HG12	1:A:66[A]:LEU:CD2	2.39	0.53
1:A:195[A]:SER:CB	4:A:383[A]:HOH:O	2.54	0.53
1:A:199:VAL:HG21	1:A:228:TYR:CD1	2.44	0.52
1:A:47:ILE:HD13	1:A:53[B]:VAL:CG2	2.39	0.52
1:A:192:GLN:NE2	3:A:246:653:C3'	2.75	0.49
1:A:188(A):LYS:O	1:A:189:ASP:HB2	2.17	0.44
1:A:192:GLN:HE21	3:A:246:653:C4'	2.31	0.44
1:A:48:ASN:ND2	1:A:51:TRP:HB2	2.33	0.43
1:A:31:VAL:CG1	1:A:66[A]:LEU:CD2	2.97	0.42
1:A:45:SER:OG	1:A:198:PRO:HB3	2.19	0.42
1:A:31:VAL:HG12	1:A:66[A]:LEU:HD23	2.01	0.42
1:A:192:GLN:NE2	3:A:246:653:CV'	2.83	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/223 (106%)	229 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	200/184 (109%)	196 (98%)	4 (2%)	48 16

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	VAL
1	A	125	THR
1	A	236[A]	SER
1	A	236[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	192	GLN
1	A	223	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	653	A	246	-	31,31,31	2.24	8 (25%)	41,44,44	2.43	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	653	A	246	-	-	5/15/15/15	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	246	653	C5'-C1B	-5.62	1.39	1.49
3	A	246	653	C4-N3	-5.05	1.29	1.38
3	A	246	653	C4-C5	-4.60	1.33	1.40
3	A	246	653	C5-N4	-3.92	1.31	1.39
3	A	246	653	C8-N3	-3.90	1.28	1.36
3	A	246	653	CV'-C3'	-3.14	1.42	1.51
3	A	246	653	C6-C1	3.11	1.44	1.39
3	A	246	653	C1'-C8	-2.50	1.43	1.47

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	246	653	CV'-C3'-C4'	-8.63	106.04	120.54
3	A	246	653	CV'-C3'-C2'	7.77	133.60	120.54
3	A	246	653	C6'-C1'-C8	4.27	124.66	119.55
3	A	246	653	C5-N4-C8	-4.12	100.36	105.17
3	A	246	653	C2'-C1'-C6'	-3.18	115.68	119.39
3	A	246	653	C3-C2-C1	2.89	123.89	120.80
3	A	246	653	C6-C1-C7	2.51	124.62	120.25
3	A	246	653	C2-C1-C6	-2.44	116.43	119.25
3	A	246	653	C1'-C6'-C5'	2.41	122.95	119.79

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	246	653	C3'-CV'-CW'-NX'
3	A	246	653	C2-C1-C7-N1
3	A	246	653	C6-C1-C7-N1
3	A	246	653	C2-C1-C7-N2
3	A	246	653	C6-C1-C7-N2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	246	653	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/223 (100%)	1.28	38 (17%) 4 4	7, 19, 30, 51	31 (13%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	115	ASN	4.2
1	A	149	THR	4.0
1	A	193	GLY	3.9
1	A	147	SER	3.5
1	A	151	TYR	3.4
1	A	203	GLY	3.4
1	A	222	LYS	3.4
1	A	125	THR	3.2
1	A	117	ARG	3.2
1	A	146	SER	3.2
1	A	111	ALA	3.1
1	A	128	CYS	3.0
1	A	187	GLY	3.0
1	A	185	LEU	3.0
1	A	116	SER	2.9
1	A	217[A]	SER	2.8
1	A	31	VAL	2.8
1	A	220	CYS	2.8
1	A	62	GLY	2.7
1	A	148	GLY	2.6
1	A	188(A)	LYS	2.6
1	A	97	ASN	2.5
1	A	243	ALA	2.5
1	A	76	VAL	2.5
1	A	51	TRP	2.5
1	A	237	TRP	2.5
1	A	158	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	THR	2.4
1	A	127	SER	2.4
1	A	132	ALA	2.3
1	A	162	ILE	2.3
1	A	219	GLY	2.2
1	A	192	GLN	2.2
1	A	16	ILE	2.1
1	A	150	SER	2.1
1	A	202	SER	2.1
1	A	85	ALA	2.0
1	A	168	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	653	A	246	28/28	0.72	0.14	22,26,36,37	7
2	CA	A	247	1/1	0.80	0.16	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 653 A 246:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.