



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:33 AM UTC

PDB ID : 2O34 / pdb_00002o34
Title : Crystal structure of protein DVU1097 from *Desulfovibrio vulgaris* Hildenborough, Pfam DUF375
Authors : Malashkevich, V.N.; Toro, R.; Sauder, J.M.; Schwinn, K.D.; Thompson, D.A.; Rutter, M.E.; Dickey, M.; Groshong, C.; Bain, K.T.; Adams, J.M.; Reyes, C.; Rooney, I.; Powell, A.; Boice, A.; Gheyi, T.; Ozyurt, S.; Atwell, S.; Wasserman, S.R.; Emtage, S.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2006-11-30
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 10% 86% 13% .
1	B	250	 21% 75% 13% . 8%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3943 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	Se	0	1	0
			1858	1161	338	347	3	9			
1	B	231	Total	C	N	O	S	Se	0	1	0
			1708	1069	308	322	2	7			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	cloning artifact	UNP Q72D34
A	1	LEU	-	cloning artifact	UNP Q72D34
A	3	MSE	MET	modified residue	UNP Q72D34
A	48	MSE	MET	modified residue	UNP Q72D34
A	63	MSE	MET	modified residue	UNP Q72D34
A	89	MSE	MET	modified residue	UNP Q72D34
A	111	MSE	MET	modified residue	UNP Q72D34
A	152	MSE	MET	modified residue	UNP Q72D34
A	205	MSE	MET	modified residue	UNP Q72D34
A	244	MSE	MET	modified residue	UNP Q72D34
B	0	SER	-	cloning artifact	UNP Q72D34
B	1	LEU	-	cloning artifact	UNP Q72D34
B	3	MSE	MET	modified residue	UNP Q72D34
B	48	MSE	MET	modified residue	UNP Q72D34
B	63	MSE	MET	modified residue	UNP Q72D34
B	89	MSE	MET	modified residue	UNP Q72D34
B	111	MSE	MET	modified residue	UNP Q72D34
B	152	MSE	MET	modified residue	UNP Q72D34
B	205	MSE	MET	modified residue	UNP Q72D34
B	244	MSE	MET	modified residue	UNP Q72D34

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Na 2	0	0
2	B	1	Total 1	Na 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	226	Total 226	O 226	0	0
3	B	148	Total 148	O 148	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hypothetical protein



• Molecule 1: Hypothetical protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.48Å 81.48Å 151.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.98 – 1.95 19.98 – 1.95	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.98-1.95) 60.3 (19.98-1.95)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 1.96Å)	Xtriage
Refinement program	REFMAC 5.2.0019, SHELX	Depositor
R, R_{free}	0.210 , 0.273 0.212 , 0.281	Depositor DCC
R_{free} test set	1717 reflections (3.13%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3943	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1884	1.01	1/2547 (0.0%)
1	B	0.80	0/1730	0.99	2/2339 (0.1%)
All	All	0.83	0/3614	1.00	3/4886 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	ASP	CA-C-N	6.73	128.26	119.84
1	B	146	ASP	C-N-CA	6.73	128.26	119.84
1	A	143	ILE	CB-CA-C	-5.89	104.85	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1858	0	1861	28	0
1	B	1708	0	1716	31	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	226	0	0	4	0
3	B	148	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3943	0	3577	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:GLY:H	1:A:175:HIS:HA	0.94	1.08
1:B:60:LYS:HA	1:B:63:MSE:HE2	1.40	1.01
1:A:174:GLY:N	1:A:175:HIS:HA	1.70	0.95
1:B:146:ASP:N	1:B:147:PRO:HD2	1.85	0.92
1:B:209:ALA:HB1	1:B:237[B]:ARG:HG3	1.59	0.83
1:B:207:ARG:HD2	3:B:639:HOH:O	1.79	0.81
1:B:146:ASP:H	1:B:147:PRO:CD	1.94	0.80
1:B:144:LEU:O	1:B:152:MSE:HB2	1.83	0.77
1:A:174:GLY:H	1:A:175:HIS:CA	1.86	0.76
1:B:146:ASP:N	1:B:147:PRO:CD	2.48	0.75
1:B:147:PRO:HA	1:B:149:SER:H	1.48	0.75
1:A:208:ARG:HB2	3:A:707:HOH:O	1.88	0.74
1:B:146:ASP:H	1:B:147:PRO:HD2	1.49	0.73
1:A:59:LEU:C	1:A:63:MSE:HE3	2.14	0.73
1:B:147:PRO:HA	1:B:149:SER:N	2.08	0.69
1:B:173:ILE:H	1:B:174:GLY:HA3	1.59	0.67
1:A:174:GLY:N	1:A:175:HIS:CA	2.53	0.66
1:B:60:LYS:HA	1:B:63:MSE:CE	2.24	0.65
1:A:59:LEU:HB3	1:A:63:MSE:CE	2.28	0.64
1:A:209:ALA:HB1	1:A:237:ARG:HG3	1.83	0.60
1:B:37:ARG:NH1	1:B:126:GLU:OE1	2.34	0.59
1:A:59:LEU:HB3	1:A:63:MSE:HE2	1.86	0.58
1:A:19:ALA:O	1:A:21:ARG:N	2.37	0.57
1:A:124:ILE:HG12	1:A:133:LEU:CD1	2.36	0.56
1:A:177:LEU:O	1:B:208:ARG:NH1	2.38	0.56
1:A:21:ARG:HD2	1:A:41:LEU:HB3	1.88	0.55
1:B:147:PRO:HG3	1:B:149:SER:O	2.08	0.53
1:B:173:ILE:H	1:B:174:GLY:CA	2.23	0.49
1:B:158:ARG:NH1	3:B:577:HOH:O	2.46	0.49
1:B:179:LEU:C	1:B:181:ASP:H	2.20	0.49
1:A:61:VAL:HG23	3:A:564:HOH:O	2.14	0.48
1:A:207:ARG:O	1:A:208:ARG:HD2	2.15	0.47
1:A:147:PRO:HA	3:A:720:HOH:O	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:194:LEU:HD22	1:B:227:ILE:HD11	1.98	0.46
1:B:32:GLU:HG2	1:B:33:GLU:OE2	2.16	0.45
1:A:59:LEU:O	1:A:63:MSE:HE3	2.17	0.44
1:B:162:ALA:HB1	1:B:163:PRO:HA	1.99	0.44
1:A:222:LEU:HB2	1:A:227:ILE:HD12	2.00	0.44
1:B:60:LYS:HE2	3:B:644:HOH:O	2.17	0.43
1:B:210:ASP:OD1	1:B:210:ASP:N	2.48	0.43
1:B:155:ILE:HG12	1:B:244:MSE:HE3	1.99	0.43
1:A:3:MSE:HE1	1:A:143:ILE:HG12	2.01	0.43
1:A:143:ILE:HD13	1:A:143:ILE:HG21	1.79	0.43
1:A:222:LEU:CB	1:A:227:ILE:HD12	2.49	0.43
1:B:134:TYR:CE2	1:B:163:PRO:HB3	2.54	0.42
1:A:124:ILE:HG12	1:A:133:LEU:HD12	2.00	0.42
1:B:175:HIS:HA	1:B:176:SER:HA	1.77	0.42
1:B:171:ALA:HB3	3:B:583:HOH:O	2.19	0.42
1:A:133:LEU:O	1:A:163:PRO:HA	2.20	0.42
1:A:190:ARG:NH1	3:A:533:HOH:O	2.54	0.41
1:B:105:ALA:HB3	1:B:130:ASP:OD2	2.21	0.41
1:A:146:ASP:HA	1:A:147:PRO:HD2	1.95	0.41
1:B:176:SER:O	1:B:177:LEU:C	2.64	0.41
1:A:17:ARG:HG2	1:A:18:CYS:N	2.36	0.40
1:A:146:ASP:HB3	1:A:149:SER:HB2	2.02	0.40
1:A:59:LEU:HD21	1:A:89:MSE:SE	2.71	0.40
1:B:31:VAL:HG22	1:B:56:ARG:HD3	2.03	0.40
1:B:89:MSE:HE3	1:B:104:VAL:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	248/250 (99%)	236 (95%)	8 (3%)	4 (2%)	7 2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	230/250 (92%)	211 (92%)	14 (6%)	5 (2%)	5	1
All	All	478/500 (96%)	447 (94%)	22 (5%)	9 (2%)	6	1

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	ARG
1	B	147	PRO
1	B	177	LEU
1	A	14	TYR
1	B	148	ALA
1	A	18	CYS
1	B	146	ASP
1	A	174	GLY
1	B	174	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/181 (104%)	182 (96%)	7 (4%)	30	20
1	B	171/181 (94%)	161 (94%)	10 (6%)	18	7
All	All	360/362 (99%)	343 (95%)	17 (5%)	23	12

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	11	VAL
1	A	18	CYS
1	A	22	GLU
1	A	32	GLU
1	A	166	LEU
1	A	183	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	31	VAL
1	B	32	GLU
1	B	43	GLU
1	B	88	ARG
1	B	133	LEU
1	B	136	GLU
1	B	149	SER
1	B	194	LEU
1	B	208	ARG
1	B	210	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	7	HIS
1	A	16	ASN
1	A	127	ASN
1	A	204	ASN
1	B	91	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	241/250 (96%)	0.66	25 (10%)	11 13	28, 41, 75, 90	0
1	B	224/250 (89%)	1.20	52 (23%)	2 2	22, 47, 87, 99	1 (0%)
All	All	465/500 (93%)	0.92	77 (16%)	4 4	22, 43, 79, 99	1 (0%)

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	19	ALA	7.6
1	B	173	ILE	6.9
1	B	177	LEU	6.5
1	A	19	ALA	6.2
1	B	249	ALA	6.0
1	A	148	ALA	5.5
1	B	174	GLY	5.5
1	A	147	PRO	5.4
1	A	171	ALA	5.3
1	A	11	VAL	4.8
1	B	175	HIS	4.7
1	B	151	ASP	4.6
1	A	16	ASN	4.4
1	A	173	ILE	4.4
1	B	148	ALA	4.3
1	B	150	GLY	4.3
1	B	176	SER	4.3
1	B	30	VAL	4.1
1	B	248	VAL	4.1
1	B	178	SER	4.1
1	A	176	SER	4.0
1	B	162	ALA	4.0
1	B	147	PRO	4.0
1	A	20	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	TYR	3.9
1	A	18	CYS	3.9
1	A	174	GLY	3.8
1	B	183	ASP	3.7
1	B	145	PRO	3.6
1	B	140	VAL	3.6
1	A	21	ARG	3.5
1	B	146	ASP	3.5
1	B	31	VAL	3.5
1	A	181	ASP	3.4
1	B	149	SER	3.3
1	A	175	HIS	3.2
1	A	183	ASP	3.2
1	B	136	GLU	3.1
1	B	172	ARG	3.1
1	B	26	VAL	3.0
1	B	182	GLY	3.0
1	A	144	LEU	2.9
1	B	144	LEU	2.9
1	B	21	ARG	2.8
1	B	29	VAL	2.8
1	B	42	ALA	2.8
1	A	208	ARG	2.8
1	B	171	ALA	2.7
1	B	181	ASP	2.6
1	A	150	GLY	2.6
1	B	210	ASP	2.6
1	B	141	VAL	2.6
1	A	178	SER	2.5
1	B	169	SER	2.5
1	B	160	GLY	2.4
1	B	123	LEU	2.4
1	B	25	THR	2.4
1	A	17	ARG	2.4
1	B	208	ARG	2.4
1	B	142	GLY	2.4
1	A	13	ASP	2.3
1	B	119	VAL	2.3
1	B	159	ALA	2.2
1	A	210	ASP	2.2
1	B	217	GLU	2.2
1	B	206	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	207	ARG	2.1
1	B	45	ALA	2.1
1	B	28	GLN	2.1
1	B	116	PHE	2.1
1	A	217	GLU	2.0
1	B	39	THR	2.0
1	B	23	GLY	2.0
1	B	154	GLY	2.0
1	B	179	LEU	2.0
1	A	81	GLY	2.0
1	B	180	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NA	A	501	1/1	0.95	0.06	34,34,34,34	0
2	NA	B	502	1/1	0.95	0.09	37,37,37,37	0
2	NA	A	503	1/1	0.98	0.04	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.