



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:54 PM UTC

PDB ID : 7O3L / pdb_00007o3l
Title : Crystal Structure of AcrB Double Mutant
Authors : Ababou, A.
Deposited on : 2021-04-02
Resolution : 3.53 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1025 (3.56-3.48)
Clashscore	190562	1079 (3.56-3.48)
Ramachandran outliers	187476	1052 (3.56-3.48)
Sidechain outliers	187428	1053 (3.56-3.48)
RSRZ outliers	180081	1024 (3.56-3.48)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1069	% 31%51%14%..
1	B	1069	2% 32%51%13%..
1	C	1069	3% 28%54%15%..
1	D	1069	38%49%10%..
1	E	1069	39%48%10%.

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	1069	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LMT	A	2000	X	-	-	-
2	LMT	B	2000	X	-	-	-
2	LMT	C	2000	X	-	-	-
2	LMT	D	2000	X	-	-	-
2	LMT	E	2000	X	-	-	-
2	LMT	F	2000	X	-	-	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 47722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Efflux pump membrane transporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1044	Total	C	N	O	S	0	0	0
			7930	5093	1315	1479	43			
1	B	1042	Total	C	N	O	S	0	0	0
			7913	5083	1311	1476	43			
1	C	1042	Total	C	N	O	S	0	0	0
			7913	5083	1311	1476	43			
1	D	1044	Total	C	N	O	S	0	0	0
			7930	5093	1315	1479	43			
1	E	1042	Total	C	N	O	S	0	0	0
			7913	5083	1311	1476	43			
1	F	1042	Total	C	N	O	S	0	0	0
			7913	5083	1311	1476	43			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP E2QH56
A	-18	GLY	-	expression tag	UNP E2QH56
A	-17	SER	-	expression tag	UNP E2QH56
A	-16	SER	-	expression tag	UNP E2QH56
A	-15	HIS	-	expression tag	UNP E2QH56
A	-14	HIS	-	expression tag	UNP E2QH56
A	-13	HIS	-	expression tag	UNP E2QH56
A	-12	HIS	-	expression tag	UNP E2QH56
A	-11	HIS	-	expression tag	UNP E2QH56
A	-10	HIS	-	expression tag	UNP E2QH56
A	-9	SER	-	expression tag	UNP E2QH56
A	-8	SER	-	expression tag	UNP E2QH56
A	-7	GLY	-	expression tag	UNP E2QH56
A	-6	LEU	-	expression tag	UNP E2QH56
A	-5	VAL	-	expression tag	UNP E2QH56
A	-4	PRO	-	expression tag	UNP E2QH56
A	-3	ARG	-	expression tag	UNP E2QH56

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP E2QH56
A	-1	SER	-	expression tag	UNP E2QH56
A	0	HIS	-	expression tag	UNP E2QH56
A	615	ALA	PHE	engineered mutation	UNP E2QH56
A	617	ALA	PHE	engineered mutation	UNP E2QH56
B	-19	MET	-	initiating methionine	UNP E2QH56
B	-18	GLY	-	expression tag	UNP E2QH56
B	-17	SER	-	expression tag	UNP E2QH56
B	-16	SER	-	expression tag	UNP E2QH56
B	-15	HIS	-	expression tag	UNP E2QH56
B	-14	HIS	-	expression tag	UNP E2QH56
B	-13	HIS	-	expression tag	UNP E2QH56
B	-12	HIS	-	expression tag	UNP E2QH56
B	-11	HIS	-	expression tag	UNP E2QH56
B	-10	HIS	-	expression tag	UNP E2QH56
B	-9	SER	-	expression tag	UNP E2QH56
B	-8	SER	-	expression tag	UNP E2QH56
B	-7	GLY	-	expression tag	UNP E2QH56
B	-6	LEU	-	expression tag	UNP E2QH56
B	-5	VAL	-	expression tag	UNP E2QH56
B	-4	PRO	-	expression tag	UNP E2QH56
B	-3	ARG	-	expression tag	UNP E2QH56
B	-2	GLY	-	expression tag	UNP E2QH56
B	-1	SER	-	expression tag	UNP E2QH56
B	0	HIS	-	expression tag	UNP E2QH56
B	615	ALA	PHE	engineered mutation	UNP E2QH56
B	617	ALA	PHE	engineered mutation	UNP E2QH56
C	-19	MET	-	initiating methionine	UNP E2QH56
C	-18	GLY	-	expression tag	UNP E2QH56
C	-17	SER	-	expression tag	UNP E2QH56
C	-16	SER	-	expression tag	UNP E2QH56
C	-15	HIS	-	expression tag	UNP E2QH56
C	-14	HIS	-	expression tag	UNP E2QH56
C	-13	HIS	-	expression tag	UNP E2QH56
C	-12	HIS	-	expression tag	UNP E2QH56
C	-11	HIS	-	expression tag	UNP E2QH56
C	-10	HIS	-	expression tag	UNP E2QH56
C	-9	SER	-	expression tag	UNP E2QH56
C	-8	SER	-	expression tag	UNP E2QH56
C	-7	GLY	-	expression tag	UNP E2QH56
C	-6	LEU	-	expression tag	UNP E2QH56
C	-5	VAL	-	expression tag	UNP E2QH56

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	PRO	-	expression tag	UNP E2QH56
C	-3	ARG	-	expression tag	UNP E2QH56
C	-2	GLY	-	expression tag	UNP E2QH56
C	-1	SER	-	expression tag	UNP E2QH56
C	0	HIS	-	expression tag	UNP E2QH56
C	615	ALA	PHE	engineered mutation	UNP E2QH56
C	617	ALA	PHE	engineered mutation	UNP E2QH56
D	-19	MET	-	initiating methionine	UNP E2QH56
D	-18	GLY	-	expression tag	UNP E2QH56
D	-17	SER	-	expression tag	UNP E2QH56
D	-16	SER	-	expression tag	UNP E2QH56
D	-15	HIS	-	expression tag	UNP E2QH56
D	-14	HIS	-	expression tag	UNP E2QH56
D	-13	HIS	-	expression tag	UNP E2QH56
D	-12	HIS	-	expression tag	UNP E2QH56
D	-11	HIS	-	expression tag	UNP E2QH56
D	-10	HIS	-	expression tag	UNP E2QH56
D	-9	SER	-	expression tag	UNP E2QH56
D	-8	SER	-	expression tag	UNP E2QH56
D	-7	GLY	-	expression tag	UNP E2QH56
D	-6	LEU	-	expression tag	UNP E2QH56
D	-5	VAL	-	expression tag	UNP E2QH56
D	-4	PRO	-	expression tag	UNP E2QH56
D	-3	ARG	-	expression tag	UNP E2QH56
D	-2	GLY	-	expression tag	UNP E2QH56
D	-1	SER	-	expression tag	UNP E2QH56
D	0	HIS	-	expression tag	UNP E2QH56
D	615	ALA	PHE	engineered mutation	UNP E2QH56
D	617	ALA	PHE	engineered mutation	UNP E2QH56
E	-19	MET	-	initiating methionine	UNP E2QH56
E	-18	GLY	-	expression tag	UNP E2QH56
E	-17	SER	-	expression tag	UNP E2QH56
E	-16	SER	-	expression tag	UNP E2QH56
E	-15	HIS	-	expression tag	UNP E2QH56
E	-14	HIS	-	expression tag	UNP E2QH56
E	-13	HIS	-	expression tag	UNP E2QH56
E	-12	HIS	-	expression tag	UNP E2QH56
E	-11	HIS	-	expression tag	UNP E2QH56
E	-10	HIS	-	expression tag	UNP E2QH56
E	-9	SER	-	expression tag	UNP E2QH56
E	-8	SER	-	expression tag	UNP E2QH56
E	-7	GLY	-	expression tag	UNP E2QH56

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-6	LEU	-	expression tag	UNP E2QH56
E	-5	VAL	-	expression tag	UNP E2QH56
E	-4	PRO	-	expression tag	UNP E2QH56
E	-3	ARG	-	expression tag	UNP E2QH56
E	-2	GLY	-	expression tag	UNP E2QH56
E	-1	SER	-	expression tag	UNP E2QH56
E	0	HIS	-	expression tag	UNP E2QH56
E	615	ALA	PHE	engineered mutation	UNP E2QH56
E	617	ALA	PHE	engineered mutation	UNP E2QH56
F	-19	MET	-	initiating methionine	UNP E2QH56
F	-18	GLY	-	expression tag	UNP E2QH56
F	-17	SER	-	expression tag	UNP E2QH56
F	-16	SER	-	expression tag	UNP E2QH56
F	-15	HIS	-	expression tag	UNP E2QH56
F	-14	HIS	-	expression tag	UNP E2QH56
F	-13	HIS	-	expression tag	UNP E2QH56
F	-12	HIS	-	expression tag	UNP E2QH56
F	-11	HIS	-	expression tag	UNP E2QH56
F	-10	HIS	-	expression tag	UNP E2QH56
F	-9	SER	-	expression tag	UNP E2QH56
F	-8	SER	-	expression tag	UNP E2QH56
F	-7	GLY	-	expression tag	UNP E2QH56
F	-6	LEU	-	expression tag	UNP E2QH56
F	-5	VAL	-	expression tag	UNP E2QH56
F	-4	PRO	-	expression tag	UNP E2QH56
F	-3	ARG	-	expression tag	UNP E2QH56
F	-2	GLY	-	expression tag	UNP E2QH56
F	-1	SER	-	expression tag	UNP E2QH56
F	0	HIS	-	expression tag	UNP E2QH56
F	615	ALA	PHE	engineered mutation	UNP E2QH56
F	617	ALA	PHE	engineered mutation	UNP E2QH56

- Molecule 2 is DODECYL-BETA-D-MALTOSIDE (CCD ID: LMT) (formula: C₂₄H₄₆O₁₁).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 35	C 24	O 11	0	0
2	B	1	Total 35	C 24	O 11	0	0
2	C	1	Total 35	C 24	O 11	0	0
2	D	1	Total 35	C 24	O 11	0	0
2	E	1	Total 35	C 24	O 11	0	0
2	F	1	Total 35	C 24	O 11	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Efflux pump membrane transporter





● Molecule 1: Efflux pump membrane transporter











Met	GLY	SER	SER	HIS	HIS	HIS	HIS	SER	SER	GLY	VAL	PRO	ARG	GLY	SER	HIS	MET	P2	N3	F4	F5	D6	I7	R8	P9	I10	F11	A12	W13	V14	I15	A16	I17	M20	G23	A26	I27	L30	P31	V32	Y35	P36	T37	I38	P41	A42	V43	T44	I45	S46
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	152.15Å 156.72Å 218.75Å 90.00° 92.40° 90.00°	Depositor
Resolution (Å)	19.93 – 3.53 19.93 – 3.53	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.93-3.53) 99.2 (19.93-3.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 3.52Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.264 , 0.354 0.266 , 0.355	Depositor DCC
R_{free} test set	6294 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	104.6	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.060 for -k,-h,-l 0.074 for k,h,-l 0.074 for h,-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	47722	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.61 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8693e-04.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	4/8080 (0.0%)	1.34	75/10972 (0.7%)
1	B	0.92	5/8062 (0.1%)	1.37	94/10947 (0.9%)
1	C	0.94	5/8062 (0.1%)	1.42	117/10947 (1.1%)
1	D	0.93	3/8080 (0.0%)	1.34	65/10972 (0.6%)
1	E	0.97	3/8062 (0.0%)	1.36	71/10947 (0.6%)
1	F	0.97	5/8062 (0.1%)	1.36	78/10947 (0.7%)
All	All	0.94	25/48408 (0.1%)	1.36	500/65732 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	F	0	1
All	All	0	2

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	897	ILE	CA-CB	14.01	1.61	1.54
1	D	905	VAL	CA-CB	11.10	1.59	1.54
1	E	367	ILE	CA-CB	-10.04	1.49	1.54
1	F	367	ILE	CA-CB	-7.79	1.50	1.54
1	C	905	VAL	CA-CB	7.05	1.58	1.54

The worst 5 of 500 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	559	LEU	CA-C-N	15.84	136.07	119.76
1	E	559	LEU	C-N-CA	15.84	136.07	119.76
1	B	559	LEU	CA-C-N	15.77	135.53	119.76
1	B	559	LEU	C-N-CA	15.77	135.53	119.76
1	D	35	TYR	CA-C-N	-12.55	107.17	119.85

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1041	GLU	Peptide
1	F	1041	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7930	0	8072	680	1
1	B	7913	0	8058	719	1
1	C	7913	0	8058	777	0
1	D	7930	0	8072	586	0
1	E	7913	0	8058	532	0
1	F	7913	0	8058	600	0
2	A	35	0	46	7	0
2	B	35	0	46	3	0
2	C	35	0	46	5	0
2	D	35	0	46	4	0
2	E	35	0	46	17	0
2	F	35	0	46	2	0
All	All	47722	0	48652	3737	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 3737 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:536:ARG:HH21	2:E:2000:LMT:C3B	1.42	1.33
1:E:536:ARG:NE	2:E:2000:LMT:H4B	1.48	1.26
1:E:536:ARG:HE	2:E:2000:LMT:C4B	1.56	1.19
1:A:541:TYR:OH	2:A:2000:LMT:H6E	1.41	1.18
1:C:74:ASN:HB3	1:C:95:GLU:HB2	1.34	1.10

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:529:ASP:OD2	1:B:525:HIS:NE2[2_555]	1.98	0.22

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1042/1069 (98%)	891 (86%)	117 (11%)	34 (3%)	3	23
1	B	1040/1069 (97%)	893 (86%)	126 (12%)	21 (2%)	6	32
1	C	1040/1069 (97%)	874 (84%)	132 (13%)	34 (3%)	3	23
1	D	1042/1069 (98%)	904 (87%)	113 (11%)	25 (2%)	4	29
1	E	1040/1069 (97%)	899 (86%)	124 (12%)	17 (2%)	7	36
1	F	1040/1069 (97%)	898 (86%)	110 (11%)	32 (3%)	3	24
All	All	6244/6414 (97%)	5359 (86%)	722 (12%)	163 (3%)	4	27

5 of 163 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	SER
1	A	507	GLU
1	A	512	PHE
1	A	992	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	995	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	848/870 (98%)	701 (83%)	147 (17%)	2	12
1	B	846/870 (97%)	705 (83%)	141 (17%)	2	13
1	C	846/870 (97%)	700 (83%)	146 (17%)	2	12
1	D	848/870 (98%)	738 (87%)	110 (13%)	4	21
1	E	846/870 (97%)	725 (86%)	121 (14%)	3	18
1	F	846/870 (97%)	718 (85%)	128 (15%)	3	17
All	All	5080/5220 (97%)	4287 (84%)	793 (16%)	2	16

5 of 793 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	362	PHE
1	E	270	LEU
1	D	463	THR
1	D	353	LEU
1	D	842	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	642	ASN
1	F	104	GLN
1	F	228	GLN
1	B	692	HIS
1	B	667	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LMT	C	2000	-	36,36,36	1.81	9 (25%)	47,47,47	1.38	8 (17%)
2	LMT	F	2000	-	36,36,36	1.90	11 (30%)	47,47,47	2.19	15 (31%)
2	LMT	E	2000	-	36,36,36	1.82	11 (30%)	47,47,47	2.51	16 (34%)
2	LMT	B	2000	-	36,36,36	1.84	10 (27%)	47,47,47	1.59	7 (14%)
2	LMT	D	2000	-	36,36,36	1.93	9 (25%)	47,47,47	1.51	9 (19%)
2	LMT	A	2000	-	36,36,36	1.97	9 (25%)	47,47,47	1.69	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LMT	C	2000	-	3/3/10/10	10/21/61/61	0/2/2/2
2	LMT	F	2000	-	3/3/10/10	15/21/61/61	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LMT	E	2000	-	2/2/10/10	12/21/61/61	0/2/2/2
2	LMT	B	2000	-	2/2/10/10	10/21/61/61	0/2/2/2
2	LMT	D	2000	-	2/2/10/10	14/21/61/61	0/2/2/2
2	LMT	A	2000	-	2/2/10/10	10/21/61/61	0/2/2/2

The worst 5 of 59 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2000	LMT	O1'-C1'	4.74	1.48	1.40
2	F	2000	LMT	O1'-C1'	4.52	1.47	1.40
2	A	2000	LMT	O5'-C5'	4.41	1.55	1.44
2	C	2000	LMT	O5'-C5'	4.25	1.54	1.44
2	F	2000	LMT	O5'-C5'	4.23	1.54	1.44

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2000	LMT	C4B-C3B-C2B	7.10	123.29	110.83
2	E	2000	LMT	O3B-C3B-C4B	-7.02	93.83	110.38
2	F	2000	LMT	O1B-C1B-C2B	5.77	122.29	108.09
2	A	2000	LMT	O5B-C5B-C4B	5.62	119.82	109.70
2	E	2000	LMT	O5B-C1B-C2B	5.45	121.57	110.37

5 of 14 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	2000	LMT	C3'
2	A	2000	LMT	C4B
2	B	2000	LMT	C3'
2	B	2000	LMT	C4B
2	C	2000	LMT	C2'

5 of 71 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2000	LMT	O5'-C1'-O1'-C1
2	C	2000	LMT	O5B-C5B-C6B-O6B
2	F	2000	LMT	O5B-C5B-C6B-O6B
2	D	2000	LMT	O5B-C5B-C6B-O6B
2	A	2000	LMT	O5'-C5'-C6'-O6'


There are no ring outliers.


6 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2000	LMT	5	0
2	F	2000	LMT	2	0
2	E	2000	LMT	17	0
2	B	2000	LMT	3	0
2	D	2000	LMT	4	0
2	A	2000	LMT	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1044/1069 (97%)	-0.18	10 (0%) 79 54	13, 49, 91, 117	0
1	B	1042/1069 (97%)	-0.04	26 (2%) 58 33	11, 49, 101, 139	0
1	C	1042/1069 (97%)	0.04	31 (2%) 52 29	8, 62, 104, 135	0
1	D	1044/1069 (97%)	-0.35	2 (0%) 91 81	9, 36, 74, 116	0
1	E	1042/1069 (97%)	-0.52	0 100 100	7, 28, 59, 88	0
1	F	1042/1069 (97%)	-0.46	1 (0%) 92 85	5, 30, 61, 103	0
All	All	6256/6414 (97%)	-0.25	70 (1%) 78 52	5, 40, 91, 139	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	715	SER	8.5
1	C	116	PRO	6.0
1	C	856	GLY	5.7
1	B	78	MET	4.7
1	B	870	GLY	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	LMT	B	2000	35/35	0.69	0.12	11,43,62,64	0
2	LMT	D	2000	35/35	0.78	0.11	20,39,53,62	0
2	LMT	E	2000	35/35	0.78	0.10	14,53,70,88	0
2	LMT	A	2000	35/35	0.81	0.10	31,50,62,70	0
2	LMT	F	2000	35/35	0.82	0.10	14,40,70,74	0
2	LMT	C	2000	35/35	0.90	0.08	12,33,43,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around LMT D 2000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT E 2000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 2000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT F 2000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LMT C 2000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.