



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:48 PM UTC

PDB ID : 4O4I / pdb_00004o4i
Title : Tubulin-Laulimalide-Epothilone A complex
Authors : Prota, A.E.; Bargsten, K.; Northcote, P.T.; Marsh, M.; Altmann, K.H.; Miller, J.H.; Diaz, J.F.; Steinmetz, M.O.
Deposited on : 2013-12-18
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	2% 75% 21% ..
1	C	451	79% 16% ..
2	B	445	2% 71% 23% ..
2	D	445	2% 75% 19% ..
3	E	143	2% 50% 28% 6% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	4% 75% 15% 9%

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 18304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	0	0
			3416	2163	581	650	22			
1	C	440	Total	C	N	O	S	0	1	0
			3440	2177	584	656	23			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	428	Total	C	N	O	S	0	1	0
			3372	2117	577	651	27			
2	D	431	Total	C	N	O	S	0	1	0
			3381	2120	579	654	28			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	2	0
			1013	625	184	199	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	ILE	cloning artifact	UNP P63043
E	4	ALA	SER	cloning artifact	UNP P63043

- Molecule 4 is a protein called Tubulin-tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	351	Total	C	N	O	S	0	8	0
			2898	1867	490	527	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	2	Total	Mg	0	0
			2	2		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		
7	B	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total 28	C 10	N 5	O 11	P 2	0	0
8	D	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 9 is EPOTHILONE A (CCD ID: EP) (formula: C₂₆H₃₉NO₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	N	O	S	0	0
			34	26	1	6	1		
9	D	1	Total	C	N	O	S	0	0
			34	26	1	6	1		


- Molecule 10 is Laulimalide (CCD ID: LLM) (formula: C₃₀H₄₂O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	5	0
			37	30	7		


- Molecule 11 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES)

(formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
11	B	1	12	6	1	4	1	0	0

- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: C₁₁H₁₈N₅O₁₂P₃).




- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	112	Total 112	O 112	0	0
13	B	90	Total 90	O 90	0	0
13	C	161	Total 161	O 161	0	0
13	D	63	Total 63	O 63	0	0
13	E	34	Total 34	O 34	0	0
13	F	47	Total 47	O 47	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tubulin alpha-1B chain




• Molecule 1: Tubulin alpha-1B chain




• Molecule 2: Tubulin beta-2B chain







● Molecule 2: Tubulin beta-2B chain




● Molecule 3: Stathmin-4



● Molecule 4: Tubulin-tyrosine ligase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.20Å 158.03Å 180.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.21 – 2.40 52.21 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (52.21-2.40) 99.7 (52.21-2.40)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.188 , 0.214 0.190 , 0.216	Depositor DCC
R_{free} test set	5886 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	46.2	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18304	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLM, ACP, EP, GDP, CA, MES, MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3494	0.76	2/4743 (0.0%)
1	C	0.43	0/3521	0.84	2/4780 (0.0%)
2	B	0.42	0/3450	0.78	1/4672 (0.0%)
2	D	0.43	0/3458	0.82	1/4684 (0.0%)
3	E	0.41	0/1027	0.76	0/1362
4	F	0.36	0/2985	0.77	1/4032 (0.0%)
All	All	0.41	0/17935	0.79	7/24273 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	71	GLU	CA-C-N	-5.97	113.81	121.04
1	C	71	GLU	C-N-CA	-5.97	113.81	121.04
2	D	98	GLY	N-CA-C	-5.89	106.35	111.95
1	A	71	GLU	CA-C-N	-5.70	114.14	121.04
1	A	71	GLU	C-N-CA	-5.70	114.14	121.04

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3416	0	3331	63	0
1	C	3440	0	3354	54	0
2	B	3372	0	3255	82	0
2	D	3381	0	3258	65	0
3	E	1013	0	1030	38	0
4	F	2898	0	2914	47	0
5	A	32	0	12	1	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	B	2	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
8	B	28	0	12	0	0
8	D	28	0	12	1	0
9	B	34	0	39	6	0
9	D	34	0	39	0	0
10	B	37	0	42	4	0
11	B	12	0	12	3	0
12	F	31	0	14	4	0
13	A	112	0	0	4	0
13	B	90	0	0	7	0
13	C	161	0	0	10	0
13	D	63	0	0	6	0
13	E	34	0	0	5	0
13	F	47	0	0	5	0
All	All	18304	0	17336	334	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 334 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:184:LYS:NZ	4:F:185:TYR:O	2.01	0.94
2:B:276:THR:OG1	9:B:503:EP:N20	2.00	0.92
4:F:74:LYS:NZ	4:F:331:GLU:OE2	2.02	0.92
1:C:163:LYS:NZ	13:C:749:HOH:O	2.03	0.91
3:E:10:GLU:O	3:E:12:ASN:ND2	2.03	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/451 (96%)	426 (98%)	9 (2%)	0	100	100
1	C	439/451 (97%)	431 (98%)	8 (2%)	0	100	100
2	B	427/445 (96%)	412 (96%)	14 (3%)	1 (0%)	43	58
2	D	430/445 (97%)	421 (98%)	9 (2%)	0	100	100
3	E	119/143 (83%)	119 (100%)	0	0	100	100
4	F	352/384 (92%)	341 (97%)	10 (3%)	1 (0%)	36	50
All	All	2202/2319 (95%)	2150 (98%)	50 (2%)	2 (0%)	48	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	321	GLY
4	F	154	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/379 (97%)	350 (95%)	18 (5%)	22	39
1	C	372/379 (98%)	360 (97%)	12 (3%)	34	56
2	B	371/383 (97%)	350 (94%)	21 (6%)	18	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	371/383 (97%)	355 (96%)	16 (4%)	26	44
3	E	110/127 (87%)	92 (84%)	18 (16%)	2	3
4	F	321/342 (94%)	313 (98%)	8 (2%)	42	64
All	All	1913/1993 (96%)	1820 (95%)	93 (5%)	23	39

5 of 93 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	79	ARG
3	E	49	GLU
2	D	138	THR
2	D	345	GLU
3	E	62	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
4	F	306	HIS
4	F	333	ASN
1	C	393	HIS
2	D	15	GLN
2	D	167	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	MES	B	507	-	12,12,12	2.25	1 (8%)	15,16,16	2.05	2 (13%)
5	GTP	C	501	6	33,34,34	0.95	1 (3%)	50,54,54	1.52	11 (22%)
8	GDP	B	501	6	29,30,30	1.14	2 (6%)	45,47,47	1.69	7 (15%)
10	LLM	B	505	-	39,40,40	2.57	15 (38%)	41,55,55	2.05	10 (24%)
8	GDP	D	501	6	29,30,30	1.15	3 (10%)	45,47,47	1.70	7 (15%)
12	ACP	F	500	6	31,33,33	1.68	10 (32%)	47,52,52	1.79	11 (23%)
5	GTP	A	501	6	33,34,34	0.91	1 (3%)	50,54,54	1.57	15 (30%)
9	EP	D	503	-	36,36,36	1.41	4 (11%)	48,53,53	1.99	13 (27%)
9	EP	B	503	-	36,36,36	1.28	5 (13%)	48,53,53	2.27	15 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	MES	B	507	-	-	4/6/14/14	0/1/1/1
5	GTP	C	501	6	-	3/22/38/38	0/3/3/3
8	GDP	B	501	6	-	5/16/32/32	0/3/3/3
10	LLM	B	505	-	-	16/39/64/64	0/3/4/4
8	GDP	D	501	6	-	5/16/32/32	0/3/3/3
12	ACP	F	500	6	-	5/19/38/38	0/3/3/3
5	GTP	A	501	6	-	5/22/38/38	0/3/3/3
9	EP	D	503	-	-	4/50/55/55	0/3/3/3
9	EP	B	503	-	-	10/50/55/55	0/3/3/3

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	B	507	MES	C8-S	-7.52	1.67	1.77
10	B	505	LLM	C22-C21	6.82	1.52	1.32
10	B	505	LLM	C5-C6	-5.38	1.40	1.50
10	B	505	LLM	C30-C25	-5.10	1.38	1.50
12	F	500	ACP	C5-C4	4.86	1.47	1.39

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	505	LLM	C20-C21-C22	-7.42	113.66	124.94
9	B	503	EP	C32-C27-C24	-6.89	113.93	123.31
11	B	507	MES	C5-N4-C3	6.47	122.78	108.84
9	D	503	EP	C27-O26-C24	6.10	65.07	60.67
9	B	503	EP	C27-O26-C24	5.80	64.85	60.67

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	501	GTP	C5'-O5'-PA-O3A
5	C	501	GTP	C5'-O5'-PA-O1A


There are no ring outliers.



7 monomers are involved in 20 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	B	507	MES	3	0
5	C	501	GTP	1	0
10	B	505	LLM	4	0
8	D	501	GDP	1	0
12	F	500	ACP	4	0
5	A	501	GTP	1	0
9	B	503	EP	6	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/451 (96%)	-0.39	3 (0%) 84 81	24, 40, 68, 119	0
1	C	440/451 (97%)	-0.57	1 (0%) 91 89	19, 32, 59, 84	1 (0%)
2	B	428/445 (96%)	-0.26	8 (1%) 66 62	23, 43, 81, 115	3 (0%)
2	D	431/445 (96%)	-0.23	7 (1%) 70 66	25, 47, 79, 107	7 (1%)
3	E	121/143 (84%)	-0.02	3 (2%) 58 54	25, 53, 100, 118	2 (1%)
4	F	351/384 (91%)	0.19	15 (4%) 40 36	27, 65, 136, 154	8 (2%)
All	All	2208/2319 (95%)	-0.26	37 (1%) 69 65	19, 44, 98, 154	21 (0%)

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	371	PRO	4.2
4	F	125	THR	4.1
2	D	282	GLN	3.5
1	A	262	TYR	3.5
2	B	438	ALA	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	LLM	B	505	37/37	0.77	0.21	72,99,120,123	5
9	EP	B	503	34/34	0.78	0.15	54,67,75,82	0
6	MG	B	502	1/1	0.80	0.17	54,54,54,54	0
6	MG	D	502	1/1	0.83	0.19	62,62,62,62	0
12	ACP	F	500	31/31	0.84	0.11	69,83,158,200	0
7	CA	B	506	1/1	0.87	0.13	80,80,80,80	0
11	MES	B	507	12/12	0.88	0.12	48,55,71,74	0
7	CA	C	503	1/1	0.88	0.11	68,68,68,68	0
6	MG	C	502	1/1	0.90	0.24	29,29,29,29	0
6	MG	F	501	1/1	0.91	0.11	77,77,77,77	0
6	MG	A	502	1/1	0.93	0.21	30,30,30,30	0
9	EP	D	503	34/34	0.94	0.08	35,46,60,63	0
7	CA	A	503	1/1	0.95	0.12	64,64,64,64	0
8	GDP	D	501	28/28	0.96	0.06	29,34,46,51	0
6	MG	B	504	1/1	0.98	0.16	28,28,28,28	0
8	GDP	B	501	28/28	0.98	0.05	18,27,32,35	0
5	GTP	A	501	32/32	0.98	0.05	19,26,34,42	0
5	GTP	C	501	32/32	0.98	0.05	16,22,27,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LLM B 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EP B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ACP F 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EP D 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.