



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:30 AM UTC

PDB ID : 4O4J / pdb_00004o4j
Title : Tubulin-Peloruside A complex
Authors : Prota, A.E.; Bargsten, K.; Northcote, P.T.; Marsh, M.; Altmann, K.H.; Miller, J.H.; Diaz, J.F.; Steinmetz, M.O.
Deposited on : 2013-12-18
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	2% 88% 8% ..
1	C	451	2% 90% 7% ..
2	B	445	2% 84% 11% ..
2	D	445	4% 81% 15% .
3	E	143	4% 78% 8% . 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	384	35% 48% 18% . 31%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 18124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	12	0
			3485	2213	587	661	24			
1	C	440	Total	C	N	O	S	0	15	0
			3496	2218	585	667	26			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	428	Total	C	N	O	S	0	12	0
			3424	2151	581	664	28			
2	D	431	Total	C	N	O	S	0	6	0
			3413	2145	580	659	29			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	123	Total	C	N	O	S	0	6	0
			1044	645	188	206	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	ILE	cloning artifact	UNP P63043
E	4	ALA	SER	cloning artifact	UNP P63043

- Molecule 4 is a protein called Tubulin-tyrosine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	265	Total	C	N	O	S	0	1	0
			2179	1409	370	386	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
9	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 10 is Peloruside A (CCD ID: POU) (formula: $C_{27}H_{48}O_{11}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			38	27	11		
10	D	1	Total	C	O	0	0
			38	27	11		

- Molecule 11 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	F	1	Total 31	C 11	N 5	O 12	P 3	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	191	Total O 191 191	0	0
12	B	210	Total O 210 210	0	0
12	C	264	Total O 264 264	0	0
12	D	116	Total O 116 116	0	0
12	E	43	Total O 43 43	0	0
12	F	21	Total O 21 21	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain





- Molecule 2: Tubulin beta-2B chain






• Molecule 3: Stathmin-4



• Molecule 4: Tubulin-tyrosine ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.37Å 157.84Å 180.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.30 – 2.20 72.30 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.30-2.20) 99.7 (72.30-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.201 , 0.239 0.204 , 0.241	Depositor DCC
R_{free} test set	7568 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18124	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GTP, MG, POU, GDP, ACP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3599	0.76	3/4887 (0.1%)
1	C	0.43	0/3619	0.77	1/4917 (0.0%)
2	B	0.40	0/3532	0.75	3/4783 (0.1%)
2	D	0.35	0/3506	0.73	2/4751 (0.0%)
3	E	0.35	0/1071	0.69	0/1423
4	F	0.30	0/2230	0.76	0/3006
All	All	0.38	0/17557	0.75	9/23767 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	197	ASN	N-CA-C	6.15	120.82	112.88
2	B	260	VAL	N-CA-C	5.37	113.46	108.15
1	A	359	PRO	O-C-N	5.24	123.61	121.15
2	B	56	ALA	CA-C-O	5.22	120.97	117.94
2	D	260	VAL	CA-C-N	5.13	124.83	119.64

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3485	0	3436	19	0
1	C	3496	0	3440	22	0
2	B	3424	0	3319	31	0
2	D	3413	0	3310	39	0
3	E	1044	0	1068	6	0
4	F	2179	0	2178	40	0
5	A	32	0	12	1	0
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
8	A	6	0	8	0	0
9	B	28	0	12	0	0
9	D	28	0	12	0	0
10	B	38	0	48	4	0
10	D	38	0	48	2	0
11	F	31	0	14	1	0
12	A	191	0	0	1	0
12	B	210	0	0	5	0
12	C	264	0	0	4	0
12	D	116	0	0	2	0
12	E	43	0	0	1	0
12	F	21	0	0	0	0
All	All	18124	0	16917	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:9:ILE:HG22	3:E:10:GLU:HG3	1.70	0.74
1:C:381:THR:HG22	1:C:383:ALA:H	1.51	0.73
2:B:298:SER:HB3	2:B:307:PRO:HD2	1.71	0.71
2:B:22:GLU:OE2	12:B:802:HOH:O	2.09	0.70
4:F:197:ARG:NH1	4:F:257:GLU:OE1	2.23	0.70

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/451 (100%)	440 (98%)	9 (2%)	0	100	100
1	C	453/451 (100%)	441 (97%)	12 (3%)	0	100	100
2	B	438/445 (98%)	430 (98%)	8 (2%)	0	100	100
2	D	435/445 (98%)	421 (97%)	14 (3%)	0	100	100
3	E	125/143 (87%)	123 (98%)	2 (2%)	0	100	100
4	F	253/384 (66%)	239 (94%)	12 (5%)	2 (1%)	16	16
All	All	2153/2319 (93%)	2094 (97%)	57 (3%)	2 (0%)	48	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	F	264	PHE
4	F	189	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/379 (101%)	371 (97%)	11 (3%)	37	51
1	C	386/379 (102%)	382 (99%)	4 (1%)	68	81
2	B	382/383 (100%)	373 (98%)	9 (2%)	43	58
2	D	378/383 (99%)	368 (97%)	10 (3%)	40	55
3	E	116/127 (91%)	110 (95%)	6 (5%)	21	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	F	241/342 (70%)	210 (87%)	31 (13%)	4	4
All	All	1885/1993 (95%)	1814 (96%)	71 (4%)	28	40

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	F	264	PHE
4	F	285	LEU
4	F	330	ILE
2	D	1	MET
1	C	384	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	339	ASN
3	E	12	ASN
4	F	348	GLN
3	E	64	GLN
1	C	50	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GTP	C	501	6	33,34,34	0.95	1 (3%)	50,54,54	1.50	9 (18%)
5	GTP	A	501	6	33,34,34	1.17	3 (9%)	50,54,54	1.51	10 (20%)
9	GDP	B	501	6	29,30,30	1.21	3 (10%)	45,47,47	1.82	8 (17%)
9	GDP	D	501	-	29,30,30	1.14	2 (6%)	45,47,47	1.67	6 (13%)
10	POU	D	503	-	38,39,39	2.15	6 (15%)	44,57,57	1.96	9 (20%)
8	GOL	A	504	-	5,5,5	0.37	0	5,5,5	0.31	0
11	ACP	F	401	-	31,33,33	1.71	9 (29%)	47,52,52	1.96	12 (25%)
10	POU	B	503	-	38,39,39	2.31	8 (21%)	44,57,57	2.12	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	C	501	6	-	6/22/38/38	0/3/3/3
5	GTP	A	501	6	-	4/22/38/38	0/3/3/3
9	GDP	B	501	6	-	5/16/32/32	0/3/3/3
9	GDP	D	501	-	-	3/16/32/32	0/3/3/3
10	POU	D	503	-	-	7/54/76/76	0/1/2/2
8	GOL	A	504	-	-	4/4/4/4	-
11	ACP	F	401	-	-	10/19/38/38	0/3/3/3
10	POU	B	503	-	-	6/54/76/76	0/1/2/2

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	B	503	POU	C22-C21	10.46	1.56	1.33
10	D	503	POU	C22-C21	9.35	1.54	1.33
10	B	503	POU	C2-C1	-5.27	1.39	1.52
10	D	503	POU	C2-C1	-5.11	1.39	1.52
11	F	401	ACP	C5-C4	4.76	1.47	1.39

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	B	503	POU	C13-C12-C11	-6.52	104.62	114.42
10	D	503	POU	C13-C12-C11	-6.28	104.98	114.42
9	B	501	GDP	C5-C4-N3	-6.19	118.54	128.39
11	F	401	ACP	C5-C4-N3	-5.78	118.76	126.72
10	B	503	POU	C15-O15-C1	5.76	126.93	116.58

There are no chirality outliers.

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	C	501	GTP	PB-O3B-PG-O2G
5	C	501	GTP	C5'-O5'-PA-O3A
5	C	501	GTP	C5'-O5'-PA-O1A
8	A	504	GOL	O1-C1-C2-O2


There are no ring outliers.

4 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	GTP	1	0
10	D	503	POU	2	0
11	F	401	ACP	1	0
10	B	503	POU	4	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand GTP C 501




Ligand GTP A 501







Ligand POU D 503



Ligand ACP F 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/451 (97%)	-0.29	7 (1%) 70 67	15, 32, 66, 112	12 (2%)
1	C	440/451 (97%)	-0.59	8 (1%) 67 64	10, 22, 45, 68	15 (3%)
2	B	428/445 (96%)	-0.36	7 (1%) 70 67	11, 27, 58, 112	14 (3%)
2	D	431/445 (96%)	0.11	16 (3%) 45 42	18, 44, 77, 102	12 (2%)
3	E	123/143 (86%)	0.29	6 (4%) 35 31	18, 44, 85, 121	6 (4%)
4	F	265/384 (69%)	2.16	136 (51%) 0 0	34, 79, 123, 153	1 (0%)
All	All	2126/2319 (91%)	0.06	180 (8%) 16 14	10, 35, 92, 153	60 (2%)

The worst 5 of 180 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	182	ILE	7.5
2	D	1	MET	6.0
3	E	143	ALA	5.8
4	F	271	LEU	5.4
4	F	275	LEU	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
11	ACP	F	401	31/31	0.70	0.15	67,106,145,166	0
6	MG	D	502	1/1	0.73	0.28	67,67,67,67	0
8	GOL	A	504	6/6	0.83	0.17	57,60,60,62	0
10	POU	D	503	38/38	0.84	0.14	35,59,74,77	0
10	POU	B	503	38/38	0.85	0.13	23,44,77,81	0
7	CA	A	503	1/1	0.87	0.11	68,68,68,68	0
9	GDP	D	501	28/28	0.95	0.08	27,36,45,53	0
6	MG	C	502	1/1	0.96	0.04	18,18,18,18	0
5	GTP	A	501	32/32	0.98	0.05	10,17,22,41	0
6	MG	A	502	1/1	0.99	0.02	14,14,14,14	0
6	MG	B	502	1/1	0.99	0.02	9,9,9,9	0
5	GTP	C	501	32/32	0.99	0.03	4,13,19,30	0
9	GDP	B	501	28/28	0.99	0.04	11,16,20,22	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP F 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around POU D 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around POU B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.