



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:47 PM UTC

PDB ID : 1O5B / pdb_00001o5b
Title : Dissecting and Designing Inhibitor Selectivity Determinants at the S1 site
Using an Artificial Ala190 Protease (Ala190 uPA)
Authors : Katz, B.A.; Luong, C.; Ho, J.D.; Somoza, J.R.; Gjerstad, E.; Tang, J.;
Williams, S.R.; Verner, E.; Mackman, R.L.; Young, W.B.; Sprengeler, P.A.;
Chan, H.; Mortara, K.; Janc, J.W.; McGrath, M.E.
Deposited on : 2003-09-09
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	23	4% 22% 9% 65%
2	B	253	2% 64% 29%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CIT	B	3	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5124 atoms, of which 2705 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	8	Total	C	H	N	O	S	0	0	0
			133	42	68	12	10	1			


- Molecule 2 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	245	Total	C	H	N	O	S	0	13	0
			3953	1255	1972	342	365	19			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	145	ALA	ASN	engineered mutation	UNP P00749
B	190	ALA	SER	engineered mutation	UNP P00749

- Molecule 3 is 4-IODOBENZO[B]THIOPHENE-2-CARBOXAMIDINE (CCD ID: ESI) (formula: C₉H₈IN₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	B	1	Total	C	H	I	N	S	0	0
			21	9	8	1	2	1		

- Molecule 4 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			18	6	5	7		
4	B	1	Total	C	H	O	0	0
			18	6	5	7		
4	B	1	Total	C	H	O	0	0
			18	6	5	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	13	Total	H	O	0	0
			39	26	13		
5	B	306	Total	H	O	0	2
			924	616	308		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Urokinase-type plasminogen activator



- Molecule 2: Urokinase-type plasminogen activator



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	81.87Å 49.78Å 66.35Å 90.00° 113.10° 90.00°	Depositor
Resolution (Å)	7.00 – 1.85 7.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	86.2 (7.00-1.85) 84.5 (7.00-1.85)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.64 (at 1.86Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.197 , 0.240 0.198 , 0.201	Depositor DCC
R_{free} test set	1730 reflections (9.69%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 90.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for $1/2^*h+3/2^*k, 1/2^*h-1/2^*k, -1/2^*h-1/2^*k-l$ 0.029 for $1/2^*h-3/2^*k, -1/2^*h-1/2^*k, -1/2^*h+1/2^*k-l$	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5124	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT, ESI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	0/65	1.71	2/83 (2.4%)
2	B	1.82	23/2078 (1.1%)	1.75	54/2814 (1.9%)
All	All	1.81	23/2143 (1.1%)	1.75	56/2897 (1.9%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	241	HIS	CG-ND1	-7.65	1.29	1.38
2	B	37	HIS	CG-ND1	-6.62	1.30	1.38
2	B	22	THR	CA-CB	6.39	1.62	1.53
2	B	57	HIS	CD2-NE2	-6.27	1.30	1.37
2	B	165	HIS	CG-ND1	-6.17	1.31	1.38
2	B	91	HIS	CG-ND1	-6.01	1.31	1.38
2	B	91	HIS	CD2-NE2	-5.98	1.31	1.37
2	B	165	HIS	CD2-NE2	-5.95	1.31	1.37
2	B	170(B)	HIS	CD2-NE2	-5.94	1.31	1.37
2	B	233	HIS	CG-ND1	-5.92	1.31	1.38
2	B	99	HIS	CG-ND1	-5.88	1.31	1.38
2	B	99	HIS	CD2-NE2	-5.86	1.31	1.37
2	B	37	HIS	CD2-NE2	-5.82	1.31	1.37
2	B	100	HIS	CG-ND1	-5.78	1.31	1.38
2	B	100	HIS	CD2-NE2	-5.62	1.31	1.37
2	B	233	HIS	CD2-NE2	-5.56	1.31	1.37
2	B	170(B)	HIS	CG-ND1	-5.53	1.32	1.38
2	B	57	HIS	CG-ND1	-5.51	1.32	1.38
2	B	56	THR	CA-CB	5.38	1.61	1.53
2	B	209	LEU	CA-CB	5.30	1.59	1.53
2	B	214	SER	CA-C	5.16	1.57	1.52
2	B	211	GLY	N-CA	5.11	1.50	1.45
2	B	129	ASP	N-CA	5.07	1.50	1.45

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	92	LYS	N-CA-C	-9.61	93.51	109.46
2	B	165	HIS	CA-CB-CG	-8.16	105.64	113.80
2	B	59	PHE	CA-CB-CG	-7.91	105.89	113.80
2	B	185	ASP	N-CA-C	-7.71	99.67	110.31
2	B	207	MET	N-CA-C	-7.63	96.12	108.41
2	B	199	LEU	N-CA-C	-7.50	94.90	108.02
2	B	17[A]	ILE	N-CA-C	-7.09	98.23	108.36
2	B	17[B]	ILE	N-CA-C	-7.09	98.23	108.36
2	B	73	LEU	N-CA-C	6.86	119.77	111.40
2	B	51	TRP	NE1-CE2-CZ2	6.62	140.03	130.10
2	B	84[A]	GLU	N-CA-C	-6.50	100.06	110.14
2	B	84[B]	GLU	N-CA-C	-6.50	100.06	110.14
2	B	107	LYS	N-CA-C	-6.45	98.76	109.46
2	B	38	VAL	N-CA-C	-6.37	99.19	108.11
2	B	37(A)	ARG	N-CA-C	-6.30	97.39	108.23
2	B	21	PHE	N-CA-C	-6.28	100.33	110.32
1	A	10	LYS	N-CA-C	-6.23	99.94	109.41
2	B	229	THR	N-CA-C	-6.16	100.47	110.20
2	B	209	LEU	N-CA-C	-6.12	99.19	108.67
2	B	234	PHE	CA-CB-CG	-6.12	107.69	113.80
2	B	125	SER	N-CA-C	-6.08	101.00	110.42
2	B	160	VAL	N-CA-CB	-5.98	101.30	112.36
2	B	99	HIS	N-CA-C	-5.92	99.25	108.90
2	B	29	TRP	CG-CD1-NE1	-5.88	102.56	110.20
2	B	193	GLY	N-CA-C	-5.86	107.31	114.92
2	B	190	ALA	N-CA-C	-5.83	102.14	110.59
2	B	121	ILE	N-CA-C	-5.80	100.20	109.20
2	B	170	GLN	CB-CG-CD	-5.78	102.77	112.60
2	B	175	GLU	N-CA-C	-5.74	105.77	112.89
2	B	123[A]	LEU	N-CA-C	-5.71	102.97	110.39
2	B	123[B]	LEU	N-CA-C	-5.71	102.97	110.39
2	B	226	GLY	N-CA-C	-5.64	101.63	111.62
1	A	12	GLN	N-CA-C	-5.62	96.60	107.44
2	B	215	TRP	NE1-CE2-CZ2	5.59	138.49	130.10
2	B	77	THR	N-CA-C	-5.50	100.92	109.72
2	B	186	TRP	NE1-CE2-CZ2	5.48	138.31	130.10
2	B	51	TRP	CG-CD1-NE1	-5.39	103.20	110.20
2	B	186	TRP	CG-CD1-NE1	-5.31	103.30	110.20
2	B	29	TRP	NE1-CE2-CZ2	5.30	138.06	130.10
2	B	116[B]	ARG	N-CA-C	-5.29	106.78	113.18
2	B	237	TRP	CG-CD1-NE1	-5.29	103.32	110.20
2	B	33	ILE	N-CA-CB	-5.27	104.08	111.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	165	HIS	N-CA-C	-5.25	105.23	111.69
2	B	241	HIS	CE1-NE2-CD2	-5.20	103.80	109.00
2	B	158	THR	CA-CB-OG1	-5.18	101.83	109.60
2	B	208	THR	CA-CB-OG1	-5.15	101.87	109.60
2	B	215	TRP	NE1-CE2-CD2	-5.14	100.72	107.40
2	B	51	TRP	NE1-CE2-CD2	-5.13	100.73	107.40
2	B	87	ASN	OD1-CG-ND2	-5.13	117.47	122.60
2	B	37	HIS	CA-CB-CG	-5.11	108.69	113.80
2	B	237	TRP	NE1-CE2-CZ2	5.11	137.76	130.10
2	B	233	HIS	CA-CB-CG	-5.08	108.72	113.80
2	B	26	ASN	CA-CB-CG	-5.06	107.54	112.60
2	B	135	SER	N-CA-C	-5.06	100.99	109.24
2	B	217	ARG	N-CA-C	-5.05	100.50	108.73
2	B	51	TRP	N-CA-C	5.01	117.07	108.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	65	68	68	1	0
2	B	1981	1972	1933	19	2
3	B	13	8	8	2	0
4	B	39	15	15	1	0
5	A	13	26	0	0	0
5	B	308	616	0	0	2
All	All	2419	2705	2024	20	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:81[A]:MET:HE1	2:B:113:GLN:H	1.33	0.92
2:B:92:LYS:H	2:B:92:LYS:HD2	1.69	0.56
2:B:27:GLN:HG3	2:B:139[A]:THR:HG21	1.85	0.56
2:B:51:TRP:CD1	2:B:242:THR:HG23	2.43	0.54
2:B:94:TYR:HH	2:B:99:HIS:HD1	1.56	0.52
2:B:60(B):TYR:CE1	2:B:61:LYS:HB2	2.45	0.51
1:A:12:GLN:HB3	1:A:15:GLN:NE2	2.28	0.49
2:B:138:ILE:C	2:B:139[A]:THR:HG23	2.38	0.48
2:B:192[A]:GLN:CG	3:B:251:ESI:I9	3.32	0.47
2:B:81[A]:MET:HE1	2:B:113:GLN:N	2.16	0.46
2:B:81[A]:MET:HE3	2:B:118:ILE:HD13	1.98	0.46
2:B:192[A]:GLN:HG3	3:B:251:ESI:I9	2.87	0.44
2:B:57:HIS:CD2	4:B:3:CIT:O1	2.72	0.43
2:B:47[B]:MET:SD	2:B:53:ILE:HB	2.59	0.43
2:B:91:HIS:ND1	2:B:92:LYS:O	2.51	0.43
2:B:27:GLN:CG	2:B:139[A]:THR:HG21	2.47	0.43
2:B:92:LYS:O	2:B:93:ASP:CB	2.67	0.43
2:B:47[B]:MET:HE1	2:B:238:ILE:CG2	2.49	0.42
2:B:72:ARG:HA	2:B:153:GLU:O	2.21	0.41
2:B:110(A):LYS:HG3	2:B:110(B):GLU:HG3	2.03	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:185(A):PRO:O	5:B:288:HOH:H1[4_546]	1.55	0.05
2:B:154:GLN:OE1	5:B:316:HOH:H2[2_656]	1.58	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	6/23 (26%)	6 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	255/253 (101%)	232 (91%)	20 (8%)	3 (1%)	10	2
All	All	261/276 (95%)	238 (91%)	20 (8%)	3 (1%)	10	3

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	97(B)	LEU
2	B	93	ASP
2	B	60	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	7/22 (32%)	6 (86%)	1 (14%)	3	0
2	B	224/218 (103%)	213 (95%)	11 (5%)	22	8
All	All	231/240 (96%)	219 (95%)	12 (5%)	19	7

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
2	B	20	GLU
2	B	56	THR
2	B	57	HIS
2	B	92	LYS
2	B	146	SER
2	B	147	THR
2	B	150	LEU
2	B	187	LYS
2	B	202	SER
2	B	209	LEU
2	B	240	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
2	B	78	GLN
2	B	101	ASN
2	B	204	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	CIT	B	2	-	12,12,12	3.17	7 (58%)	17,17,17	1.69	4 (23%)
4	CIT	B	1	-	12,12,12	2.32	5 (41%)	17,17,17	1.24	1 (5%)
4	CIT	B	3	-	12,12,12	3.55	7 (58%)	17,17,17	2.47	10 (58%)
3	ESI	B	251	-	14,14,14	1.59	3 (21%)	14,20,20	2.10	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	B	2	-	-	3/16/16/16	-
4	CIT	B	1	-	-	2/16/16/16	-
4	CIT	B	3	-	-	8/16/16/16	-
3	ESI	B	251	-	-	0/0/4/4	0/2/2/2

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	3	CIT	C2-C3	8.49	1.64	1.54
4	B	2	CIT	C2-C3	7.93	1.63	1.54
4	B	3	CIT	C3-C6	5.10	1.58	1.53
4	B	3	CIT	C4-C3	5.03	1.60	1.54
4	B	1	CIT	C2-C3	4.97	1.60	1.54
4	B	2	CIT	C4-C3	4.19	1.59	1.54
3	B	251	ESI	C1-C0	-3.76	1.38	1.45
4	B	1	CIT	C4-C3	3.52	1.58	1.54
4	B	2	CIT	C2-C1	3.18	1.60	1.50
3	B	251	ESI	C9-I9	-3.11	2.03	2.10
4	B	2	CIT	C3-C6	2.92	1.56	1.53
4	B	3	CIT	O2-C1	-2.74	1.21	1.30
4	B	1	CIT	O4-C5	-2.70	1.21	1.30
4	B	3	CIT	O4-C5	-2.67	1.22	1.30
4	B	2	CIT	C4-C5	2.57	1.58	1.50
4	B	1	CIT	O2-C1	-2.57	1.22	1.30
4	B	3	CIT	C2-C1	2.52	1.58	1.50
4	B	2	CIT	O4-C5	-2.34	1.23	1.30
4	B	1	CIT	O6-C6	-2.28	1.22	1.30
4	B	2	CIT	O2-C1	-2.16	1.23	1.30
4	B	3	CIT	O6-C6	-2.08	1.23	1.30
3	B	251	ESI	C4-C5	-2.02	1.37	1.44

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3	CIT	O7-C3-C2	-4.93	98.12	109.38
4	B	3	CIT	O7-C3-C4	-3.94	100.39	109.38
3	B	251	ESI	C4-C3-S2	-3.89	107.61	111.40
3	B	251	ESI	C3-C4-C5	3.38	115.16	111.42
3	B	251	ESI	C8-C9-C4	-3.37	118.57	121.63
4	B	3	CIT	C4-C3-C2	3.15	117.40	109.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3	CIT	O6-C6-C3	3.11	119.11	113.14
4	B	2	CIT	O5-C6-C3	-3.08	116.12	122.09
4	B	2	CIT	O6-C6-C3	2.91	118.72	113.14
4	B	3	CIT	C4-C3-C6	2.89	116.43	110.03
4	B	3	CIT	O3-C5-C4	-2.85	114.87	122.95
4	B	3	CIT	C3-C2-C1	2.66	121.20	113.92
3	B	251	ESI	C3-S2-C1	2.55	94.69	90.97
4	B	2	CIT	C2-C3-C6	2.48	115.52	110.03
4	B	3	CIT	C3-C4-C5	2.43	120.56	113.92
4	B	1	CIT	O5-C6-C3	-2.43	117.39	122.09
4	B	3	CIT	O4-C5-C4	2.28	121.56	114.35
3	B	251	ESI	C5-C1-S2	-2.21	106.32	113.14
3	B	251	ESI	C0-C1-S2	2.16	123.02	118.68
4	B	3	CIT	O5-C6-C3	-2.11	118.00	122.09
4	B	2	CIT	C3-C4-C5	2.03	119.48	113.92

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	3	CIT	C1-C2-C3-C4
4	B	3	CIT	C1-C2-C3-C6
4	B	3	CIT	C1-C2-C3-O7
4	B	3	CIT	O7-C3-C4-C5
4	B	3	CIT	C6-C3-C4-C5
4	B	3	CIT	C2-C3-C4-C5
4	B	2	CIT	C1-C2-C3-C4
4	B	1	CIT	C4-C3-C6-O5
4	B	1	CIT	C4-C3-C6-O6
4	B	2	CIT	C1-C2-C3-O7
4	B	3	CIT	C3-C4-C5-O4
4	B	2	CIT	C1-C2-C3-C6
4	B	3	CIT	C3-C4-C5-O3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	3	CIT	1	0
3	B	251	ESI	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	8/23 (34%)	0.33	1 (12%) 8 7	12, 26, 42, 47	1 (12%)
2	B	245/253 (96%)	-0.42	4 (1%) 70 74	6, 17, 33, 49	30 (12%)
All	All	253/276 (91%)	-0.39	5 (1%) 65 68	6, 17, 34, 49	31 (12%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	LEU	3.8
2	B	242	THR	2.9
2	B	37(A)	ARG	2.8
2	B	97(A)	THR	2.5
2	B	97	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CIT	B	2	13/13	0.71	0.14	39,44,49,50	0
4	CIT	B	3	13/13	0.79	0.14	44,48,52,54	6
4	CIT	B	1	13/13	0.89	0.07	17,24,30,31	0
3	ESI	B	251	13/13	0.99	0.05	6,13,17,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.