



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:33 PM UTC

PDB ID : 1O5D / pdb_00001o5d
Title : Dissecting and Designing Inhibitor Selectivity Determinants at the S1 site
Using an Artificial Ala190 Protease (Ala190 uPA)
Authors : Katz, B.A.; Luong, C.; Ho, J.D.; Somoza, J.R.; Gjerstad, E.; Tang, J.;
Williams, S.R.; Verner, E.; Mackman, R.L.; Young, W.B.; Sprengeler, P.A.;
Chan, H.; Mortara, K.; Janc, J.W.; McGrath, M.E.
Deposited on : 2003-09-09
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	152	3% 42% 17% • • 37%
2	H	254	3% 64% 29% 6% •
3	T	218	5% 37% 27% 6% 31%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CR9	H	258	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9593 atoms, of which 5038 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	L	96	1395	439	668	126	149	13	0	2	0


- Molecule 2 is a protein called Coagulation factor VII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	254	3965	1263	1977	353	359	13	0	3	0

- Molecule 3 is a protein called Tissue factor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	T	151	2406	775	1186	198	245	2	0	1	0

- Molecule 4 is 2-{5-[AMINO(IMINIO)METHYL]-6-FLUORO-1H-BENZIMIDAZOL-2-YL}-6-[(2-METHYLCYCLOHEXYL)OXY]BENZENOLATE (CCD ID: CR9) (formula: C₂₁H₂₃FN₄O₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	H	1	Total	C	F	H	N	O	0	0
			51	21	1	23	4	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	110	Total	H	O	0	0
			330	220	110		
5	H	292	Total	H	O	0	3
			885	590	295		
5	T	186	Total	H	O	0	1
			561	374	187		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Coagulation factor VII



• Molecule 2: Coagulation factor VII



• Molecule 3: Tissue factor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.54Å 69.03Å 79.12Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	7.00 – 2.05 7.00 – 2.05	Depositor EDS
% Data completeness (in resolution range)	77.3 (7.00-2.05) 75.3 (7.00-2.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 2.05Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.226 , 0.262 0.203 , 0.200	Depositor DCC
R_{free} test set	4073 reflections (10.15%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.894	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 122.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.003 for -l,k,h 0.028 for h,-k,-l 0.022 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9593	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CR9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	1.82	7/748 (0.9%)	1.77	13/1009 (1.3%)
2	H	1.99	31/2046 (1.5%)	1.93	63/2784 (2.3%)
3	T	1.69	4/1250 (0.3%)	1.87	39/1701 (2.3%)
All	All	1.87	42/4044 (1.0%)	1.89	115/5494 (2.1%)

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	115	HIS	CG-ND1	-8.55	1.28	1.38
2	H	76	HIS	CG-ND1	-7.98	1.29	1.38
2	H	101	HIS	CG-ND1	-7.97	1.29	1.38
3	T	41	LYS	C-N	-7.87	1.23	1.33
2	H	117	HIS	CG-ND1	-7.78	1.29	1.38
2	H	202	HIS	CG-ND1	-7.77	1.29	1.38
1	L	84	HIS	CG-ND1	-7.62	1.29	1.38
2	H	224	HIS	CG-ND1	-7.52	1.29	1.38
2	H	52	VAL	CA-CB	7.49	1.65	1.54
2	H	71	HIS	CD2-NE2	-7.40	1.29	1.37
1	L	105	HIS	CG-ND1	-7.29	1.30	1.38
2	H	109	HIS	CG-ND1	-7.13	1.30	1.38
2	H	199	HIS	CG-ND1	-7.10	1.30	1.38
2	H	57	HIS	CG-ND1	-6.90	1.30	1.38
2	H	117	HIS	CA-CB	6.78	1.64	1.53
2	H	129(F)	PHE	CA-CB	6.71	1.62	1.53
3	T	43	GLY	N-CA	6.42	1.50	1.44
1	L	125	VAL	CA-C	6.39	1.58	1.52
2	H	141	TRP	CA-CB	6.35	1.61	1.53
2	H	202	HIS	CA-CB	-6.32	1.44	1.53
2	H	191	CYS	CA-CB	6.08	1.62	1.53
1	L	105	HIS	CD2-NE2	-6.05	1.31	1.37
2	H	57	HIS	CD2-NE2	-5.84	1.31	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	21	VAL	CA-CB	5.72	1.61	1.54
2	H	78	GLY	N-CA	5.70	1.52	1.45
2	H	101	HIS	CD2-NE2	-5.60	1.31	1.37
2	H	71	HIS	CG-ND1	-5.53	1.32	1.38
2	H	224	HIS	CD2-NE2	-5.41	1.31	1.37
2	H	51	TRP	NE1-CE2	-5.35	1.31	1.37
2	H	76	HIS	CD2-NE2	-5.26	1.32	1.37
3	T	32	GLN	CA-CB	5.25	1.63	1.53
2	H	29	TRP	CA-CB	5.25	1.63	1.53
2	H	114	LEU	CA-C	-5.23	1.46	1.52
2	H	138[B]	VAL	CA-C	5.22	1.58	1.52
2	H	138[A]	VAL	CA-C	5.22	1.58	1.52
1	L	84	HIS	CD2-NE2	-5.17	1.32	1.37
1	L	115	HIS	CD2-NE2	-5.11	1.32	1.37
2	H	67	VAL	CA-CB	5.11	1.61	1.54
3	T	134	VAL	CA-CB	5.11	1.60	1.54
2	H	113	VAL	CA-CB	5.08	1.61	1.54
2	H	211	GLY	N-CA	5.03	1.50	1.45
2	H	249	GLY	N-CA	5.02	1.50	1.45

All (115) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	129(G)	VAL	N-CA-C	-11.51	91.37	108.87
3	T	40	THR	N-CA-C	-10.32	93.54	109.85
2	H	199	HIS	N-CA-C	-10.22	90.11	107.80
2	H	181	PHE	CA-CB-CG	9.78	123.58	113.80
3	T	28	LYS	N-CA-C	-9.50	95.36	109.42
2	H	256	PHE	CA-CB-CG	-9.16	104.64	113.80
2	H	117	HIS	CA-CB-CG	-8.61	105.19	113.80
3	T	197	THR	N-CA-C	-8.21	102.01	112.93
3	T	20	LYS	N-CA-C	-7.83	97.88	109.15
2	H	170(F)	GLY	N-CA-C	-7.72	100.00	110.97
3	T	65	LYS	N-CA-C	-7.64	102.88	111.14
2	H	190[A]	SER	N-CA-C	-7.63	99.16	110.46
2	H	190[B]	SER	N-CA-C	-7.63	99.16	110.46
3	T	19	PHE	CA-CB-CG	-7.62	106.18	113.80
2	H	250	VAL	N-CA-C	-7.58	104.47	111.67
2	H	164	MET	N-CA-C	-7.51	98.38	110.32
2	H	224	HIS	CA-CB-CG	-7.45	106.35	113.80
2	H	112	VAL	N-CA-C	-7.28	99.42	109.46
2	H	170(D)	LYS	N-CA-C	7.27	126.28	110.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	129(F)	PHE	CA-CB-CG	-7.19	106.61	113.80
2	H	109	HIS	CA-CB-CG	-7.18	106.62	113.80
2	H	100	ASN	OD1-CG-ND2	-7.13	115.47	122.60
3	T	84	GLU	N-CA-C	-7.09	103.20	112.41
1	L	48	ASP	N-CA-C	-7.05	100.37	110.59
3	T	156	TYR	N-CA-C	-7.03	95.83	110.80
1	L	92	VAL	N-CA-CB	-6.97	101.47	112.07
3	T	86	THR	N-CA-CB	-6.96	100.56	111.56
3	T	150	ASP	N-CA-C	-6.88	104.89	113.28
3	T	82	ASN	CA-CB-CG	-6.86	105.74	112.60
2	H	229	THR	N-CA-C	-6.77	100.27	110.28
3	T	61	ASP	CA-CB-CG	-6.75	105.85	112.60
3	T	157	TYR	N-CA-C	-6.74	92.12	111.00
2	H	101	HIS	CA-CB-CG	-6.72	107.08	113.80
3	T	72	LEU	N-CA-C	-6.69	98.86	109.23
3	T	99	GLU	N-CA-C	-6.66	101.56	110.55
2	H	29	TRP	NE1-CE2-CZ2	6.66	140.09	130.10
2	H	79	ASP	N-CA-C	6.61	124.89	110.80
2	H	165	THR	CA-CB-OG1	-6.44	99.94	109.60
2	H	207	TRP	NE1-CE2-CZ2	6.44	139.76	130.10
2	H	129(B)	ARG	N-CA-C	6.43	121.33	112.90
3	T	70	THR	N-CA-C	-6.36	98.54	108.90
2	H	43	GLY	N-CA-C	-6.35	102.55	112.58
2	H	29	TRP	CG-CD1-NE1	-6.34	101.96	110.20
2	H	214	SER	N-CA-C	6.34	120.61	112.12
2	H	143	GLN	OE1-CD-NE2	-6.33	116.27	122.60
3	T	136	ARG	N-CA-C	-6.33	99.42	109.23
1	L	71	PHE	CA-CB-CG	-6.32	107.48	113.80
1	L	92	VAL	N-CA-C	-6.31	105.45	113.22
2	H	134	ARG	CA-CB-CG	-6.30	101.49	114.10
1	L	125	VAL	N-CA-C	6.30	117.41	112.12
2	H	237	TRP	NE1-CE2-CZ2	6.27	139.51	130.10
2	H	61	TRP	CG-CD1-NE1	-6.23	102.11	110.20
3	T	7	VAL	N-CA-C	-6.21	99.30	108.99
3	T	190	GLN	N-CA-C	-6.21	99.24	108.99
3	T	53	THR	N-CA-C	-6.03	105.88	113.72
2	H	59	PHE	CA-CB-CG	-5.97	107.83	113.80
2	H	125	GLU	N-CA-C	-5.88	101.77	110.48
2	H	251	LEU	N-CA-C	-5.88	99.65	109.24
3	T	25	TRP	CG-CD1-NE1	-5.88	102.55	110.20
1	L	49	GLN	OE1-CD-NE2	-5.84	116.76	122.60
2	H	207	TRP	CG-CD1-NE1	-5.81	102.65	110.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	138[B]	VAL	N-CA-C	-5.72	100.05	108.85
2	H	138[A]	VAL	N-CA-C	-5.72	100.05	108.85
2	H	224	HIS	CE1-NE2-CD2	-5.71	103.29	109.00
3	T	45	TRP	NE1-CE2-CZ2	5.70	138.65	130.10
2	H	84	ARG	N-CA-C	-5.70	102.09	110.52
3	T	14	TRP	NE1-CE2-CZ2	5.68	138.63	130.10
3	T	45	TRP	CG-CD1-NE1	-5.65	102.86	110.20
2	H	76	HIS	CE1-NE2-CD2	-5.64	103.36	109.00
3	T	35	THR	N-CA-C	-5.63	100.50	109.23
2	H	141	TRP	NE1-CE2-CZ2	5.63	138.55	130.10
2	H	237	TRP	CG-CD1-NE1	-5.60	102.92	110.20
3	T	18	ASN	CA-CB-CG	-5.57	107.03	112.60
3	T	31	ASN	OD1-CG-ND2	-5.53	117.07	122.60
1	L	104	ASP	N-CA-C	-5.52	101.28	110.17
2	H	30	GLN	OE1-CD-NE2	-5.52	117.08	122.60
3	T	21	THR	N-CA-C	5.52	117.51	108.52
2	H	69	GLY	N-CA-C	-5.51	108.13	115.40
2	H	215	TRP	CG-CD1-NE1	-5.50	103.04	110.20
2	H	51	TRP	CG-CD1-NE1	-5.49	103.06	110.20
3	T	191	ALA	N-CA-C	-5.46	100.80	109.76
2	H	29	TRP	NE1-CE2-CD2	-5.45	100.32	107.40
2	H	217	GLN	N-CA-C	-5.43	99.67	108.41
2	H	159	ASN	OD1-CG-ND2	-5.39	117.21	122.60
1	L	62	LYS	N-CA-C	-5.37	99.69	108.76
2	H	29	TRP	CD1-NE1-CE2	5.33	118.50	108.90
2	H	206	THR	CA-CB-OG1	-5.32	101.62	109.60
3	T	14	TRP	CG-CD1-NE1	-5.30	103.31	110.20
2	H	134	ARG	N-CA-C	5.28	117.84	111.40
3	T	44	ASP	N-CA-C	-5.23	103.49	110.55
2	H	151	THR	N-CA-C	-5.23	103.98	110.41
1	L	93	ASN	OD1-CG-ND2	-5.22	117.38	122.60
3	T	23	LEU	N-CA-C	-5.22	101.21	109.76
2	H	117	HIS	CE1-NE2-CD2	-5.21	103.78	109.00
1	L	91	CYS	N-CA-C	5.20	119.25	113.01
2	H	141	TRP	CG-CD1-NE1	-5.20	103.44	110.20
3	T	193	ILE	N-CA-C	-5.19	97.66	108.88
2	H	61	TRP	CD1-NE1-CE2	5.18	118.23	108.90
1	L	130	THR	N-CA-C	-5.17	105.65	112.94
3	T	15	LYS	N-CA-C	-5.17	98.97	108.02
2	H	56	ALA	N-CA-C	5.17	117.81	111.82
1	L	88	GLN	CB-CG-CD	-5.16	103.83	112.60
3	T	95	GLU	N-CA-C	-5.14	100.92	108.99

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	237	TRP	NE1-CE2-CD2	-5.13	100.73	107.40
2	H	61	TRP	NE1-CE2-CZ2	5.12	137.78	130.10
3	T	41	LYS	N-CA-C	5.10	118.28	111.75
2	H	29	TRP	CG-CD2-CE3	-5.09	128.81	133.90
3	T	138	ASN	OD1-CG-ND2	-5.06	117.54	122.60
1	L	97	GLY	N-CA-C	-5.05	108.31	115.43
3	T	188	SER	CA-CB-OG	-5.05	101.00	111.10
2	H	207	TRP	NE1-CE2-CD2	-5.04	100.85	107.40
2	H	235	ILE	N-CA-CB	-5.04	104.92	110.51
2	H	247	ARG	N-CA-C	-5.04	101.36	109.82
3	T	45	TRP	CD1-NE1-CE2	5.01	117.91	108.90
2	H	183	ALA	N-CA-C	-5.00	101.47	109.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	727	668	663	10	1
2	H	1988	1977	1967	27	0
3	T	1220	1186	1182	22	0
4	H	28	23	23	2	0
5	H	295	590	0	0	0
5	L	110	220	0	0	0
5	T	187	374	0	0	1
All	All	4555	5038	3835	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:195:SER:OG	4:H:258:CR9:O6'	2.07	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:236:GLU:H	2:H:236:GLU:CD	2.02	0.67
1:L:61:CYS:SG	1:L:68:TYR:HB2	2.36	0.65
1:L:71:PHE:CE2	3:T:131:ARG:HD3	2.31	0.65
2:H:143:GLN:HE22	2:H:147:ARG:H	1.50	0.59
2:H:136:SER:CB	2:H:199:HIS:CE1	2.90	0.55
2:H:136:SER:HB2	2:H:199:HIS:CE1	2.44	0.53
3:T:155:LEU:O	3:T:188:SER:N	2.43	0.51
2:H:228:TYR:N	2:H:228:TYR:CD1	2.78	0.51
3:T:87:GLY:O	3:T:89:ALA:N	2.43	0.51
3:T:51:TYR:CE1	3:T:83:VAL:HG13	2.46	0.49
2:H:50:ILE:HG13	2:H:51:TRP:CD1	2.48	0.48
2:H:135:PHE:HB3	2:H:159:ASN:ND2	2.28	0.48
2:H:124:PRO:O	2:H:235:ILE:HD13	2.13	0.47
1:L:121[B]:LEU:HD11	1:L:128:THR:HB	1.95	0.47
3:T:194:PRO:HA	3:T:200:ARG:NH1	2.28	0.47
4:H:258:CR9:O6'	4:H:258:CR9:N3	2.46	0.47
3:T:108:LEU:HB2	3:T:202:SER:HB3	1.96	0.46
1:L:62:LYS:O	1:L:68:TYR:HA	2.16	0.46
2:H:245:GLU:CD	2:H:245:GLU:N	2.74	0.45
3:T:86:THR:OG1	3:T:90:GLY:C	2.60	0.45
2:H:45:THR:OG1	2:H:198:PRO:HB3	2.17	0.45
2:H:165:THR:O	2:H:166:GLN:C	2.60	0.44
1:L:71:PHE:CE2	3:T:131:ARG:CD	3.00	0.44
2:H:143:GLN:HE22	2:H:147:ARG:N	2.14	0.44
2:H:27:CYS:N	2:H:28:PRO:CD	2.81	0.43
2:H:245:GLU:O	2:H:247:ARG:CD	2.66	0.43
3:T:45:TRP:CE2	3:T:74:ARG:NH2	2.86	0.43
3:T:150:ASP:O	3:T:193:ILE:HA	2.19	0.43
1:L:65:LEU:HD11	3:T:203:THR:HG21	2.01	0.43
2:H:91:PRO:O	2:H:253:ARG:NH2	2.52	0.43
3:T:136:ARG:O	3:T:137:ASN:C	2.61	0.43
2:H:48:ASN:HD21	2:H:51:TRP:CD1	2.37	0.43
3:T:51:TYR:CD1	3:T:83:VAL:HG22	2.53	0.42
3:T:154:THR:O	3:T:189:VAL:HA	2.19	0.42
3:T:63:ILE:HD12	3:T:63:ILE:C	2.44	0.42
2:H:92:SER:OG	2:H:255:PRO:HA	2.19	0.42
1:L:90:ILE:HD12	1:L:92:VAL:HG12	2.02	0.42
3:T:101:THR:O	3:T:102:PRO:C	2.63	0.42
2:H:161:PRO:O	2:H:183:ALA:HA	2.20	0.41
3:T:59:LEU:O	3:T:60:THR:C	2.63	0.41
3:T:135:ARG:HA	3:T:139:THR:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:127:THR:O	2:H:129(B):ARG:HB3	2.21	0.41
1:L:61:CYS:SG	1:L:68:TYR:CB	3.06	0.41
2:H:245:GLU:O	2:H:247:ARG:HD3	2.19	0.41
3:T:144:ARG:HD2	3:T:148:GLY:O	2.20	0.41
1:L:129:PRO:HB3	1:L:134:PRO:HB3	2.02	0.41
2:H:76:HIS:ND1	2:H:77:ASP:O	2.54	0.41
2:H:121:LEU:HD11	2:H:207:TRP:HB3	2.03	0.41
2:H:166:GLN:HB2	3:T:94:TYR:OH	2.21	0.41
2:H:239:GLN:O	2:H:243:ARG:HG3	2.21	0.41
3:T:51:TYR:CD1	3:T:83:VAL:CG2	3.05	0.40
3:T:142:SER:O	3:T:145:ASP:N	2.54	0.40
2:H:215:TRP:CE2	2:H:227:VAL:HG11	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:118:TYR:O	5:T:303:HOH:H1[2_646]	1.53	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	96/152 (63%)	81 (84%)	9 (9%)	6 (6%)	1	0
2	H	255/254 (100%)	228 (89%)	23 (9%)	4 (2%)	7	2
3	T	146/218 (67%)	125 (86%)	17 (12%)	4 (3%)	4	0
All	All	497/624 (80%)	434 (87%)	49 (10%)	14 (3%)	3	0

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	50	CYS
2	H	79	ASP
3	T	88	SER
3	T	148	GLY
1	L	57	ASN
2	H	49	THR
3	T	156	TYR
1	L	66	GLN
1	L	51	ALA
2	H	170(I)	PRO
3	T	137	ASN
1	L	81	CYS
2	H	255	PRO
1	L	74	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	85/132 (64%)	79 (93%)	6 (7%)	13	7
2	H	219/216 (101%)	210 (96%)	9 (4%)	27	21
3	T	140/199 (70%)	132 (94%)	8 (6%)	18	11
All	All	444/547 (81%)	421 (95%)	23 (5%)	20	14

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	49	GLN
1	L	65	LEU
1	L	83	THR
1	L	88	GLN
1	L	92	VAL
1	L	103	SER
2	H	68	LEU
2	H	75	GLU
2	H	84	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	92	SER
2	H	159	ASN
2	H	195	SER
2	H	233	GLN
2	H	236	GLU
2	H	255	PRO
3	T	28	LYS
3	T	53	THR
3	T	83	VAL
3	T	85	SER
3	T	86	THR
3	T	131	ARG
3	T	143	LEU
3	T	195	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	88	GLN
2	H	30	GLN
2	H	87	GLN
2	H	100	ASN
2	H	117	HIS
2	H	143	GLN
2	H	159	ASN
2	H	170(A)	GLN
2	H	217	GLN
2	H	233	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CR9	H	258	-	31,31,31	2.34	9 (29%)	43,45,45	1.70	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CR9	H	258	-	2/2/3/3	1/11/23/23	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	258	CR9	C1-C7	-5.89	1.39	1.48
4	H	258	CR9	C4-N3	-5.62	1.28	1.38
4	H	258	CR9	C1-C2	4.17	1.44	1.38
4	H	258	CR9	C1'-C8	-4.15	1.40	1.47
4	H	258	CR9	C4-C5	-3.53	1.35	1.40
4	H	258	CR9	C5-N4	-3.43	1.32	1.39
4	H	258	CR9	C8-N3	-3.06	1.30	1.36
4	H	258	CR9	C4B-C5B	2.85	1.60	1.53
4	H	258	CR9	C5B-C6B	2.66	1.57	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	258	CR9	C5-N4-C8	-6.24	97.89	105.17
4	H	258	CR9	C1'-C6'-C5'	3.33	122.28	119.56
4	H	258	CR9	C4-C5-N4	3.05	113.60	109.42
4	H	258	CR9	N3-C8-N4	2.73	117.09	112.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	258	CR9	C1A-C1B-C6B	-2.69	108.11	112.62
4	H	258	CR9	C6-C5-N4	-2.13	124.73	129.31

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	H	258	CR9	C6B
4	H	258	CR9	C1B

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	258	CR9	C6-C1-C7-N1

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	258	CR9	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	L	96/152 (63%)	-0.16	5 (5%)	33 33	1, 30, 54, 66	19 (19%)
2	H	254/254 (100%)	-0.50	8 (3%)	51 52	1, 25, 54, 84	24 (9%)
3	T	151/218 (69%)	0.02	10 (6%)	24 24	9, 34, 73, 89	21 (13%)
All	All	501/624 (80%)	-0.28	23 (4%)	37 37	1, 28, 64, 89	64 (12%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	170(F)	GLY	7.8
3	T	157	TYR	6.9
3	T	137	ASN	6.8
3	T	188	SER	5.5
1	L	48	ASP	4.8
1	L	66	GLN	4.8
3	T	109	GLY	4.6
3	T	156	TYR	3.9
2	H	170(H)	SER	3.8
1	L	65	LEU	3.7
3	T	87	GLY	3.6
3	T	138	ASN	3.2
2	H	170(G)	ASP	3.0
2	H	170(D)	LYS	3.0
2	H	60(D)	ASN	2.9
1	L	51	ALA	2.8
2	H	170(E)	VAL	2.7
2	H	248	PRO	2.6
1	L	50	CYS	2.5
3	T	203	THR	2.3
3	T	129	ASP	2.3
3	T	81	GLY	2.3
2	H	62	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CR9	H	258	28/28	0.93	0.06	22,44,47,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CR9 H 258:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.