



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:26 PM UTC

PDB ID : 2O54 / pdb_00002o54
Title : Structure of E. coli topoisomerase III in complex with an 8-base single stranded oligonucleotide. Frozen in glycerol at pH 7.0
Authors : Changela, A.; Digate, R.J.; Mondragon, A.
Deposited on : 2006-12-05
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	8	
1	D	8	
2	A	659	
2	B	659	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	7	Total	C	N	O	P	0	0	0
			138	67	26	39	6			
1	D	8	Total	C	N	O	P	0	0	0
			158	77	28	46	7			

- Molecule 2 is a protein called DNA topoisomerase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	630	Total	C	N	O	S	0	0	0
			5013	3165	912	916	20			
2	B	631	Total	C	N	O	S	0	0	0
			5016	3168	913	915	20			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	654	HIS	-	expression tag	UNP P14294
A	655	HIS	-	expression tag	UNP P14294
A	656	HIS	-	expression tag	UNP P14294
A	657	HIS	-	expression tag	UNP P14294
A	658	HIS	-	expression tag	UNP P14294
A	659	HIS	-	expression tag	UNP P14294
B	654	HIS	-	expression tag	UNP P14294
B	655	HIS	-	expression tag	UNP P14294
B	656	HIS	-	expression tag	UNP P14294
B	657	HIS	-	expression tag	UNP P14294
B	658	HIS	-	expression tag	UNP P14294
B	659	HIS	-	expression tag	UNP P14294

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is ACETIC ACID (CCD ID: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		
5	D	10	Total	O	0	0
			10	10		
5	A	79	Total	O	0	0
			79	79		
5	B	117	Total	O	0	0
			117	117		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'

Chain C: 




- Molecule 1: 5'-D(*CP*GP*CP*AP*AP*CP*TP*T)-3'

Chain D: 




- Molecule 2: DNA topoisomerase 3


Chain A: 



- Molecule 2: DNA topoisomerase 3

Chain B: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.83Å 101.83Å 452.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 2.50 29.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.7 (29.10-2.50) 97.7 (29.10-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.222 , 0.269 0.223 , 0.264	Depositor DCC
R_{free} test set	4104 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	45.2	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10538	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.50	0/154	1.28	0/235
1	D	0.48	0/176	1.25	0/269
2	A	0.61	1/5123 (0.0%)	0.87	5/6945 (0.1%)
2	B	0.65	0/5127	0.87	3/6952 (0.0%)
All	All	0.62	1/10580 (0.0%)	0.89	8/14401 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	156	VAL	CA-CB	5.31	1.56	1.54

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	365	ASP	CA-C-N	5.68	125.39	119.82
2	B	365	ASP	C-N-CA	5.68	125.39	119.82
2	B	435	VAL	N-CA-C	5.56	116.12	108.12
2	A	288	ALA	CA-C-N	-5.28	114.80	120.03
2	A	288	ALA	C-N-CA	-5.28	114.80	120.03
2	A	454	GLU	N-CA-C	5.25	119.17	112.87
2	A	546	ARG	N-CA-C	-5.21	106.70	113.16
2	A	222	PHE	N-CA-C	5.19	117.91	109.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	138	0	80	2	0
1	D	158	0	92	1	0
2	A	5013	0	5014	53	0
2	B	5016	0	5020	41	0
3	A	2	0	0	0	0
4	B	4	0	3	0	0
5	A	79	0	0	1	0
5	B	117	0	0	0	0
5	C	1	0	0	0	0
5	D	10	0	0	0	0
All	All	10538	0	10209	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:396:THR:OG1	2:A:399:GLU:HG3	1.78	0.83
2:A:588:TYR:HB3	2:A:656:HIS:HE1	1.47	0.80
2:A:197:ARG:HD2	2:A:574:TRP:CZ3	2.17	0.79
2:A:588:TYR:HB3	2:A:656:HIS:CE1	2.23	0.74
2:A:169:ASP:OD2	2:A:197:ARG:HG2	1.89	0.72
2:A:520:ASP:HB3	2:A:525:GLU:OE2	1.89	0.72
2:B:525:GLU:OE2	2:B:525:GLU:HA	1.91	0.70
2:B:219:LYS:NZ	2:B:251:ASP:OD1	2.25	0.69
2:A:646:GLU:HB2	2:A:649:SER:HB3	1.77	0.67
2:A:187:ALA:HB2	2:A:611:VAL:HG13	1.76	0.66
2:B:87:VAL:HG22	2:B:90:ARG:HH21	1.61	0.66
2:B:259:ARG:NH1	2:B:263:GLU:OE2	2.30	0.65
2:B:92:LEU:HG	2:B:121:LEU:HD13	1.77	0.64
2:A:197:ARG:CD	2:A:574:TRP:CZ3	2.81	0.64
2:A:21:LYS:HG3	2:A:22:PRO:HA	1.81	0.62
2:A:126:ARG:HH12	2:A:152:ASN:HD22	1.50	0.59
2:A:515:ILE:HD13	2:A:576:SER:HB2	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:98:ILE:HB	2:B:129:VAL:HG22	1.86	0.58
2:A:169:ASP:OD2	2:A:197:ARG:CG	2.51	0.58
1:D:7:DT:H5''	2:B:8:LYS:HE2	1.86	0.58
2:A:503:ILE:HG23	2:A:516:LEU:HD13	1.87	0.57
2:A:577:VAL:O	2:A:581:ILE:HG13	2.04	0.57
2:A:169:ASP:OD2	2:A:197:ARG:HD3	2.04	0.56
2:B:106:ARG:HG3	2:B:162:ALA:HB2	1.87	0.56
2:B:221:PHE:CD1	2:B:255:ARG:HB3	2.41	0.56
2:B:150:ARG:HH11	2:B:150:ARG:HG2	1.70	0.55
2:A:58:TYR:O	2:A:178:ARG:NH2	2.26	0.55
2:B:427:LEU:HD21	2:B:473:LEU:HD12	1.89	0.55
2:B:583:GLU:O	2:B:584:LYS:HB2	2.08	0.54
2:B:72:LYS:NZ	2:B:649:SER:HB2	2.23	0.54
2:A:1:MET:HG2	2:A:35:GLY:O	2.08	0.54
2:B:7:GLU:HB2	2:B:102:GLY:HA2	1.90	0.54
2:A:419:VAL:HB	2:A:443:GLU:HB2	1.89	0.53
2:A:453:LYS:O	2:A:457:GLU:HG2	2.09	0.52
1:C:6:DC:H3'	1:C:7:DT:H5'	1.90	0.52
2:A:176:MET:HA	2:A:176:MET:HE2	1.92	0.52
2:B:277:THR:O	2:B:469:LYS:HD2	2.10	0.52
2:A:520:ASP:CB	2:A:525:GLU:OE2	2.58	0.52
2:A:106:ARG:HB2	2:A:162:ALA:HB2	1.92	0.51
2:B:177:THR:HG23	2:B:194:SER:HA	1.92	0.51
2:B:455:ARG:HD2	2:B:456:ASP:OD2	2.11	0.51
1:C:2:DG:N7	2:A:178:ARG:HD2	2.26	0.51
2:B:100:HIS:CE1	2:B:112:VAL:HB	2.45	0.50
2:A:361:GLN:HG2	2:A:449:LEU:HD21	1.94	0.50
2:B:46:LEU:HD12	2:B:114:GLU:HG3	1.93	0.50
2:B:574:TRP:CZ2	2:B:595:LEU:CD1	2.96	0.49
2:A:83:LYS:O	2:A:87:VAL:HG23	2.13	0.48
2:A:197:ARG:HD2	2:A:574:TRP:CE3	2.49	0.48
2:B:187:ALA:HB2	2:B:611:VAL:HB	1.94	0.48
2:B:202:VAL:HG21	2:B:500:MET:HE3	1.95	0.48
2:B:88:ILE:O	2:B:92:LEU:HB2	2.13	0.48
2:A:276:VAL:HG21	2:A:467:VAL:HG13	1.96	0.48
2:A:615:ARG:O	2:A:617:ILE:HG13	2.15	0.47
2:B:100:HIS:NE2	2:B:112:VAL:HB	2.29	0.47
2:A:509:ASP:HB3	2:A:512:LEU:HB2	1.95	0.47
2:B:327:THR:O	2:B:328:TYR:C	2.57	0.47
2:A:202:VAL:O	2:A:203:LEU:C	2.58	0.46
2:A:500:MET:HE2	2:A:500:MET:HB3	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:208:ARG:NH2	2:B:566:THR:O	2.48	0.46
2:B:507:VAL:CG2	2:B:568:PRO:HG2	2.45	0.46
2:B:232:ALA:HB3	2:B:234:GLU:OE1	2.15	0.46
2:B:72:LYS:HZ2	2:B:649:SER:HB2	1.81	0.46
2:A:240:TRP:O	2:A:242:PRO:HD3	2.16	0.45
2:B:229:VAL:HG12	2:B:474:LEU:HB3	1.98	0.45
2:A:169:ASP:OD2	2:A:197:ARG:CD	2.65	0.45
2:A:277:THR:O	2:A:469:LYS:HE2	2.17	0.45
2:B:525:GLU:OE2	2:B:525:GLU:CA	2.62	0.45
2:A:98:ILE:HB	2:A:129:VAL:HG22	1.99	0.45
2:B:7:GLU:HG3	2:B:111:LEU:HD11	1.98	0.44
2:B:444:ALA:O	2:B:447:ARG:HB2	2.17	0.44
2:B:247:GLU:N	2:B:248:PRO:HD2	2.32	0.44
2:B:574:TRP:CZ2	2:B:595:LEU:HD11	2.53	0.44
2:A:133:LEU:HD11	2:A:155:PHE:CZ	2.52	0.44
2:A:197:ARG:HH21	2:A:571:THR:HG23	1.83	0.44
2:A:268:ARG:HD3	5:A:861:HOH:O	2.17	0.44
2:A:291:PRO:HD3	2:A:413:GLN:CD	2.43	0.44
2:B:89:LYS:HD3	2:B:119:LEU:HD22	1.98	0.44
2:A:450:LEU:HB3	2:A:454:GLU:HG3	2.00	0.43
2:A:649:SER:C	2:A:651:ALA:H	2.26	0.43
2:A:588:TYR:CB	2:A:656:HIS:HE1	2.24	0.43
2:A:613:GLN:C	2:A:615:ARG:H	2.27	0.43
2:A:151:SER:O	2:A:154:GLU:HG2	2.20	0.42
2:B:252:GLU:CD	2:B:252:GLU:H	2.27	0.42
2:B:579:THR:O	2:B:583:GLU:HG2	2.20	0.42
2:A:133:LEU:HD11	2:A:155:PHE:HZ	1.85	0.41
2:A:117:ASP:HB3	2:A:648:GLY:HA2	2.00	0.41
2:A:1:MET:HE1	2:A:3:LEU:HB2	2.01	0.41
2:A:69:VAL:HB	2:B:593:GLN:OE1	2.20	0.41
2:A:265:VAL:O	2:A:269:ILE:HG12	2.21	0.41
2:A:88:ILE:O	2:A:92:LEU:HB2	2.20	0.41
2:B:6:ALA:HA	2:B:101:ALA:O	2.20	0.41
2:B:422:LYS:HA	2:B:438:ALA:O	2.21	0.41
2:A:276:VAL:HG21	2:A:467:VAL:CG1	2.52	0.40
2:A:428:ASP:HA	2:A:432:GLY:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	626/659 (95%)	591 (94%)	28 (4%)	7 (1%)	11	22
2	B	627/659 (95%)	608 (97%)	19 (3%)	0	100	100
All	All	1253/1318 (95%)	1199 (96%)	47 (4%)	7 (1%)	21	38

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	455	ARG
2	A	615	ARG
2	A	616	GLY
2	A	463	PRO
2	A	525	GLU
2	A	453	LYS
2	A	647	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	535/555 (96%)	516 (96%)	19 (4%)	31	58
2	B	535/555 (96%)	514 (96%)	21 (4%)	28	55
All	All	1070/1110 (96%)	1030 (96%)	40 (4%)	30	57

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	57	ARG
2	A	100	HIS
2	A	126	ARG
2	A	140	GLN
2	A	153	SER
2	A	158	LEU
2	A	178	ARG
2	A	251	ASP
2	A	280	ASN
2	A	458	GLU
2	A	462	THR
2	A	467	VAL
2	A	480	VAL
2	A	520	ASP
2	A	527	THR
2	A	586	CYS
2	A	611	VAL
2	A	617	ILE
2	A	652	ILE
2	B	18	VAL
2	B	38	VAL
2	B	48	GLN
2	B	80	SER
2	B	98	ILE
2	B	127	GLN
2	B	246	CYS
2	B	263	GLU
2	B	270	SER
2	B	277	THR
2	B	282	LYS
2	B	304	ARG
2	B	317	GLN
2	B	318	LYS
2	B	372	CYS
2	B	435	VAL
2	B	469	LYS
2	B	481	VAL
2	B	482	GLU
2	B	497	LEU
2	B	508	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	23	HIS
2	A	50	GLN
2	A	74	GLN
2	A	93	HIS
2	A	241	GLN
2	A	311	ASN
2	A	317	GLN
2	A	404	ASN
2	A	656	HIS
2	B	76	GLN
2	B	580	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACY	B	803	-	3,3,3	1.17	0	3,3,3	0.72	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	7/8 (87%)	-0.29	0 100 100	24, 28, 62, 71	0
1	D	8/8 (100%)	-1.21	0 100 100	17, 24, 36, 44	0
2	A	630/659 (95%)	0.31	58 (9%) 14 13	14, 51, 104, 145	0
2	B	631/659 (95%)	-0.25	16 (2%) 58 54	15, 35, 79, 138	0
All	All	1276/1334 (95%)	0.02	74 (5%) 29 25	14, 42, 95, 145	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	647	VAL	8.1
2	B	658	HIS	6.7
2	A	652	ILE	5.7
2	A	617	ILE	5.4
2	B	648	GLY	5.1
2	B	656	HIS	5.0
2	A	455	ARG	5.0
2	A	658	HIS	4.7
2	A	217	VAL	4.6
2	B	651	ALA	4.6
2	B	618	VAL	4.6
2	A	452	SER	4.5
2	A	651	ALA	4.4
2	A	454	GLU	4.4
2	A	656	HIS	4.2
2	B	650	GLY	4.1
2	B	652	ILE	4.1
2	A	650	GLY	4.0
2	A	216	PHE	4.0
2	A	218	ALA	4.0
2	A	648	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	654	HIS	3.8
2	A	523	GLY	3.7
2	B	653	ALA	3.6
2	A	249	TYR	3.6
2	B	654	HIS	3.6
2	B	657	HIS	3.6
2	A	657	HIS	3.5
2	A	544	LYS	3.5
2	A	655	HIS	2.9
2	A	524	THR	2.9
2	A	257	LEU	2.9
2	A	649	SER	2.9
2	A	254	GLY	2.9
2	A	616	GLY	2.9
2	A	646	GLU	2.9
2	B	655	HIS	2.8
2	A	481	VAL	2.8
2	A	22	PRO	2.7
2	A	63	LEU	2.7
2	A	458	GLU	2.7
2	A	213	ILE	2.6
2	A	512	LEU	2.6
2	B	649	SER	2.6
2	A	507	VAL	2.6
2	A	219	LYS	2.6
2	B	123	PRO	2.6
2	A	480	VAL	2.6
2	A	245	ALA	2.6
2	A	547	TYR	2.6
2	A	123	PRO	2.5
2	A	463	PRO	2.5
2	B	619	ALA	2.5
2	A	614	PHE	2.4
2	A	453	LYS	2.4
2	A	264	HIS	2.4
2	A	520	ASP	2.4
2	A	462	THR	2.4
2	A	122	ALA	2.4
2	A	456	ASP	2.3
2	A	504	ALA	2.3
2	A	258	HIS	2.3
2	A	256	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	260	PRO	2.3
2	B	122	ALA	2.2
2	A	473	LEU	2.2
2	B	620	PRO	2.2
2	A	611	VAL	2.2
2	A	93	HIS	2.1
2	A	252	GLU	2.1
2	A	459	ASN	2.1
2	A	519	THR	2.1
2	A	526	ALA	2.0
2	A	240	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ACY	B	803	4/4	0.82	0.12	23,23,24,25	0
3	CL	A	800	1/1	0.98	0.03	23,23,23,23	0
3	CL	A	801	1/1	0.99	0.03	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.