



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 12:54 PM UTC

PDB ID : 5O56 / pdb_00005o56
Title : Glycogen Phosphorylase b in complex with 29b
Authors : Solovou, T.G.A.; Kyriakis, E.; Stravodimos, G.A.; Kantsadi, A.L.; Chatzileontiadou, D.S.; Skamnaki, V.T.; Leonidas, D.D.
Deposited on : 2017-05-31
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	843	 81% 13% . .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen phosphorylase, muscle form.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	810	Total	C	N	O	S	0	3	0
			6611	4216	1167	1199	29			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is (2 {R},3 {S},4 {R},5 {R},6 {R})-5-azanyl-2-(hydroxymethyl)-6-(3-naphthale n-2-yl-1 {H}-1,2,4-triazol-5-yl)oxane-3,4-diol (CCD ID: 9L8) (formula: $C_{18}H_{20}N_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			26	18	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	199	Total	O	0	0
			199	199		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycogen phosphorylase, muscle form



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	128.55Å 128.55Å 116.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	13.56 – 2.45 13.56 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.5 (13.56-2.45) 97.5 (13.56-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.44Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.164 , 0.229 0.165 , 0.229	Depositor DCC
R_{free} test set	1776 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6851	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, 9L8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.40	17/6769 (0.3%)	1.39	33/9160 (0.4%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	701	GLU	N-CA	-6.55	1.38	1.46
1	A	64	VAL	CA-CB	6.00	1.61	1.54
1	A	461	GLU	C-O	-5.88	1.17	1.24
1	A	655	LYS	C-O	-5.76	1.15	1.23
1	A	249	PRO	N-CA	5.74	1.53	1.47
1	A	465	LYS	C-O	-5.68	1.16	1.24
1	A	162	GLU	CA-CB	-5.48	1.44	1.53
1	A	378	THR	CB-CG2	5.36	1.70	1.52
1	A	284	ASN	CA-C	5.34	1.59	1.53
1	A	674	SER	N-CA	-5.32	1.39	1.46
1	A	283	ASP	C-O	5.21	1.30	1.24
1	A	576	GLN	CA-C	5.09	1.59	1.52
1	A	211	GLN	CA-C	5.09	1.59	1.52
1	A	240	THR	CA-C	-5.08	1.46	1.52
1	A	60[A]	ARG	C-O	-5.04	1.18	1.24
1	A	60[B]	ARG	C-O	-5.04	1.18	1.24
1	A	335	ILE	C-O	5.04	1.29	1.24

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	325	ASN	N-CA-C	9.41	124.29	108.20
1	A	507	ILE	N-CA-C	7.75	119.53	112.17
1	A	211	GLN	N-CA-C	7.51	120.42	111.33
1	A	47	THR	CA-C-N	-7.36	110.93	119.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	THR	C-N-CA	-7.36	110.93	119.47
1	A	487	THR	CA-C-N	-7.22	112.44	120.45
1	A	487	THR	C-N-CA	-7.22	112.44	120.45
1	A	467	ILE	CB-CA-C	7.12	121.36	112.04
1	A	556	HIS	N-CA-C	6.95	120.08	108.20
1	A	789	ALA	N-CA-C	-6.51	104.11	111.14
1	A	278	VAL	N-CA-C	6.26	116.84	108.27
1	A	551	ARG	N-CA-C	6.15	117.67	110.97
1	A	20	GLY	N-CA-C	6.13	118.53	111.36
1	A	250	ASN	N-CA-C	6.00	120.33	112.89
1	A	91	MET	N-CA-C	5.80	118.83	111.69
1	A	499	LEU	N-CA-C	-5.63	104.53	111.40
1	A	81	ARG	CG-CD-NE	5.62	124.38	112.00
1	A	14	SER	N-CA-C	5.60	118.57	111.69
1	A	647	ASN	N-CA-C	5.48	118.95	111.39
1	A	140	ALA	N-CA-C	-5.42	105.29	111.14
1	A	346	ILE	CA-C-N	-5.41	114.10	119.56
1	A	346	ILE	C-N-CA	-5.41	114.10	119.56
1	A	828	GLU	CB-CA-C	5.31	115.91	109.85
1	A	394	THR	CB-CA-C	-5.28	101.04	110.70
1	A	230	VAL	CB-CA-C	-5.26	106.08	111.08
1	A	705	GLU	N-CA-C	5.26	117.76	111.71
1	A	677	GLY	N-CA-C	-5.18	106.51	112.83
1	A	611	PRO	N-CA-C	5.18	121.47	113.75
1	A	362	ASP	N-CA-C	5.10	116.84	111.28
1	A	394	THR	N-CA-C	-5.07	105.45	111.69
1	A	763	ASN	N-CA-C	5.06	116.79	111.28
1	A	797	TRP	N-CA-C	-5.01	105.71	111.07
1	A	832	GLN	N-CA-C	5.01	117.42	109.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6611	0	6560	43	0
2	A	15	0	7	1	0
3	A	26	0	0	0	0
4	A	199	0	0	3	0
All	All	6851	0	6567	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:HIS:HD2	1:A:218:THR:OG1	1.59	0.85
1:A:549:LEU:HD12	1:A:557:ILE:HD13	1.60	0.81
1:A:539:GLN:OE1	4:A:1001:HOH:O	2.15	0.64
1:A:144:LEU:HD23	1:A:147:MET:HE1	1.84	0.60
1:A:379:VAL:HG22	4:A:1130:HOH:O	2.01	0.59
1:A:201:HIS:CD2	1:A:218:THR:OG1	2.50	0.59
1:A:546:ALA:HA	1:A:557:ILE:HD11	1.86	0.58
1:A:152:LEU:HD22	1:A:827:VAL:HG11	1.87	0.56
1:A:67:TRP:CH2	1:A:229:PRO:HD3	2.43	0.54
1:A:764:MET:HA	1:A:768[B]:HIS:CD2	2.43	0.53
1:A:122:LEU:HA	1:A:125:ILE:HD12	1.92	0.52
1:A:152:LEU:HD22	1:A:827:VAL:CG1	2.40	0.52
1:A:128:ASP:OD1	1:A:651:SER:HB3	2.11	0.51
1:A:283:ASP:OD2	1:A:571[A]:HIS:HE1	1.94	0.50
1:A:144:LEU:HA	1:A:147:MET:HE3	1.94	0.50
1:A:284:ASN:ND2	1:A:380:LEU:HD12	2.26	0.50
1:A:571[A]:HIS:HB2	1:A:574:LYS:HD2	1.95	0.48
1:A:170:ILE:HG12	1:A:646:GLU:HG3	1.96	0.48
1:A:591:LYS:HE3	1:A:633:ASP:OD2	2.15	0.47
1:A:611:PRO:O	1:A:617:LYS:HE3	2.15	0.47
1:A:271:LEU:N	1:A:271:LEU:HD23	2.31	0.46
1:A:568:LYS:HZ1	1:A:665:GLN:HE22	1.62	0.46
1:A:198:LEU:HD13	1:A:305:GLN:HB2	1.98	0.46
1:A:13:ILE:HD12	1:A:501:GLU:HB2	1.97	0.46
1:A:249:PRO:O	1:A:252:PHE:HB2	2.16	0.45
1:A:708:PHE:CD2	1:A:710:ILE:HD11	2.52	0.45
1:A:15:VAL:HG12	1:A:509:GLU:HG2	2.00	0.44
1:A:568:LYS:NZ	1:A:665:GLN:HE22	2.15	0.44
1:A:144:LEU:HD23	1:A:147:MET:CE	2.46	0.44
1:A:60[A]:ARG:NE	4:A:1003:HOH:O	2.41	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:730:GLU:O	1:A:734:ARG:HD2	2.20	0.42
1:A:745:LEU:HD23	1:A:745:LEU:HA	1.85	0.42
1:A:143:PHE:C	1:A:147:MET:HE2	2.44	0.42
1:A:549:LEU:HD23	1:A:549:LEU:HA	1.89	0.41
1:A:81:ARG:HG3	1:A:81:ARG:HH11	1.86	0.41
1:A:753:LYS:HB2	1:A:753:LYS:HE3	1.94	0.41
1:A:591:LYS:CE	1:A:633:ASP:OD2	2.69	0.41
1:A:618:MET:HB3	1:A:761:ILE:HD11	2.02	0.41
1:A:732:TYR:CZ	1:A:739:ARG:HG3	2.56	0.41
1:A:494:LEU:C	1:A:494:LEU:HD23	2.46	0.40
1:A:90:TYR:CE1	1:A:130:GLY:HA2	2.56	0.40
1:A:391:LEU:HD23	1:A:391:LEU:HA	1.93	0.40
1:A:680:LYS:NZ	2:A:901:PLP:O3	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	807/843 (96%)	772 (96%)	34 (4%)	1 (0%)	48 61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	554	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	704/732 (96%)	667 (95%)	37 (5%)	20	30

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	13	ILE
1	A	22	GLU
1	A	23	ASN
1	A	29	LYS
1	A	47	THR
1	A	48	PRO
1	A	87	LEU
1	A	169	LYS
1	A	210	SER
1	A	214	LYS
1	A	245	SER
1	A	253	ASN
1	A	271	LEU
1	A	284	ASN
1	A	313	SER
1	A	314	SER
1	A	358	ARG
1	A	384	LEU
1	A	467	ILE
1	A	513	SER
1	A	550	GLU
1	A	554	LYS
1	A	562	LEU
1	A	565	VAL
1	A	568	LYS
1	A	571[A]	HIS
1	A	571[B]	HIS
1	A	598	VAL
1	A	706	GLU
1	A	720	ARG
1	A	723	GLN
1	A	754	GLN
1	A	761	ILE
1	A	763	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	779	GLU
1	A	827	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	HIS
1	A	71	GLN
1	A	73	HIS
1	A	97	ASN
1	A	201	HIS
1	A	284	ASN
1	A	305	GLN
1	A	325	ASN
1	A	481	ASN
1	A	556	HIS
1	A	665	GLN
1	A	727	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	9L8	A	902	-	29,29,29	1.79	7 (24%)	34,42,42	1.75	7 (20%)
2	PLP	A	901	1	15,15,16	2.64	3 (20%)	21,22,23	1.97	7 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9L8	A	902	-	-	0/6/30/30	0/4/4/4
2	PLP	A	901	1	-	1/6/6/8	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	PLP	C3-C2	6.77	1.48	1.41
2	A	901	PLP	C5-C4	6.35	1.47	1.40
3	A	902	9L8	C6-C4	5.12	1.55	1.47
2	A	901	PLP	C3-C4	3.06	1.46	1.40
3	A	902	9L8	C10-C11	2.65	1.42	1.36
3	A	902	9L8	C7-C6	2.59	1.42	1.37
3	A	902	9L8	C2'-N2'	2.45	1.51	1.47
3	A	902	9L8	C13-C12	2.26	1.41	1.36
3	A	902	9L8	C14-C15	2.24	1.41	1.36
3	A	902	9L8	O3'-C3'	2.19	1.48	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	9L8	N3-C4-N5	-5.15	106.83	114.17
2	A	901	PLP	C6-C5-C4	3.97	121.35	118.10
2	A	901	PLP	C6-N1-C2	3.84	126.16	119.20
2	A	901	PLP	C5-C6-N1	-3.49	118.14	123.83
2	A	901	PLP	O4P-C5A-C5	-3.46	102.87	109.36
3	A	902	9L8	C4-N3-N2	3.35	109.14	102.97
3	A	902	9L8	C6-C4-N5	3.14	128.42	122.92
2	A	901	PLP	C4-C3-C2	-2.64	115.94	119.89
3	A	902	9L8	C1'-C1-N2	2.62	126.93	122.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	9L8	O4'-C4'-C3'	-2.54	104.39	110.38
2	A	901	PLP	C5A-C5-C6	-2.44	115.38	119.36
3	A	902	9L8	O5'-C1'-C2'	2.38	114.23	109.07
2	A	901	PLP	O3-C3-C2	2.18	122.10	117.58
3	A	902	9L8	O3'-C3'-C2'	2.02	113.84	110.22

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	901	PLP	C4-C5-C5A-O4P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	PLP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	810/843 (96%)	-0.85	3 (0%) 88 89	11, 26, 54, 90	3 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	THR	2.7
1	A	555	VAL	2.3
1	A	253	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9L8	A	902	26/26	0.97	0.04	13,17,19,20	0
2	PLP	A	901	15/16	0.98	0.05	12,13,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.