



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:45 PM UTC

PDB ID : 5O59 / pdb_00005o59
Title : Cellobiohydrolase Cel7A from *T. atroviride*
Authors : Borisova, A.S.; Stahlberg, J.; Hansson, H.
Deposited on : 2017-06-01
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	430	 99%
1	B	430	 99%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	14	0
			3272	2008	551	686	27			
1	B	430	Total	C	N	O	S	0	12	0
			3259	2000	549	683	27			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ni	0	0
			2	2		
3	B	2	Total	Ni	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 1-thio-beta-D-glucopyranose (CCD ID: GS1) (formula: $C_6H_{12}O_5S$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			11	6	4	1		
6	A	1	Total	C	O	S	0	0
			12	6	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			11	6	4	1		
6	A	1	Total	C	O	S	0	0
			12	6	5	1		
6	B	1	Total	C	O	S	0	0
			11	6	4	1		
6	B	1	Total	C	O	S	0	0
			12	6	5	1		
6	B	1	Total	C	O	S	0	0
			11	6	4	1		
6	B	1	Total	C	O	S	0	0
			12	6	5	1		

- Molecule 7 is alpha-D-glucopyranose (CCD ID: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			11	6	5		
7	A	1	Total	C	O	0	0
			11	6	5		
7	B	1	Total	C	O	0	0
			11	6	5		
7	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	434	Total	O	0	8
			442	442		
9	B	415	Total	O	0	5
			420	420		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucanase

Chain A:  99%



- Molecule 1: Glucanase

Chain B:  99%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.85Å 71.34Å 102.91Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	58.63 – 1.75 58.63 – 1.75	Depositor EDS
% Data completeness (in resolution range)	82.6 (58.63-1.75) 82.6 (58.63-1.75)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.156 , 0.190 0.165 , 0.197	Depositor DCC
R_{free} test set	3413 reflections (4.20%)	wwPDB-VP
Wilson B-factor (Å ²)	15.9	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 29.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.478 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7614	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NI, PEG, GOL, BTB, GS1, NAG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3339	0.74	0/4555
1	B	0.55	0/3326	0.73	0/4538
All	All	0.55	0/6665	0.73	0/9093

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3272	0	2993	4	0
1	B	3259	0	2977	6	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	6	0	8	0	0
4	B	12	0	16	0	0
5	A	14	0	18	4	0
5	B	14	0	18	6	0
6	A	46	0	40	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	46	0	40	0	0
7	A	22	0	20	0	0
7	B	22	0	20	0	0
8	B	7	0	10	1	0
9	A	442	0	0	0	0
9	B	420	0	0	0	0
All	All	7614	0	6186	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:507:BTB:H61	5:B:507:BTB:H11	1.15	1.12
5:B:507:BTB:H11	5:B:507:BTB:C6	1.97	0.94
1:B:249[B]:ASN:HD22	1:B:249[B]:ASN:C	1.76	0.91
1:B:249[B]:ASN:C	1:B:249[B]:ASN:ND2	2.31	0.88
1:B:249[B]:ASN:ND2	1:B:249[B]:ASN:O	2.13	0.80
5:B:507:BTB:H41	5:B:507:BTB:C8	2.18	0.72
1:A:206:HIS:CE1	5:A:605:BTB:H32	2.30	0.66
5:B:507:BTB:HO1	5:B:507:BTB:HO4	1.46	0.63
1:A:206:HIS:HE1	5:A:605:BTB:H32	1.70	0.54
5:B:507:BTB:H61	5:B:507:BTB:C1	2.10	0.50
5:A:605:BTB:H11	5:A:605:BTB:H52	1.51	0.48
5:A:605:BTB:H42	5:A:605:BTB:H71	1.65	0.46
1:B:426:ILE:HB	8:B:506:PEG:H11	1.98	0.46
1:B:144:LEU:C	1:B:144:LEU:HD23	2.40	0.46
5:B:507:BTB:H41	5:B:507:BTB:H82	1.95	0.45
1:A:177:PRO:HD2	1:A:208:SER:O	2.18	0.44
1:A:2:GLN:HB2	1:A:72:CYS:HB3	2.00	0.42
1:B:163:ALA:HB1	1:B:167:TYR:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/430 (103%)	432 (98%)	10 (2%)	0	100	100
1	B	440/430 (102%)	433 (98%)	7 (2%)	0	100	100
All	All	882/860 (103%)	865 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/344 (104%)	357 (100%)	1 (0%)	86	83
1	B	356/344 (104%)	355 (100%)	1 (0%)	86	83
All	All	714/688 (104%)	712 (100%)	2 (0%)	86	83

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	B	37	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	306	ASN
1	A	335	GLN
1	B	52	ASN
1	B	124	ASN
1	B	186	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	B	1	1	7,8,9	0.44	0	9,10,12	0.94	0
1	PCA	A	1	1	7,8,9	0.50	0	9,10,12	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	B	1	1	-	0/0/11/13	0/1/1/1
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 4 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GS1	B	511	-	10,11,12	0.97	0	13,15,17	0.98	0
6	GS1	B	512	-	11,12,12	0.62	0	15,17,17	0.68	0
4	GOL	B	505	-	5,5,5	0.53	0	5,5,5	0.67	0
7	GLC	A	611	-	11,11,12	1.46	3 (27%)	15,15,17	1.26	2 (13%)
2	NAG	A	601	1	14,14,15	0.30	0	17,19,21	0.83	0
5	BTB	B	507	3	13,13,13	0.98	0	7,16,16	1.18	1 (14%)
4	GOL	B	504	-	5,5,5	0.28	0	5,5,5	0.33	0
7	GLC	A	608	-	11,11,12	0.81	0	15,15,17	0.68	0
6	GS1	A	606	-	10,11,12	0.86	0	13,15,17	0.59	0
6	GS1	A	610	-	11,12,12	0.72	0	15,17,17	0.67	0
7	GLC	B	513	-	11,11,12	1.32	2 (18%)	15,15,17	1.43	2 (13%)
5	BTB	A	605	3	13,13,13	0.55	0	7,16,16	0.46	0
6	GS1	B	509	-	11,12,12	1.07	1 (9%)	15,17,17	0.61	0
7	GLC	B	510	-	11,11,12	0.80	0	15,15,17	0.68	0
4	GOL	A	604	-	5,5,5	0.53	0	5,5,5	0.62	0
6	GS1	B	508	-	10,11,12	1.04	1 (10%)	13,15,17	0.74	0
6	GS1	A	609	-	10,11,12	0.99	0	13,15,17	0.79	0
6	GS1	A	607	-	11,12,12	1.00	1 (9%)	15,17,17	0.62	0
8	PEG	B	506	-	6,6,6	0.45	0	5,5,5	0.29	0
2	NAG	B	501	1	14,14,15	0.37	0	17,19,21	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GS1	B	511	-	-	0/2/18/22	0/1/1/1
6	GS1	B	512	-	-	0/2/22/22	0/1/1/1
4	GOL	B	505	-	-	4/4/4/4	-
7	GLC	A	611	-	-	0/2/18/22	0/1/1/1
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
5	BTB	B	507	3	-	17/21/21/21	-
4	GOL	B	504	-	-	4/4/4/4	-
7	GLC	A	608	-	-	0/2/18/22	0/1/1/1
6	GS1	A	606	-	-	0/2/18/22	0/1/1/1
6	GS1	A	610	-	-	0/2/22/22	0/1/1/1
7	GLC	B	513	-	-	0/2/18/22	0/1/1/1
5	BTB	A	605	3	-	0/21/21/21	-
6	GS1	B	509	-	-	1/2/22/22	0/1/1/1
7	GLC	B	510	-	-	0/2/18/22	0/1/1/1
4	GOL	A	604	-	-	4/4/4/4	-
6	GS1	B	508	-	-	0/2/18/22	0/1/1/1
6	GS1	A	609	-	-	0/2/18/22	0/1/1/1
6	GS1	A	607	-	-	1/2/22/22	0/1/1/1
8	PEG	B	506	-	-	2/4/4/4	-
2	NAG	B	501	1	-	0/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	611	GLC	O5-C5	2.64	1.48	1.44
6	B	509	GS1	O5-C1	2.61	1.46	1.42
6	B	508	GS1	O5-C1	2.31	1.46	1.42
6	A	607	GS1	O5-C1	2.30	1.46	1.42
7	A	611	GLC	O1-C1	2.26	1.46	1.39
7	B	513	GLC	O5-C5	2.15	1.47	1.44
7	B	513	GLC	O1-C1	2.09	1.46	1.39
7	A	611	GLC	C1-C2	2.07	1.57	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	513	GLC	C1-O5-C5	3.61	120.56	113.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	611	GLC	C1-O5-C5	2.78	118.97	113.65
7	A	611	GLC	O5-C5-C6	2.58	110.97	106.83
5	B	507	BTB	C8-C7-N	2.29	120.54	111.59
7	B	513	GLC	O5-C1-C2	2.11	114.02	110.30

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	604	GOL	C1-C2-C3-O3
4	B	504	GOL	O1-C1-C2-C3
4	B	504	GOL	C1-C2-C3-O3
4	B	504	GOL	O2-C2-C3-O3
4	B	505	GOL	C1-C2-C3-O3
5	B	507	BTB	O1-C1-C2-C3
5	B	507	BTB	O1-C1-C2-C4
5	B	507	BTB	O1-C1-C2-N
5	B	507	BTB	C1-C2-C3-O3
5	B	507	BTB	C4-C2-C3-O3
5	B	507	BTB	N-C2-C3-O3
5	B	507	BTB	C1-C2-C4-O4
5	B	507	BTB	C3-C2-C4-O4
5	B	507	BTB	N-C2-C4-O4
5	B	507	BTB	C1-C2-N-C7
5	B	507	BTB	C3-C2-N-C7
5	B	507	BTB	C4-C2-N-C7
5	B	507	BTB	C8-C7-N-C2
5	B	507	BTB	N-C7-C8-O8
4	A	604	GOL	O1-C1-C2-C3
4	B	505	GOL	O1-C1-C2-C3
8	B	506	PEG	O2-C3-C4-O4
4	A	604	GOL	O2-C2-C3-O3
4	B	505	GOL	O2-C2-C3-O3
4	B	505	GOL	O1-C1-C2-O2
4	B	504	GOL	O1-C1-C2-O2
4	A	604	GOL	O1-C1-C2-O2
5	B	507	BTB	C6-C5-N-C2
5	B	507	BTB	C1-C2-N-C5
8	B	506	PEG	C1-C2-O2-C3
6	B	509	GS1	O5-C5-C6-O6
6	A	607	GS1	O5-C5-C6-O6
5	B	507	BTB	N-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	507	BTB	6	0
5	A	605	BTB	4	0
8	B	506	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/430 (99%)	-1.36	0 100 100	6, 16, 28, 50	16 (3%)
1	B	429/430 (99%)	-1.36	0 100 100	6, 16, 28, 54	14 (3%)
All	All	858/860 (99%)	-1.36	0 100 100	6, 16, 28, 54	30 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	1	8/9	0.99	0.03	19,21,21,22	0
1	PCA	B	1	8/9	0.99	0.02	20,20,21,21	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NI	B	502	1/1	0.97	0.04	57,57,57,57	0
4	GOL	B	504	6/6	0.97	0.09	55,58,59,59	0
8	PEG	B	506	7/7	0.97	0.07	49,52,52,53	0
7	GLC	B	513	11/12	0.98	0.04	30,31,32,33	0
7	GLC	A	611	11/12	0.98	0.04	30,31,32,32	0
2	NAG	B	501	14/15	0.99	0.03	20,21,22,23	0
4	GOL	B	505	6/6	0.99	0.03	27,29,30,30	0
5	BTB	A	605	14/14	0.99	0.06	42,49,53,55	0
5	BTB	B	507	14/14	0.99	0.06	37,54,60,62	0
6	GS1	A	606	11/12	0.99	0.03	23,24,24,25	0
6	GS1	A	607	12/12	0.99	0.03	24,26,26,27	0
6	GS1	A	609	11/12	0.99	0.03	24,24,27,29	0
6	GS1	A	610	12/12	0.99	0.03	19,20,21,21	0
6	GS1	B	508	11/12	0.99	0.03	23,24,24,24	0
6	GS1	B	509	12/12	0.99	0.03	25,27,28,29	0
6	GS1	B	511	11/12	0.99	0.03	23,24,27,29	0
7	GLC	A	608	11/12	0.99	0.03	24,24,25,25	0
3	NI	A	602	1/1	0.99	0.03	51,51,51,51	0
7	GLC	B	510	11/12	0.99	0.03	23,24,24,25	0
2	NAG	A	601	14/15	0.99	0.02	19,20,21,22	0
4	GOL	A	604	6/6	0.99	0.05	26,29,30,30	0
6	GS1	B	512	12/12	1.00	0.02	20,21,22,23	0
3	NI	A	603	1/1	1.00	0.03	33,33,33,33	0
3	NI	B	503	1/1	1.00	0.02	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.