



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:35 AM UTC

PDB ID : 7O5W / pdb_00007o5w
Title : Crystal structure of holo-F210W mutant of Hydroxy ketone aldolase (SwHKA) from *Sphingomonas wittichii* RW1
Authors : Laustsen, J.; Justo, I.; Marsden, S.R.; Hanefeld, U.; Bento, I.
Deposited on : 2021-04-09
Resolution : 1.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.20-1.20)
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	252	 92% 7% .
1	BBB	252	 92% 8%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8263 atoms, of which 3836 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HpcH/HpaI aldolase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	250	Total	C	H	N	O	S	48	4	0
			3781	1190	1894	334	354	9			
1	BBB	252	Total	C	H	N	O	S	45	3	0
			3817	1201	1912	341	354	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	0	HIS	-	expression tag	UNP A5VH82
AAA	1	HIS	-	expression tag	UNP A5VH82
AAA	210	TRP	PHE	engineered mutation	UNP A5VH82
BBB	0	HIS	-	expression tag	UNP A5VH82
BBB	1	HIS	-	expression tag	UNP A5VH82
BBB	210	TRP	PHE	engineered mutation	UNP A5VH82

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	AAA	1	Total C H O 9 2 5 2	1	0
2	AAA	1	Total C H O 9 2 5 2	1	0
2	BBB	1	Total C H O 9 2 5 2	1	0
2	BBB	1	Total C H O 9 2 5 2	1	0
2	BBB	1	Total C H O 17 4 10 3	1	0

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Mg 1 1	0	0
3	BBB	1	Total Mg 1 1	0	0

- Molecule 4 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	2	Total Br 2 2	0	0
4	BBB	1	Total Br 1 1	0	0

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AAA	1	Total K 1 1	0	0
5	BBB	1	Total K 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	311	Total O 311 311	0	0
6	BBB	294	Total O 294 294	0	2

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HpcH/HpaI aldolase

Chain AAA:  92% 7%



- Molecule 1: HpcH/HpaI aldolase

Chain BBB:  92% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	71.00Å 71.00Å 222.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.89 – 1.20 53.89 – 1.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (53.89-1.20) 99.9 (53.89-1.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 1.20Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.118 , 0.146 0.118 , 0.146	Depositor DCC
R_{free} test set	6604 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	9.9	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.043 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	8263	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, MG, BR, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.07	0/1942	1.03	2/2641 (0.1%)
1	BBB	1.16	4/1952 (0.2%)	1.10	3/2653 (0.1%)
All	All	1.11	4/3894 (0.1%)	1.07	5/5294 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	242	ARG	NE-CZ	7.17	1.41	1.33
1	BBB	195	HIS	CE1-NE2	6.37	1.39	1.32
1	BBB	184	THR	C-O	5.77	1.30	1.23
1	BBB	128	SER	C-O	-5.01	1.17	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	250	ALA	CA-C-O	-8.62	106.15	120.80
1	BBB	242	ARG	NE-CZ-NH2	6.55	125.10	119.20
1	AAA	233	ASP	CA-CB-CG	5.58	118.18	112.60
1	BBB	233	ASP	CA-CB-CG	5.34	117.94	112.60
1	BBB	178	HIS	CA-CB-CG	-5.29	108.51	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1887	1894	1895	9	1
1	BBB	1905	1912	1907	11	0
2	AAA	8	10	9	1	1
2	BBB	15	20	20	2	0
3	AAA	1	0	0	0	0
3	BBB	1	0	0	0	0
4	AAA	2	0	0	1	0
4	BBB	1	0	0	1	0
5	AAA	1	0	0	0	0
5	BBB	1	0	0	0	0
6	AAA	311	0	0	7	1
6	BBB	294	0	0	8	4
All	All	4427	3836	3831	24	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:101:GLN:NE2	6:AAA:401:HOH:O	1.94	0.99
1:BBB:185:GLU:HG2	6:BBB:429:HOH:O	1.86	0.75
6:AAA:541:HOH:O	4:BBB:305:BR:BR	2.65	0.69
2:AAA:302:PEG:O2	6:AAA:402:HOH:O	2.11	0.68
2:BBB:303:PEG:H21	6:BBB:495:HOH:O	1.95	0.66
1:AAA:101:GLN:HG2	6:AAA:658:HOH:O	2.05	0.56
1:BBB:197[A]:ARG:NH2	6:BBB:402:HOH:O	2.40	0.55
4:AAA:305:BR:BR	6:BBB:522:HOH:O	2.73	0.54
1:BBB:178:HIS:CG	1:BBB:186:GLU:HG2	2.43	0.53
1:BBB:3:LYS:NZ	6:BBB:403:HOH:O	2.42	0.53
1:BBB:1:HIS:HE1	6:BBB:616:HOH:O	1.91	0.52
1:AAA:1:HIS:N	6:AAA:406:HOH:O	2.42	0.51
1:BBB:185:GLU:HA	6:BBB:429:HOH:O	2.13	0.48
1:BBB:101:GLN:NE2	6:BBB:401:HOH:O	2.31	0.47
1:AAA:12:ARG:HB2	1:AAA:13:PRO:HD2	1.97	0.46
1:AAA:97:ARG:NH1	1:AAA:158:SER:OG	2.49	0.45
1:AAA:238[C]:SER:OG	1:AAA:242:ARG:NH1	2.50	0.44
1:BBB:97:ARG:HB2	1:BBB:155[A]:GLU:HB3	1.99	0.44
1:AAA:136:GLU:HG2	6:AAA:620:HOH:O	2.17	0.44
1:BBB:208:GLY:HA2	1:BBB:227:PHE:O	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:208:GLY:HA2	1:AAA:227:PHE:O	2.18	0.43
1:BBB:232:PRO:HG2	2:BBB:303:PEG:C1	2.49	0.42
1:AAA:5:ARG:HG3	6:AAA:594:HOH:O	2.18	0.42
1:BBB:143[B]:MET:HA	1:BBB:166:PHE:O	2.21	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:BBB:657:HOH:O	6:BBB:682:HOH:O[3_555]	1.92	0.28
6:BBB:482:HOH:O	6:BBB:649:HOH:O[3_555]	1.93	0.27
6:BBB:592:HOH:O	6:BBB:685:HOH:O[2_655]	2.08	0.12
1:AAA:223:GLU:OE2	2:AAA:301:PEG:H32[2_655]	1.55	0.05
6:AAA:490:HOH:O	6:BBB:401:HOH:O[5_555]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	254/252 (101%)	247 (97%)	6 (2%)	1 (0%)	30	10
1	BBB	253/252 (100%)	245 (97%)	7 (3%)	1 (0%)	30	10
All	All	507/504 (101%)	492 (97%)	13 (3%)	2 (0%)	30	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	118	GLY
1	BBB	118	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	193/189 (102%)	192 (100%)	1 (0%)	81	59
1	BBB	192/189 (102%)	192 (100%)	0	100	100
All	All	385/378 (102%)	384 (100%)	1 (0%)	86	67

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	114	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	BBB	302	-	3,3,6	0.42	0	2,2,5	0.50	0
2	PEG	AAA	302	-	3,3,6	1.05	0	2,2,5	0.72	0
2	PEG	BBB	303	-	6,6,6	1.17	1 (16%)	5,5,5	1.43	0
2	PEG	BBB	301	-	3,3,6	0.25	0	2,2,5	0.37	0
2	PEG	AAA	301	-	3,3,6	0.54	0	2,2,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	BBB	302	-	-	0/1/1/4	-
2	PEG	AAA	302	-	-	0/1/1/4	-
2	PEG	BBB	303	-	-	2/4/4/4	-
2	PEG	BBB	301	-	-	1/1/1/4	-
2	PEG	AAA	301	-	-	0/1/1/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	303	PEG	O4-C4	2.04	1.52	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	BBB	303	PEG	O1-C1-C2-O2
2	BBB	303	PEG	O2-C3-C4-O4
2	BBB	301	PEG	O2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	302	PEG	1	0
2	BBB	303	PEG	2	0
2	AAA	301	PEG	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	250/252 (99%)	-0.60	1 (0%) 88 91	4, 10, 20, 26	4 (1%)
1	BBB	252/252 (100%)	-0.38	0 100 100	5, 10, 20, 32	3 (1%)
All	All	502/504 (99%)	-0.49	1 (0%) 91 92	4, 10, 20, 32	7 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	250	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	BBB	301	4/7	0.82	0.13	0,28,30,31	9
2	PEG	BBB	303	7/7	0.88	0.16	6,17,27,29	17
2	PEG	AAA	301	4/7	0.91	0.12	0,17,18,19	9
2	PEG	BBB	302	4/7	0.94	0.09	0,21,29,30	9

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PEG	AAA	302	4/7	0.94	0.11	0,16,24,26	9
3	MG	AAA	303	1/1	1.00	0.04	9,9,9,9	0
3	MG	BBB	304	1/1	1.00	0.03	10,10,10,10	0
4	BR	AAA	304	1/1	1.00	0.01	15,15,15,15	1
4	BR	AAA	305	1/1	1.00	0.01	16,16,16,16	1
4	BR	BBB	305	1/1	1.00	0.07	20,20,20,20	1
5	K	AAA	306	1/1	1.00	0.01	9,9,9,9	0
5	K	BBB	306	1/1	1.00	0.02	9,9,9,9	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MG AAA 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.