



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:10 PM UTC

PDB ID : 5O6H / pdb_00005o6h
Title : Human NMT1 in complex with myristoyl-CoA and inhibitor IMP-917
Authors : Brannigan, J.A.; Wilkinson, A.J.
Deposited on : 2017-06-06
Resolution : 1.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	2% 81% 16% ..
1	B	391	4% 82% 14% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	2002	-	X	-	-
3	PO4	A	2003	-	X	-	-
3	PO4	B	2002	-	X	-	-
6	GOL	B	2010	-	X	-	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	28	0
			3298	2145	553	583	17			
1	B	382	Total	C	N	O	S	0	23	0
			3261	2122	545	577	17			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	106	GLY	-	expression tag	UNP P30419
A	107	PRO	-	expression tag	UNP P30419
A	108	HIS	-	expression tag	UNP P30419
B	106	GLY	-	expression tag	UNP P30419
B	107	PRO	-	expression tag	UNP P30419
B	108	HIS	-	expression tag	UNP P30419

- Molecule 2 is TETRADECANOYL-COA (CCD ID: MYA) (formula: $C_{35}H_{62}N_7O_{17}P_3S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	
			63	35	7	17	3	1	0
2	B	1	Total	C	N	O	P	S	
			63	35	7	17	3	1	0

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P		
			5	4	1	0	0
3	A	1	Total	O	P		
			5	4	1	0	0
3	A	1	Total	O	P		
			5	4	1	0	0
3	B	1	Total	O	P		
			5	4	1	0	0
3	B	1	Total	O	P		
			5	4	1	0	0
3	B	1	Total	O	P		
			5	4	1	0	0
3	B	1	Total	O	P		
			5	4	1	0	0

- Molecule 4 is 1-[5-[4-fluoranyl-2-[2-(1,3,5-trimethylpyrazol-4-yl)ethoxy]phenyl]-2 {H}-indazol-3-yl]- {N}, {N}-dimethyl-methanamine (CCD ID: 9M2) (formula: C₂₄H₂₈FN₅O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			31	24	1	5	1		
4	B	1	Total	C	F	N	O	0	0
			31	24	1	5	1		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	497	Total 497	O 497	0	0
8	B	521	Total 521	O 521	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1



• Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.14Å 178.85Å 58.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.00 – 1.29 55.00 – 1.29	Depositor EDS
% Data completeness (in resolution range)	98.7 (55.00-1.29) 98.7 (55.00-1.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.29Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.164 , 0.193 0.164 , 0.193	Depositor DCC
R_{free} test set	10337 reflections (4.02%)	wwPDB-VP
Wilson B-factor (Å ²)	11.1	Xtriage
Anisotropy	0.904	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7829	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MYA, 9M2, GOL, MG, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.67	34/3469 (1.0%)	1.36	13/4703 (0.3%)
1	B	1.67	26/3417 (0.8%)	1.41	13/4638 (0.3%)
All	All	1.67	60/6886 (0.9%)	1.38	26/9341 (0.3%)

All (60) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	189	ARG	CD-NE	-9.60	1.32	1.46
1	B	139	GLU	C-N	7.74	1.43	1.33
1	B	270	ARG	NE-CZ	-7.32	1.25	1.33
1	A	346	ILE	CA-C	7.18	1.58	1.52
1	B	189	ARG	CZ-NH2	-7.12	1.24	1.33
1	A	292	GLY	N-CA	7.10	1.52	1.45
1	A	426	HIS	N-CA	6.92	1.54	1.46
1	A	297	TRP	C-O	6.21	1.31	1.23
1	A	180	TYR	C-O	6.19	1.30	1.23
1	B	484	PRO	C-O	6.08	1.30	1.23
1	B	414	LYS	CA-C	6.07	1.60	1.52
1	B	168	VAL	N-CA	6.05	1.53	1.46
1	B	141	ASP	N-CA	6.00	1.53	1.46
1	B	286	VAL	C-O	5.93	1.31	1.24
1	A	139	GLU	N-CA	5.83	1.51	1.45
1	A	327	TYR	N-CA	5.79	1.53	1.46
1	A	148	GLU	CA-CB	-5.75	1.45	1.53
1	A	151	THR	CA-CB	5.73	1.61	1.53
1	B	460	THR	C-O	5.73	1.31	1.24
1	B	173	TYR	CA-CB	5.70	1.62	1.53
1	A	287	LEU	N-CA	5.68	1.50	1.45
1	B	166	ARG	N-CA	-5.68	1.39	1.46
1	B	185	ASP	N-CA	5.66	1.53	1.46
1	B	139	GLU	C-O	-5.65	1.19	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	135	HIS	N-CA	5.61	1.53	1.46
1	A	198	LEU	C-O	5.55	1.30	1.24
1	B	428	GLN	CA-C	-5.53	1.44	1.52
1	A	468	GLY	C-O	5.52	1.30	1.23
1	B	457	GLU	N-CA	5.50	1.52	1.46
1	A	380	ASN	N-CA	5.45	1.54	1.46
1	A	406	THR	N-CA	5.43	1.52	1.46
1	B	236	TYR	N-CA	5.36	1.53	1.46
1	B	494	VAL	C-O	5.34	1.30	1.24
1	B	385	PHE	N-CA	5.33	1.52	1.46
1	A	373	HIS	C-O	5.32	1.30	1.24
1	A	488	ALA	N-CA	5.31	1.52	1.46
1	A	188	PHE	N-CA	5.29	1.52	1.46
1	A	388	GLU	CD-OE1	5.27	1.35	1.25
1	A	139	GLU	C-N	5.26	1.39	1.33
1	A	318	MET	CA-C	-5.26	1.46	1.52
1	A	299	ARG	N-CA	5.24	1.52	1.46
1	B	491	VAL	N-CA	5.23	1.52	1.46
1	A	224	GLY	N-CA	5.22	1.50	1.45
1	A	300[A]	SER	C-O	5.21	1.30	1.23
1	A	300[B]	SER	C-O	5.21	1.30	1.23
1	B	282	THR	C-O	5.20	1.30	1.23
1	A	457	GLU	C-O	5.18	1.30	1.24
1	B	391	ASN	CG-OD1	-5.17	1.13	1.23
1	A	491	VAL	N-CA	5.16	1.52	1.46
1	A	387	VAL	N-CA	5.13	1.52	1.46
1	B	165	ASP	C-O	5.09	1.30	1.24
1	A	217	VAL	CA-C	5.06	1.59	1.52
1	A	297	TRP	CA-C	-5.06	1.46	1.52
1	B	483	CYS	C-O	-5.05	1.19	1.24
1	A	457	GLU	N-CA	5.04	1.51	1.46
1	A	495	LEU	N-CA	5.04	1.52	1.46
1	B	484	PRO	C-N	-5.04	1.26	1.33
1	A	150	TYR	C-O	-5.02	1.17	1.23
1	A	290	PRO	C-O	5.01	1.29	1.23
1	A	156	PHE	N-CA	5.01	1.52	1.45

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	189	ARG	NE-CZ-NH2	-21.92	99.47	119.20
1	B	189	ARG	NE-CZ-NH1	19.73	141.23	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	189	ARG	CD-NE-CZ	13.56	143.39	124.40
1	A	403	LEU	O-C-N	8.93	126.63	121.27
1	A	307	ILE	CA-C-O	-8.12	112.24	120.85
1	A	189	ARG	NE-CZ-NH1	-7.31	114.19	121.50
1	B	130	GLU	N-CA-C	-6.94	99.76	110.10
1	A	125	VAL	N-CA-C	-6.78	103.63	109.19
1	A	483	CYS	N-CA-C	-6.62	102.18	108.07
1	A	299	ARG	NE-CZ-NH2	-5.93	113.86	119.20
1	B	140	PRO	CA-C-O	-5.82	114.57	121.67
1	B	240	LYS	CG-CD-CE	-5.77	98.03	111.30
1	B	428	GLN	CB-CG-CD	5.76	122.40	112.60
1	A	217	VAL	O-C-N	5.73	127.43	121.87
1	B	152	LEU	CA-C-O	-5.69	114.51	120.88
1	A	388	GLU	CG-CD-OE2	5.51	131.07	118.40
1	B	316	ARG	N-CA-C	-5.44	102.00	110.10
1	A	440	VAL	CA-C-O	5.41	126.91	121.17
1	B	356	TYR	N-CA-C	5.39	117.15	111.28
1	B	240	LYS	N-CA-CB	-5.29	101.68	111.52
1	A	274	GLU	N-CA-C	-5.29	106.68	113.02
1	B	152	LEU	O-C-N	5.28	127.44	121.53
1	A	457	GLU	N-CA-C	-5.23	106.76	112.72
1	B	147	GLN	OE1-CD-NE2	5.22	127.82	122.60
1	A	152	LEU	CA-C-N	-5.15	114.41	119.92
1	A	152	LEU	C-N-CA	-5.15	114.41	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3298	0	3362	30	0
1	B	3261	0	3313	29	0
2	A	63	0	58	0	0
2	B	63	0	58	0	0
3	A	15	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	25	0	0	0	0
4	A	31	0	0	0	0
4	B	31	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	6	0	8	0	0
6	B	12	0	16	0	0
7	A	4	0	6	0	0
8	A	497	0	0	6	0
8	B	521	0	0	5	0
All	All	7829	0	6821	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:MET:CE	1:A:322:ARG:HG3	1.71	1.20
1:A:318:MET:HE3	1:A:322:ARG:CG	1.86	1.05
1:A:144[B]:ASN:HD21	1:B:351:GLN:NE2	1.60	0.99
1:A:318:MET:HE3	1:A:322:ARG:HG3	0.94	0.89
1:B:280[B]:VAL:HG22	1:B:478:LEU:HD23	1.55	0.87
1:A:144[B]:ASN:ND2	1:B:351:GLN:NE2	2.23	0.85
1:B:280[B]:VAL:HG22	1:B:478:LEU:CD2	2.06	0.84
1:B:393[B]:GLU:HG3	8:B:2420:HOH:O	1.83	0.77
1:B:187[A]:MET:HE2	8:B:2211:HOH:O	1.87	0.75
1:A:437[B]:ASP:OD1	8:A:2106:HOH:O	2.09	0.69
1:A:351:GLN:HE21	1:A:355:ARG:NH2	1.90	0.69
1:A:144[A]:ASN:HD22	1:A:144[A]:ASN:H	1.41	0.67
1:B:187[A]:MET:HE1	1:B:310:LYS:O	1.95	0.66
1:A:333:PRO:HB3	1:A:437[B]:ASP:CG	2.21	0.66
1:B:130:GLU:O	1:B:289[B]:LYS:HE2	1.98	0.63
1:B:412:THR:HB	8:B:2485:HOH:O	2.00	0.60
1:A:135:HIS:HE1	8:A:2472:HOH:O	1.85	0.60
1:B:187[A]:MET:HE3	1:B:188:PHE:CE2	2.37	0.59
1:B:482:LYS:HB3	8:B:2207:HOH:O	2.01	0.59
1:A:144[B]:ASN:HD21	1:B:351:GLN:CD	2.12	0.56
1:A:187:MET:HE1	1:A:310:LYS:HB2	1.88	0.56
1:B:319:THR:HB	1:B:321[B]:GLN:HE21	1.72	0.54
1:B:132:VAL:CG1	1:B:482:LYS:HG2	2.41	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:458:ASN:HA	1:B:461:PHE:CE2	2.47	0.50
1:B:318:MET:HG3	1:B:322:ARG:HG2	1.93	0.50
1:A:417[A]:LYS:HG2	1:A:447:PHE:CD1	2.49	0.48
1:B:315:SER:HB2	1:B:318:MET:HE3	1.96	0.48
1:B:434:LEU:O	1:B:437[A]:ASP:HB2	2.14	0.48
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.49	0.48
1:B:230:PRO:HB2	1:B:241[A]:LYS:HE3	1.96	0.47
1:B:132:VAL:HG11	1:B:482:LYS:HG2	1.96	0.47
1:B:126:PRO:HG3	1:B:482:LYS:HG3	1.97	0.47
1:B:417[A]:LYS:HA	1:B:417[A]:LYS:HD2	1.57	0.46
1:A:240:LYS:HE3	8:A:2480:HOH:O	2.16	0.45
1:A:490:LYS:NZ	8:A:2110:HOH:O	2.36	0.45
1:B:144:ASN:C	1:B:145[A]:ILE:HD12	2.42	0.45
1:A:234[B]:HIS:ND1	1:A:239[B]:GLU:CG	2.80	0.45
1:A:320:MET:O	1:A:324[B]:MET:HG3	2.16	0.45
1:A:334:LYS:N	1:A:437[B]:ASP:OD1	2.49	0.45
1:A:140:PRO:C	8:A:2109:HOH:O	2.61	0.44
1:A:132:VAL:CG1	1:A:482[A]:LYS:HG2	2.48	0.43
1:A:135:HIS:CE1	8:A:2472:HOH:O	2.67	0.43
1:B:437[B]:ASP:OD1	8:B:2102:HOH:O	2.21	0.43
1:B:134:THR:O	1:B:483:CYS:HA	2.20	0.42
1:B:254[A]:LEU:HD23	1:B:254[A]:LEU:HA	1.79	0.42
1:A:338[B]:LEU:HD11	1:A:437[B]:ASP:OD2	2.20	0.42
1:A:324[B]:MET:HE2	1:A:324[B]:MET:HB3	1.77	0.41
1:B:333:PRO:HB3	1:B:437[B]:ASP:CG	2.45	0.41
1:B:188:PHE:HE1	1:B:407[A]:ILE:HD12	1.86	0.41
1:A:320:MET:O	1:A:324[B]:MET:CG	2.70	0.40
1:B:187[A]:MET:HE3	1:B:188:PHE:HE2	1.80	0.40
1:A:132:VAL:HG13	1:A:482[A]:LYS:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/391 (104%)	399 (98%)	9 (2%)	0	100	100
1	B	403/391 (103%)	394 (98%)	9 (2%)	0	100	100
All	All	811/782 (104%)	793 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/355 (106%)	368 (98%)	8 (2%)	47	11
1	B	371/355 (104%)	368 (99%)	3 (1%)	73	45
All	All	747/710 (105%)	736 (98%)	11 (2%)	61	21

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144[A]	ASN
1	A	144[B]	ASN
1	A	304[A]	ARG
1	A	304[B]	ARG
1	A	310	LYS
1	A	325	LYS
1	A	334	LYS
1	A	455	LEU
1	B	316	ARG
1	B	412	THR
1	B	414	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	135	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	147	GLN
1	A	232	ASN
1	A	351	GLN
1	A	380	ASN
1	A	413	HIS
1	A	458	ASN
1	B	118	GLN
1	B	147	GLN
1	B	186	ASN
1	B	351	GLN
1	B	458	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	DMS	A	2008	-	3,3,3	0.62	0	3,3,3	1.09	0
3	PO4	B	2002	-	4,4,4	1.60	2 (50%)	6,6,6	1.74	2 (33%)
6	GOL	A	2007	-	5,5,5	1.08	0	5,5,5	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	B	2003	-	4,4,4	2.85	2 (50%)	6,6,6	1.84	1 (16%)
3	PO4	B	2005	-	4,4,4	0.88	0	6,6,6	1.05	0
3	PO4	A	2003	-	4,4,4	2.68	3 (75%)	6,6,6	2.15	1 (16%)
4	9M2	A	2005	-	32,34,34	1.95	9 (28%)	38,49,49	2.85	14 (36%)
4	9M2	B	2007	-	32,34,34	2.05	11 (34%)	38,49,49	2.39	11 (28%)
6	GOL	B	2010	-	5,5,5	1.06	0	5,5,5	2.04	2 (40%)
6	GOL	B	2009	-	5,5,5	1.47	1 (20%)	5,5,5	1.11	1 (20%)
2	MYA	B	2001	5	63,65,65	1.19	6 (9%)	85,91,91	1.33	8 (9%)
2	MYA	A	2001	5	63,65,65	1.21	7 (11%)	85,91,91	1.27	9 (10%)
3	PO4	A	2002	-	4,4,4	2.06	2 (50%)	6,6,6	1.88	2 (33%)
3	PO4	B	2006	-	4,4,4	0.62	0	6,6,6	1.09	0
3	PO4	A	2004	-	4,4,4	0.52	0	6,6,6	1.72	2 (33%)
3	PO4	B	2004	-	4,4,4	0.82	0	6,6,6	1.94	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	2007	-	-	0/4/4/4	-
4	9M2	A	2005	-	-	0/14/14/14	0/4/4/4
4	9M2	B	2007	-	-	0/14/14/14	0/4/4/4
6	GOL	B	2010	-	-	4/4/4/4	-
6	GOL	B	2009	-	-	0/4/4/4	-
2	MYA	B	2001	5	-	1/64/80/80	0/3/3/3
2	MYA	A	2001	5	-	0/64/80/80	0/3/3/3

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	2007	9M2	C4-N2	6.62	1.41	1.33
4	A	2005	9M2	C3-N1	-5.19	1.30	1.35
3	B	2003	PO4	P-O1	4.72	1.61	1.50
2	B	2001	MYA	P1A-O3A	4.14	1.64	1.59
3	A	2003	PO4	P-O3	-4.01	1.43	1.54
4	A	2005	9M2	C18-C22	3.99	1.49	1.40
2	A	2001	MYA	P1A-O3A	3.56	1.63	1.59
2	A	2001	MYA	O10-C10	3.50	1.48	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2005	9M2	C4-N2	3.33	1.37	1.33
4	A	2005	9M2	C10-C15	3.28	1.47	1.40
2	B	2001	MYA	O5-C5	3.22	1.29	1.23
4	B	2007	9M2	C18-C22	3.20	1.48	1.40
3	A	2002	PO4	P-O1	3.08	1.57	1.50
4	B	2007	9M2	C5-C4	2.99	1.44	1.40
4	A	2005	9M2	C19-C18	2.98	1.46	1.40
4	B	2007	9M2	C6-C5	2.93	1.43	1.38
2	B	2001	MYA	O10-C10	2.83	1.47	1.42
2	A	2001	MYA	C6-C5	-2.81	1.45	1.51
4	B	2007	9M2	C1-N	2.75	1.54	1.46
3	B	2003	PO4	P-O3	-2.57	1.47	1.54
4	A	2005	9M2	C14-C13	2.52	1.41	1.37
4	B	2007	9M2	C22-N4	2.49	1.37	1.32
2	B	2001	MYA	C3M-C2M	2.41	1.53	1.50
3	B	2002	PO4	P-O3	-2.40	1.47	1.54
4	B	2007	9M2	C11-C10	2.38	1.43	1.40
4	B	2007	9M2	C19-C18	2.38	1.45	1.40
3	A	2003	PO4	P-O2	-2.35	1.47	1.54
4	A	2005	9M2	C1-N	2.33	1.53	1.46
6	B	2009	GOL	O1-C1	2.29	1.52	1.42
3	A	2003	PO4	P-O4	2.28	1.61	1.54
4	B	2007	9M2	C8-C7	2.26	1.45	1.38
2	B	2001	MYA	P2A-O5A	-2.25	1.44	1.55
4	A	2005	9M2	C11-C10	2.23	1.43	1.40
4	B	2007	9M2	C14-C13	2.23	1.41	1.37
2	A	2001	MYA	C5A-N7A	-2.22	1.35	1.39
2	A	2001	MYA	C12-C11	2.20	1.56	1.52
2	A	2001	MYA	C6A-N6A	2.18	1.39	1.34
4	A	2005	9M2	C10-C7	-2.16	1.45	1.49
2	A	2001	MYA	P2A-O5A	-2.16	1.45	1.55
2	B	2001	MYA	C2A-N3A	2.14	1.37	1.33
3	A	2002	PO4	P-O4	-2.07	1.48	1.54
3	B	2002	PO4	P-O1	-2.02	1.46	1.50
4	B	2007	9M2	C6-C7	2.01	1.43	1.39

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2005	9M2	C18-C19-N3	-8.59	100.07	106.78
4	B	2007	9M2	C22-N4-N3	-7.12	97.59	105.55
4	A	2005	9M2	C22-N4-N3	-6.91	97.83	105.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2005	9M2	C19-N3-N4	6.04	123.11	112.75
4	A	2005	9M2	C21-N3-N4	-5.78	110.64	119.29
4	B	2007	9M2	C21-N3-N4	-5.45	111.13	119.29
3	A	2003	PO4	O3-P-O2	4.72	122.61	107.91
4	B	2007	9M2	C18-C22-N4	4.46	116.47	111.37
4	B	2007	9M2	C6-C7-C8	4.42	124.31	118.23
4	A	2005	9M2	C6-C5-C4	-4.26	115.70	120.80
4	B	2007	9M2	C19-N3-N4	3.98	119.58	112.75
2	B	2001	MYA	C3M-C2M-S1	3.89	118.04	113.40
3	B	2003	PO4	O3-P-O2	3.85	119.90	107.91
2	B	2001	MYA	C4A-N9A-C8A	3.85	109.78	105.74
4	B	2007	9M2	C19-C18-C22	-3.75	100.17	105.06
2	B	2001	MYA	C13-C11-C10	3.72	115.11	108.77
4	B	2007	9M2	C17-C18-C19	-3.68	120.66	126.74
4	A	2005	9M2	C18-C22-N4	3.60	115.48	111.37
2	A	2001	MYA	C13-C11-C10	3.56	114.84	108.77
2	A	2001	MYA	C3M-C2M-S1	3.48	117.55	113.40
3	A	2002	PO4	O4-P-O3	3.40	118.49	107.91
6	B	2010	GOL	C3-C2-C1	-3.39	99.38	111.80
4	B	2007	9M2	C8-C7-C10	-3.30	115.07	120.61
4	A	2005	9M2	C8-C7-C10	-3.24	115.17	120.61
3	B	2004	PO4	O4-P-O1	-3.15	99.82	110.95
4	A	2005	9M2	C-N-C2	3.11	115.47	110.38
4	A	2005	9M2	C20-C19-C18	3.07	133.62	128.20
2	B	2001	MYA	N9A-C8A-N7A	-3.07	109.58	113.94
2	B	2001	MYA	O2M-C2M-C3M	-3.05	120.45	123.98
2	A	2001	MYA	O2M-C2M-S1	-3.01	118.85	122.68
4	A	2005	9M2	C4-N2-N1	2.94	109.37	106.29
4	A	2005	9M2	C6-C7-C10	2.88	125.68	120.91
2	A	2001	MYA	O5A-P2A-O3A	2.76	114.74	107.27
3	A	2004	PO4	O4-P-O3	2.74	116.44	107.91
4	B	2007	9M2	C5-C6-C7	-2.74	117.70	121.43
3	A	2004	PO4	O4-P-O1	-2.61	101.74	110.95
3	A	2002	PO4	O2-P-O1	-2.57	101.88	110.95
2	A	2001	MYA	C4M-C3M-C2M	-2.55	106.64	112.27
3	B	2002	PO4	O4-P-O1	-2.53	102.02	110.95
2	B	2001	MYA	P3X-O3X-C3X	-2.50	116.75	123.43
2	A	2001	MYA	P3X-O3X-C3X	-2.49	116.78	123.43
3	B	2004	PO4	O4-P-O2	2.48	115.62	107.91
4	B	2007	9M2	C-N-C2	2.48	114.43	110.38
4	A	2005	9M2	C5-C6-C7	2.43	124.74	121.43
6	B	2010	GOL	O3-C3-C2	2.42	121.26	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	MYA	N3A-C2A-N1A	-2.40	124.94	128.58
2	B	2001	MYA	C14-C11-C10	-2.39	104.69	108.77
2	B	2001	MYA	N3A-C2A-N1A	-2.36	125.01	128.58
2	A	2001	MYA	O2X-C2X-C3X	2.33	117.70	111.19
4	A	2005	9M2	C17-C18-C19	-2.28	122.98	126.74
3	B	2002	PO4	O4-P-O3	2.27	114.99	107.91
4	B	2007	9M2	C12-C13-C14	2.06	125.95	123.23
6	B	2009	GOL	O2-C2-C3	-2.05	100.68	109.18
2	A	2001	MYA	C5A-C4A-N3A	-2.05	123.90	126.72
4	A	2005	9M2	C21-N3-C19	-2.04	126.22	128.09

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	2010	GOL	O1-C1-C2-C3
6	B	2010	GOL	C1-C2-C3-O3
6	B	2010	GOL	O1-C1-C2-O2
6	B	2010	GOL	O2-C2-C3-O3
2	B	2001	MYA	C6-C7-N8-C9

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/391 (97%)	-0.06	7 (1%) 67 72	7, 13, 28, 72	28 (7%)
1	B	382/391 (97%)	-0.09	15 (3%) 43 49	5, 12, 29, 66	27 (7%)
All	All	764/782 (97%)	-0.08	22 (2%) 53 60	5, 13, 29, 72	55 (7%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	316	ARG	4.6
1	B	412	THR	4.4
1	A	131[A]	VAL	4.1
1	B	414	LYS	4.0
1	B	133	ASN	3.4
1	B	131	VAL	3.3
1	B	129	GLY	2.8
1	A	115	ARG	2.7
1	B	315	SER	2.6
1	B	317	ASN	2.6
1	B	116	SER	2.5
1	A	133	ASN	2.5
1	A	313	HIS	2.3
1	A	134	THR	2.3
1	B	115	ARG	2.2
1	B	128	LEU	2.2
1	B	132	VAL	2.2
1	B	117	TYR	2.1
1	A	132	VAL	2.1
1	B	314	LEU	2.1
1	A	129	GLY	2.0
1	B	411	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	A	2004	5/5	0.75	0.15	46,54,62,64	0
3	PO4	B	2006	5/5	0.76	0.13	44,56,58,62	0
3	PO4	B	2005	5/5	0.78	0.13	42,44,61,61	0
3	PO4	B	2004	5/5	0.79	0.16	48,48,64,64	0
6	GOL	B	2010	6/6	0.82	0.14	23,33,38,49	0
7	DMS	A	2008	4/4	0.82	0.19	36,37,39,40	0
6	GOL	B	2009	6/6	0.92	0.09	16,17,18,20	0
6	GOL	A	2007	6/6	0.96	0.07	17,18,20,20	0
4	9M2	A	2005	31/31	0.96	0.07	12,14,20,24	0
4	9M2	B	2007	31/31	0.97	0.06	11,13,17,18	0
2	MYA	A	2001	63/63	0.98	0.04	8,12,16,23	0
3	PO4	B	2002	5/5	0.98	0.06	16,18,23,23	0
3	PO4	B	2003	5/5	0.98	0.07	16,16,18,20	0
3	PO4	A	2003	5/5	0.98	0.06	16,16,19,19	0
3	PO4	A	2002	5/5	0.99	0.05	17,18,20,21	0
2	MYA	B	2001	63/63	0.99	0.04	8,11,14,17	0
5	MG	A	2006	1/1	0.99	0.11	21,21,21,21	0
5	MG	B	2008	1/1	0.99	0.10	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 9M2 A 2005:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9M2 B 2007:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around MYA A 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around MYA B 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.