



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:03 PM UTC

PDB ID : 3O8O / pdb_00003o8o
Title : Structure of phosphofructokinase from *Saccharomyces cerevisiae*
Authors : Banaszak, K.; Mechin, I.; Kopperschlager, G.; Rypniewski, W.
Deposited on : 2010-08-03
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	787	
1	C	787	
1	E	787	
1	G	787	
2	B	766	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	766	2% 51% 40% 8% .
2	F	766	% 51% 39% 8% ..
2	H	766	2% 51% 40% 8% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 46645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 6-phosphofructokinase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	750	Total	C	N	O	S	273	0	0
			5759	3620	1019	1098	22			
1	C	750	Total	C	N	O	S	196	0	0
			5759	3620	1019	1098	22			
1	E	752	Total	C	N	O	S	260	0	0
			5777	3631	1023	1101	22			
1	G	746	Total	C	N	O	S	289	0	0
			5733	3604	1015	1092	22			

- Molecule 2 is a protein called 6-phosphofructokinase subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	763	Total	C	N	O	S	364	0	0
			5834	3652	1034	1118	30			
2	D	763	Total	C	N	O	S	266	0	0
			5834	3652	1034	1118	30			
2	F	762	Total	C	N	O	S	279	0	0
			5827	3647	1033	1117	30			
2	H	763	Total	C	N	O	S	173	0	0
			5834	3652	1034	1118	30			

- Molecule 3 is 6-O-phosphono-beta-D-fructofuranose (CCD ID: F6P) (formula: C₆H₁₃O₉P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			16	6	9	1		
3	B	1	Total	C	O	P	0	0
			16	6	9	1		
3	C	1	Total	C	O	P	0	0
			16	6	9	1		
3	D	1	Total	C	O	P	0	0
			16	6	9	1		
3	E	1	Total	C	O	P	0	0
			16	6	9	1		
3	F	1	Total	C	O	P	0	0
			16	6	9	1		
3	G	1	Total	C	O	P	0	0
			16	6	9	1		
3	H	1	Total	C	O	P	0	0
			16	6	9	1		

- Molecule 4 is 2,6-di-O-phosphono-beta-D-fructofuranose (CCD ID: FDP) (formula: $C_6H_{14}O_{12}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			20	6	12	2		
4	B	1	Total	C	O	P	0	0
			20	6	12	2		
4	C	1	Total	C	O	P	0	0
			20	6	12	2		
4	D	1	Total	C	O	P	0	0
			20	6	12	2		
4	E	1	Total	C	O	P	0	0
			20	6	12	2		
4	F	1	Total	C	O	P	0	0
			20	6	12	2		
4	G	1	Total	C	O	P	0	0
			20	6	12	2		
4	H	1	Total	C	O	P	0	0
			20	6	12	2		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 6-phosphofructokinase subunit alpha





- Molecule 1: 6-phosphofructokinase subunit alpha






• Molecule 1: 6-phosphofructokinase subunit alpha






• Molecule 2: 6-phosphofructokinase subunit beta





• Molecule 2: 6-phosphofructokinase subunit beta





• Molecule 2: 6-phosphofructokinase subunit beta





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	180.05Å 186.21Å 236.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.90 35.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (35.00-2.90) 98.3 (35.00-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.90Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.259 , 0.309 0.257 , 0.305	Depositor DCC
R_{free} test set	3446 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	70.0	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	46645	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F6P, FDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	2/5860 (0.0%)	1.11	41/7925 (0.5%)
1	C	0.72	1/5860 (0.0%)	1.12	38/7925 (0.5%)
1	E	0.73	0/5878	1.14	47/7948 (0.6%)
1	G	0.75	0/5834	1.13	37/7889 (0.5%)
2	B	0.65	2/5940 (0.0%)	1.11	32/8038 (0.4%)
2	D	0.70	3/5940 (0.1%)	1.13	40/8038 (0.5%)
2	F	0.64	1/5932 (0.0%)	1.11	36/8027 (0.4%)
2	H	0.75	1/5940 (0.0%)	1.16	45/8038 (0.6%)
All	All	0.70	10/47184 (0.0%)	1.13	316/63828 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
2	F	0	1
2	H	0	1
All	All	0	4

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	836	ILE	CA-CB	6.96	1.62	1.54
1	A	719	ILE	CA-CB	6.09	1.60	1.53
2	D	733	ALA	CA-CB	5.88	1.62	1.53
2	H	736	ALA	CA-CB	-5.83	1.44	1.53
2	B	783	VAL	CA-CB	-5.58	1.50	1.55

The worst 5 of 316 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	592	ASP	N-CA-C	-13.36	94.65	110.41
2	H	811	GLY	N-CA-C	11.95	128.62	114.16
2	B	811	GLY	N-CA-C	11.75	128.38	114.16
2	H	352	SER	N-CA-C	11.53	125.28	111.33
2	D	811	GLY	N-CA-C	11.10	128.42	114.66

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	771	TYR	Sidechain
2	D	771	TYR	Sidechain
2	F	771	TYR	Sidechain
2	H	771	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5759	0	5758	355	0
1	C	5759	0	5758	349	0
1	E	5777	0	5779	343	0
1	G	5733	0	5733	330	0
2	B	5834	0	5799	385	0
2	D	5834	0	5799	373	0
2	F	5827	0	5791	349	1
2	H	5834	0	5799	392	1
3	A	16	0	11	5	0
3	B	16	0	11	4	0
3	C	16	0	11	1	0
3	D	16	0	11	3	0
3	E	16	0	11	2	0
3	F	16	0	11	4	0
3	G	16	0	11	2	0
3	H	16	0	11	5	0
4	A	20	0	10	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	10	3	0
4	C	20	0	10	1	0
4	D	20	0	10	2	0
4	E	20	0	10	0	0
4	F	20	0	10	3	0
4	G	20	0	10	0	0
4	H	20	0	10	2	0
All	All	46645	0	46384	2783	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 2783 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:427:HIS:HB3	1:C:460:ASP:HB2	1.33	1.10
1:A:665:ARG:HH21	4:A:1:FDP:H62	1.12	1.07
2:F:418:THR:HG1	2:F:422:TRP:CD1	1.72	1.07
1:A:427:HIS:HB3	1:A:460:ASP:HB2	1.35	1.06
2:H:874:LYS:HD3	2:H:917:VAL:HG11	1.34	1.06

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:333:GLU:OE2	2:H:578:GLU:OE1[4_446]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	746/787 (95%)	630 (84%)	87 (12%)	29 (4%)	2	10
1	C	746/787 (95%)	634 (85%)	85 (11%)	27 (4%)	2	11
1	E	748/787 (95%)	643 (86%)	74 (10%)	31 (4%)	2	9
1	G	742/787 (94%)	632 (85%)	79 (11%)	31 (4%)	2	9
2	B	761/766 (99%)	675 (89%)	61 (8%)	25 (3%)	3	13
2	D	761/766 (99%)	672 (88%)	59 (8%)	30 (4%)	2	10
2	F	760/766 (99%)	674 (89%)	58 (8%)	28 (4%)	2	11
2	H	761/766 (99%)	680 (89%)	53 (7%)	28 (4%)	2	11
All	All	6025/6212 (97%)	5240 (87%)	556 (9%)	229 (4%)	2	10

5 of 229 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	252	GLY
1	A	321	GLU
1	A	331	ALA
1	A	585	SER
1	A	587	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	614/645 (95%)	559 (91%)	55 (9%)	9	28
1	C	614/645 (95%)	556 (91%)	58 (9%)	8	26
1	E	616/645 (96%)	558 (91%)	58 (9%)	8	26
1	G	612/645 (95%)	551 (90%)	61 (10%)	7	24
2	B	612/615 (100%)	547 (89%)	65 (11%)	6	22
2	D	612/615 (100%)	547 (89%)	65 (11%)	6	22
2	F	611/615 (99%)	549 (90%)	62 (10%)	7	24
2	H	612/615 (100%)	552 (90%)	60 (10%)	7	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	4903/5040 (97%)	4419 (90%)	484 (10%)	7 24

5 of 484 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	890	ARG
2	H	472	LEU
1	E	809	ASN
2	H	366	ARG
2	H	869	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 175 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	331	GLN
1	G	817	ASN
2	F	522	ASN
1	G	431	GLN
1	G	930	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	FDP	F	6	-	19,20,20	0.94	0	29,32,32	0.91	2 (6%)
4	FDP	A	1	-	19,20,20	0.93	0	29,32,32	0.77	0
4	FDP	B	2	-	19,20,20	0.92	0	29,32,32	0.85	1 (3%)
4	FDP	G	7	-	19,20,20	0.93	0	29,32,32	0.81	1 (3%)
4	FDP	E	5	-	19,20,20	0.95	0	29,32,32	0.89	1 (3%)
3	F6P	E	988	-	15,16,16	0.84	0	16,25,25	0.81	0
4	FDP	C	3	-	19,20,20	0.96	0	29,32,32	0.80	0
4	FDP	H	8	-	19,20,20	0.94	0	29,32,32	0.93	1 (3%)
3	F6P	H	986	-	15,16,16	0.83	0	16,25,25	0.79	1 (6%)
3	F6P	D	982	-	15,16,16	0.90	0	16,25,25	0.78	0
3	F6P	A	988	-	15,16,16	0.92	0	16,25,25	3.82	5 (31%)
4	FDP	D	4	-	19,20,20	0.93	0	29,32,32	0.85	1 (3%)
3	F6P	F	984	-	15,16,16	0.89	0	16,25,25	0.89	0
3	F6P	B	980	-	15,16,16	1.04	1 (6%)	16,25,25	1.03	1 (6%)
3	F6P	C	988	-	15,16,16	0.90	0	16,25,25	0.88	0
3	F6P	G	988	-	15,16,16	1.09	1 (6%)	16,25,25	3.74	5 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FDP	F	6	-	-	6/12/34/34	0/1/1/1
4	FDP	A	1	-	-	2/12/34/34	0/1/1/1
4	FDP	B	2	-	-	0/12/34/34	0/1/1/1
4	FDP	G	7	-	-	1/12/34/34	0/1/1/1
4	FDP	E	5	-	-	3/12/34/34	0/1/1/1
3	F6P	E	988	-	-	1/9/28/28	0/1/1/1
4	FDP	C	3	-	-	0/12/34/34	0/1/1/1
4	FDP	H	8	-	-	1/12/34/34	0/1/1/1
3	F6P	H	986	-	-	1/9/28/28	0/1/1/1
3	F6P	D	982	-	-	1/9/28/28	0/1/1/1
3	F6P	A	988	-	-	3/9/28/28	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FDP	D	4	-	-	2/12/34/34	0/1/1/1
3	F6P	F	984	-	-	1/9/28/28	0/1/1/1
3	F6P	B	980	-	-	1/9/28/28	0/1/1/1
3	F6P	C	988	-	-	1/9/28/28	0/1/1/1
3	F6P	G	988	-	-	3/9/28/28	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	988	F6P	O5-C2	-2.35	1.39	1.43
3	B	980	F6P	P-O3P	2.25	1.63	1.54

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	988	F6P	O3P-P-O6	-8.96	83.32	106.67
3	G	988	F6P	O3P-P-O6	-8.88	83.52	106.67
3	A	988	F6P	O6-P-O1P	6.86	124.98	106.44
3	A	988	F6P	O3P-P-O1P	-6.86	84.12	110.83
3	G	988	F6P	O6-P-O1P	6.79	124.80	106.44

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	988	F6P	C6-O6-P-O3P
3	G	988	F6P	C6-O6-P-O1P
3	G	988	F6P	C6-O6-P-O3P
4	A	1	FDP	C6-O6-P2-O4P
4	A	1	FDP	C6-O6-P2-O5P

There are no ring outliers.

14 monomers are involved in 43 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	6	FDP	3	0
4	A	1	FDP	6	0
4	B	2	FDP	3	0
3	E	988	F6P	2	0
4	C	3	FDP	1	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	8	FDP	2	0
3	H	986	F6P	5	0
3	D	982	F6P	3	0
3	A	988	F6P	5	0
4	D	4	FDP	2	0
3	F	984	F6P	4	0
3	B	980	F6P	4	0
3	C	988	F6P	1	0
3	G	988	F6P	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









Ligand FDP G 7




Ligand FDP E 5




Ligand F6P E 988




Ligand FDP H 8




Ligand F6P H 986




Ligand F6P D 982




Ligand F6P A 988




Ligand FDP D 4




Ligand F6P F 984




Ligand F6P B 980



Ligand F6P C 988





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	750/787 (95%)	0.27	29 (3%)	43	35	17, 56, 81, 81	67 (8%)
1	C	750/787 (95%)	0.34	35 (4%)	36	28	16, 53, 81, 81	49 (6%)
1	E	752/787 (95%)	0.15	18 (2%)	59	50	12, 50, 81, 81	58 (7%)
1	G	744/787 (94%)	0.19	39 (5%)	33	26	19, 49, 81, 81	60 (8%)
2	B	763/766 (99%)	0.43	37 (4%)	35	28	17, 65, 81, 81	88 (11%)
2	D	761/766 (99%)	0.15	14 (1%)	67	59	20, 57, 81, 81	54 (7%)
2	F	762/766 (99%)	0.12	11 (1%)	73	65	24, 58, 81, 81	61 (8%)
2	H	763/766 (99%)	-0.14	12 (1%)	70	62	17, 45, 77, 81	37 (4%)
All	All	6045/6212 (97%)	0.19	195 (3%)	50	41	12, 54, 81, 81	474 (7%)

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	805	LEU	7.1
1	C	788	TYR	7.0
1	G	326	VAL	6.9
2	B	896	PHE	6.6
1	G	327	ASP	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FDP	H	8	20/20	0.71	0.12	81,82,82,82	0
4	FDP	C	3	20/20	0.73	0.14	79,82,82,82	0
4	FDP	F	6	20/20	0.75	0.11	81,82,82,82	0
3	F6P	A	988	16/16	0.79	0.12	79,82,82,82	0
3	F6P	B	980	16/16	0.85	0.11	77,80,82,82	0
3	F6P	C	988	16/16	0.86	0.15	75,78,80,80	0
4	FDP	A	1	20/20	0.88	0.10	77,82,82,82	0
3	F6P	E	988	16/16	0.89	0.11	62,66,66,68	0
4	FDP	G	7	20/20	0.90	0.09	80,82,82,82	0
3	F6P	G	988	16/16	0.91	0.12	61,62,64,68	0
3	F6P	F	984	16/16	0.91	0.11	56,57,63,64	0
3	F6P	D	982	16/16	0.92	0.11	65,67,70,71	0
4	FDP	B	2	20/20	0.93	0.12	65,68,70,71	0
4	FDP	E	5	20/20	0.94	0.08	66,70,74,74	0
3	F6P	H	986	16/16	0.95	0.08	41,47,47,49	0
4	FDP	D	4	20/20	0.97	0.06	52,54,56,57	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FDP H 8:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FDP F 6:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P A 988:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around F6P B 980:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P C 988:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FDP A 1:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P E 988:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FDP G 7:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P G 988:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P F 984:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around F6P D 982:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FDP B 2:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FDP E 5:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around F6P H 986:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.