



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:42 PM UTC

PDB ID : 4O8G / pdb_00004o8g
Title : Structure of Infrared Fluorescent Protein 1.4
Authors : Bhattacharya, S.; Forest, K.T.
Deposited on : 2013-12-27
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	13% 90% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bacteriophytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	10	0
			2432	1553	429	439	11			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	SER	conflict	UNP Q9RZA4
A	54	VAL	MET	engineered mutation	UNP Q9RZA4
A	119	ALA	GLY	engineered mutation	UNP Q9RZA4
A	135	ILE	THR	conflict	UNP Q9RZA4
A	186	MET	VAL	engineered mutation	UNP Q9RZA4
A	195	MET	LEU	engineered mutation	UNP Q9RZA4
A	196	GLN	HIS	engineered mutation	UNP Q9RZA4
A	207	HIS	ASP	engineered mutation	UNP Q9RZA4
A	208	THR	ILE	engineered mutation	UNP Q9RZA4
A	288	VAL	ALA	engineered mutation	UNP Q9RZA4
A	307	GLU	TYR	engineered mutation	UNP Q9RZA4
A	311	LYS	LEU	engineered mutation	UNP Q9RZA4
A	314	GLY	LEU	engineered mutation	UNP Q9RZA4
A	318	ARG	VAL	engineered mutation	UNP Q9RZA4
A	322	LEU	ALA	conflict	UNP Q9RZA4
A	323	GLU	ASP	conflict	UNP Q9RZA4
A	324	HIS	-	expression tag	UNP Q9RZA4
A	325	HIS	-	expression tag	UNP Q9RZA4
A	326	HIS	-	expression tag	UNP Q9RZA4
A	327	HIS	-	expression tag	UNP Q9RZA4
A	328	HIS	-	expression tag	UNP Q9RZA4
A	329	HIS	-	expression tag	UNP Q9RZA4

- Molecule 2 is 3-[2-[(Z)-[3-(2-carboxyethyl)-5-[(Z)-(4-ethenyl-3-methyl-5-oxidanylidene-pyrro-2-ylidene)methyl]-4-methyl-pyrrol-1-ium-2-ylidene]methyl]-5-[(Z)-[(3E)-3-ethylidene-4-methyl-5-oxidanylidene-pyrrolidin-2-ylidene]methyl]-4-methyl-1H-pyrrol-3-yl]propanoic acid (CCD ID: LBV) (formula: C₃₃H₃₇N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	1
			43	33	4	6		

- Molecule 3 is 3-[2-[(Z)-[5-[(Z)-(4-ethenyl-3-methyl-5-oxidanylidene-pyrrol-2-ylidene)methyl]-3-(3-hydroxy-3-oxopropyl)-4-methyl-pyrrol-1-ium-2-ylidene]methyl]-5-[(Z)-[(3E,4R)-3-ethylidene-4-methyl-5-oxidanylidene-pyrrolidin-2-ylidene]methyl]-4-methyl-1H-pyrrol-3-yl]propanoic acid (CCD ID: LBW) (formula: $C_{33}H_{37}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			43	33	4	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total 227	O 227	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacteriophytochrome



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	96.16Å 53.23Å 66.80Å 90.00° 90.60° 90.00°	Depositor
Resolution (Å)	24.04 – 1.65 24.04 – 1.65	Depositor EDS
% Data completeness (in resolution range)	93.4 (24.04-1.65) 93.4 (24.04-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.03 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.185 , 0.211 0.195 , 0.220	Depositor DCC
R_{free} test set	1876 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.007 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.009 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.014 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.013 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.016 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2745	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LBW, LBV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2523	0.82	0/3447

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2462	14	0
2	A	43	0	33	1	0
3	A	43	0	33	2	0
4	A	227	0	0	2	0
All	All	2745	0	2528	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:MET:HB2	1:A:288[A]:VAL:CG1	2.29	0.62
1:A:174:MET:HE3	1:A:288[A]:VAL:CG1	2.36	0.56
1:A:259:MET:HG2	4:A:698:HOH:O	2.06	0.55
1:A:174:MET:HB2	1:A:288[A]:VAL:HG13	1.90	0.54
1:A:288[A]:VAL:HG23	4:A:510:HOH:O	2.08	0.54
1:A:203:PHE:CZ	2:A:401[A]:LBV:HBD2	2.48	0.49
1:A:290:HIS:CE1	3:A:402[B]:LBW:O_D	2.66	0.49
1:A:143:ALA:O	1:A:147[A]:LEU:HD23	2.13	0.49
1:A:174:MET:HE3	1:A:288[A]:VAL:HG13	1.95	0.47
1:A:98:GLN:HG2	1:A:116[B]:HIS:NE2	2.29	0.47
1:A:174:MET:HE3	1:A:288[A]:VAL:HG11	1.98	0.44
1:A:193:GLU:HA	1:A:194:GLY:HA2	1.79	0.43
1:A:147[A]:LEU:HD22	1:A:159:VAL:HG12	2.02	0.41
1:A:221:LEU:C	1:A:221:LEU:HD23	2.46	0.41
3:A:402[B]:LBW:O_D	3:A:402[B]:LBW:HBD1	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/329 (96%)	312 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	261/265 (98%)	260 (100%)	1 (0%)	84 77

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LBV	A	401[A]	1	46,46,46	3.79	14 (30%)	58,67,67	2.10	13 (22%)
3	LBW	A	402[B]	1	46,46,46	3.60	15 (32%)	58,67,67	2.04	17 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LBV	A	401[A]	1	-	11/26/74/74	0/4/4/4
3	LBW	A	402[B]	1	-	7/26/74/74	0/4/4/4

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[A]	LBV	C1A-N_A	14.18	1.56	1.37
3	A	402[B]	LBW	C1A-N_A	13.03	1.54	1.37
2	A	401[A]	LBV	C4D-N_D	8.85	1.58	1.38
2	A	401[A]	LBV	C2A-C1A	8.67	1.62	1.52
3	A	402[B]	LBW	C2A-C1A	8.39	1.61	1.52
3	A	402[B]	LBW	C4D-N_D	8.25	1.57	1.38
3	A	402[B]	LBW	C2A-C3A	-8.24	1.40	1.51
2	A	401[A]	LBV	C2A-C3A	-8.10	1.41	1.51
2	A	401[A]	LBV	C3D-C2D	-6.66	1.23	1.37
2	A	401[A]	LBV	CHB-C1B	6.49	1.55	1.40
3	A	402[B]	LBW	C3D-C2D	-6.18	1.24	1.37
3	A	402[B]	LBW	CHB-C1B	5.68	1.53	1.40
2	A	401[A]	LBV	CHC-C4B	5.31	1.52	1.40
3	A	402[B]	LBW	C4A-C3A	-5.20	1.35	1.45
3	A	402[B]	LBW	CHC-C4B	4.99	1.51	1.40
2	A	401[A]	LBV	C4A-C3A	-4.78	1.36	1.45
3	A	402[B]	LBW	CHD-C4C	4.29	1.50	1.40
2	A	401[A]	LBV	C1D-C2D	-4.29	1.37	1.45
2	A	401[A]	LBV	CHD-C4C	4.19	1.50	1.40
3	A	402[B]	LBW	C1D-C2D	-3.85	1.38	1.45
2	A	401[A]	LBV	C1D-N_D	3.64	1.44	1.37
3	A	402[B]	LBW	C1D-N_D	2.89	1.42	1.37
2	A	401[A]	LBV	CAA-C3A	2.73	1.40	1.33
3	A	402[B]	LBW	CMA-C2A	-2.68	1.45	1.53
3	A	402[B]	LBW	C4C-C3C	2.63	1.51	1.45
3	A	402[B]	LBW	O_D-C4D	-2.59	1.18	1.23
2	A	401[A]	LBV	C4C-C3C	2.45	1.51	1.45
2	A	401[A]	LBV	CMA-C2A	-2.42	1.46	1.53
3	A	402[B]	LBW	C4A-N_A	-2.32	1.33	1.37

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[A]	LBV	C3D-C4D-N_D	-6.51	97.95	106.13
3	A	402[B]	LBW	O_A-C1A-C2A	5.82	132.91	126.74
3	A	402[B]	LBW	C3D-C4D-N_D	-5.42	99.33	106.13
2	A	401[A]	LBV	O_A-C1A-N_A	4.79	130.57	124.93
2	A	401[A]	LBV	C4D-C3D-C2D	4.75	114.03	107.92
2	A	401[A]	LBV	C3D-C2D-C1D	4.75	113.21	107.92
3	A	402[B]	LBW	C1D-N_D-C4D	-4.75	104.83	110.66
2	A	401[A]	LBV	C1D-N_D-C4D	-4.46	105.18	110.66
3	A	402[B]	LBW	C3D-C2D-C1D	4.42	112.84	107.92
2	A	401[A]	LBV	O_A-C1A-C2A	4.33	131.33	126.74
3	A	402[B]	LBW	C4D-C3D-C2D	3.84	112.86	107.92
2	A	401[A]	LBV	CHC-C4B-C3B	-3.18	120.38	127.22
3	A	402[B]	LBW	CAA-C3A-C4A	-3.09	122.62	126.36
3	A	402[B]	LBW	CHC-C4B-C3B	-3.05	120.67	127.22
3	A	402[B]	LBW	O_A-C1A-N_A	3.02	128.49	124.93
2	A	401[A]	LBV	CAA-C3A-C4A	-2.93	122.81	126.36
2	A	401[A]	LBV	CHC-C4B-N_B	2.65	131.27	125.29
3	A	402[B]	LBW	CHC-C4B-N_B	2.56	131.07	125.29
3	A	402[B]	LBW	CHC-C1C-N_C	2.51	129.86	124.60
2	A	401[A]	LBV	C2C-C1C-N_C	-2.46	106.56	110.04
3	A	402[B]	LBW	C2C-C1C-N_C	-2.43	106.61	110.04
2	A	401[A]	LBV	CHB-C1B-C2B	-2.34	121.79	127.53
3	A	402[B]	LBW	CHD-C1D-N_D	-2.28	121.19	126.06
3	A	402[B]	LBW	CBB-CAB-C3B	-2.27	106.25	112.53
2	A	401[A]	LBV	CHB-C1B-N_B	2.26	130.40	125.29
3	A	402[B]	LBW	CHB-C1B-C2B	-2.21	122.11	127.53
3	A	402[B]	LBW	CHB-C4A-N_A	-2.17	121.44	126.06
2	A	401[A]	LBV	CHD-C4C-C3C	2.15	130.40	124.87
3	A	402[B]	LBW	C2D-C1D-N_D	2.15	110.10	106.97
3	A	402[B]	LBW	CHB-C1B-N_B	2.07	129.96	125.29

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401[A]	LBV	C4A-C3A-CAA-CBA
2	A	401[A]	LBV	C3C-C4C-CHD-C1D
2	A	401[A]	LBV	N_C-C4C-CHD-C1D
3	A	402[B]	LBW	C2A-C3A-CAA-CBA
3	A	402[B]	LBW	C3C-C4C-CHD-C1D
3	A	402[B]	LBW	N_C-C4C-CHD-C1D
3	A	402[B]	LBW	N_A-C4A-CHB-C1B
2	A	401[A]	LBV	N_A-C4A-CHB-C1B

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401[A]	LBV	C2C-CAC-CBC-CGC
2	A	401[A]	LBV	N_B-C1B-CHB-C4A
2	A	401[A]	LBV	CAB-CBB-CGB-O2B
2	A	401[A]	LBV	CAB-CBB-CGB-O1B
3	A	402[B]	LBW	CAB-CBB-CGB-O2B
3	A	402[B]	LBW	CAB-CBB-CGB-O1B
2	A	401[A]	LBV	C2A-C3A-CAA-CBA
2	A	401[A]	LBV	CAC-CBC-CGC-O1C
3	A	402[B]	LBW	CAC-CBC-CGC-O1C
2	A	401[A]	LBV	CAC-CBC-CGC-O2C


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[A]	LBV	1	0
3	A	402[B]	LBW	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LBV A 401 (A)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/329 (94%)	0.50	42 (13%) 7 7	6, 16, 41, 77	14 (4%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	ALA	7.5
1	A	194	GLY	6.7
1	A	105	TRP	6.1
1	A	96	ALA	5.1
1	A	106	PRO	5.1
1	A	91	PRO	4.5
1	A	195	MET	4.0
1	A	5	PRO	3.8
1	A	322	LEU	3.8
1	A	192	ARG	3.7
1	A	97	LEU	3.6
1	A	93	CYS	3.6
1	A	87	ALA	3.6
1	A	138	HIS	3.5
1	A	90[A]	PRO	3.5
1	A	193	GLU	3.4
1	A	196	GLN	3.4
1	A	6	LEU	3.2
1	A	139	ALA	3.2
1	A	77	LEU	2.9
1	A	141	ARG	2.9
1	A	321	ALA	2.9
1	A	109	GLY	2.9
1	A	140	LEU	2.9
1	A	92	GLY	2.8
1	A	110	HIS	2.8
1	A	198	PHE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	100	ARG	2.6
1	A	88	ALA	2.6
1	A	95	ASP	2.6
1	A	86	GLN	2.5
1	A	85	LEU	2.5
1	A	4	ASP	2.4
1	A	82	TRP	2.3
1	A	323	GLU	2.2
1	A	94	PRO	2.2
1	A	84	ALA	2.1
1	A	80	GLU	2.1
1	A	145	PHE	2.1
1	A	130	GLU	2.0
1	A	142	ASN	2.0
1	A	76	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LBV	A	401[A]	43/43	0.94	0.07	10,11,14,17	43
3	LBW	A	402[B]	43/43	0.95	0.07	9,10,13,17	43

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LBV A 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LBW A 402 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.