



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:40 PM UTC

PDB ID : 4O8Z / pdb_00004o8z
Title : Crystal structure of human SIRT3 in complex with compound (2-butylbenzofuran-3-yl)(4-(2-(diethylamino)ethoxy)-3,5-diiodophenyl)methanone
Authors : Liu, B.
Deposited on : 2013-12-31
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	 7% 81% 8% • 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	272	Total	C	N	O	S	0	1	0
			2142	1381	367	385	9			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	97	HIS	-	expression tag	UNP Q9NTG7
A	98	HIS	-	expression tag	UNP Q9NTG7
A	99	HIS	-	expression tag	UNP Q9NTG7
A	100	HIS	-	expression tag	UNP Q9NTG7
A	101	HIS	-	expression tag	UNP Q9NTG7
A	102	HIS	-	expression tag	UNP Q9NTG7
A	103	ASP	-	expression tag	UNP Q9NTG7
A	104	TYR	-	expression tag	UNP Q9NTG7
A	105	ASP	-	expression tag	UNP Q9NTG7
A	106	ILE	-	expression tag	UNP Q9NTG7
A	107	PRO	-	expression tag	UNP Q9NTG7
A	108	THR	-	expression tag	UNP Q9NTG7
A	109	THR	-	expression tag	UNP Q9NTG7
A	110	GLU	-	expression tag	UNP Q9NTG7
A	111	ASN	-	expression tag	UNP Q9NTG7
A	112	LEU	-	expression tag	UNP Q9NTG7
A	113	TYR	-	expression tag	UNP Q9NTG7
A	114	PHE	-	expression tag	UNP Q9NTG7
A	115	GLN	-	expression tag	UNP Q9NTG7

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

-
- The chemical structure of BBI (Bibenzylidene) is shown with various substituents labeled with three-letter codes. The structure consists of two benzene rings connected by a central carbon atom. The substituents are labeled as follows:
- Top ring: CAH, GAG, OAF, CAE, CAD, CAB, CAA, CAC, CAM, CAL, CAI, CAJ, CAK, CBE, GAP, CAQ, CAR, IAS, CAV, CAW, CBA, CBB, CAY, NAX, CAZ.
 - Bottom ring: IBD, CBC, CAT, OAU, CAV, CAW, CBA, CBB, CAY, NAX, CAZ.


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	217	Total O 217 217	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	116.32Å 116.32Å 45.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	100.73 – 2.00 100.73 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.6 (100.73-2.00) 98.6 (100.73-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.73 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.167 , 0.204 0.178 , 0.217	Depositor DCC
R_{free} test set	1216 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.034 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2391	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BBI, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	4/2196 (0.2%)	1.14	9/2991 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	154	ILE	CA-C	6.67	1.58	1.52
1	A	172	ASP	CA-C	6.35	1.61	1.52
1	A	204	TYR	C-O	-5.63	1.17	1.23
1	A	154	ILE	C-N	5.04	1.39	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	ILE	CB-CA-C	-10.64	99.25	110.53
1	A	154	ILE	CA-C-N	-7.64	112.47	120.03
1	A	154	ILE	C-N-CA	-7.64	112.47	120.03
1	A	172	ASP	N-CA-C	6.97	125.64	110.80
1	A	349	GLY	CA-C-N	6.74	127.00	119.32
1	A	349	GLY	C-N-CA	6.74	127.00	119.32
1	A	154	ILE	N-CA-C	6.62	115.86	108.05
1	A	175	TYR	N-CA-C	-5.85	100.56	108.11
1	A	138	GLN	CB-CA-C	-5.09	100.82	109.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2142	0	2153	14	0
2	A	1	0	0	0	0
3	A	31	0	29	2	0
4	A	217	0	0	2	1
All	All	2391	0	2182	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ARG:NH1	4:A:711:HOH:O	2.22	0.71
1:A:207:ASN:H	1:A:210:HIS:HD2	1.42	0.67
1:A:316:LEU:HD13	1:A:341:LEU:HD23	1.84	0.60
1:A:164:LEU:HD11	1:A:168:LEU:HD21	1.84	0.59
1:A:166:SER:HA	1:A:170:GLN:OE1	2.04	0.58
3:A:402:BB1:CAE	3:A:402:BB1:H14	2.37	0.53
1:A:171:TYR:O	1:A:172:ASP:HB2	2.08	0.52
1:A:165:TYR:CE1	1:A:169:GLN:HG3	2.47	0.50
1:A:167:ASN:O	1:A:171:TYR:HB2	2.11	0.49
1:A:296:GLU:HB3	1:A:297:PRO:HD2	1.94	0.49
1:A:164:LEU:CD1	1:A:168:LEU:HD21	2.45	0.46
1:A:210:HIS:HE1	4:A:501:HOH:O	2.00	0.45
1:A:207:ASN:H	1:A:210:HIS:CD2	2.29	0.44
1:A:311:MET:HE3	1:A:311:MET:HB2	1.87	0.43
3:A:402:BB1:H14	3:A:402:BB1:CAD	2.50	0.42
1:A:177:GLU:CD	1:A:177:GLU:H	2.29	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:508:HOH:O	4:A:680:HOH:O[1_556]	1.97	0.23

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/303 (89%)	263 (98%)	5 (2%)	1 (0%)	30	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	172	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/264 (90%)	232 (98%)	5 (2%)	47	52

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	ILE
1	A	168	LEU
1	A	200	TYR
1	A	324	VAL
1	A	381	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	GLN
1	A	203	ASN
1	A	210	HIS
1	A	228	GLN
1	A	248	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	BBI	A	402	-	33,33,33	2.03	9 (27%)	44,45,45	1.79	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BBI	A	402	-	-	6/22/22/22	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	BBI	CAP-CAN	-5.02	1.39	1.49
3	A	402	BBI	CAL-CAM	-5.01	1.34	1.46
3	A	402	BBI	CAM-CAN	-4.61	1.34	1.49
3	A	402	BBI	CAL-CAG	-4.23	1.34	1.39
3	A	402	BBI	CAH-CAG	-3.19	1.33	1.39
3	A	402	BBI	CAK-CAL	-3.09	1.35	1.39
3	A	402	BBI	CAD-CAE	2.92	1.57	1.49
3	A	402	BBI	CAR-IAS	-2.49	2.04	2.10
3	A	402	BBI	CBC-IBD	-2.12	2.05	2.10

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	BBI	OAF-CAE-CAD	5.49	123.55	115.22
3	A	402	BBI	CAD-CAE-CAM	-5.22	128.77	133.84
3	A	402	BBI	CAL-CAM-CAE	2.99	109.48	106.75
3	A	402	BBI	OAF-CAE-CAM	-2.91	107.67	111.19
3	A	402	BBI	CAH-CAG-CAL	-2.78	120.39	123.58
3	A	402	BBI	CAG-CAL-CAM	2.48	107.77	105.64
3	A	402	BBI	CAI-CAJ-CAK	-2.32	117.38	120.24
3	A	402	BBI	OAF-CAG-CAL	-2.26	108.62	110.84
3	A	402	BBI	CAT-CBC-IBD	2.14	122.84	119.32
3	A	402	BBI	OAF-CAG-CAH	2.14	130.98	126.49
3	A	402	BBI	CAV-CAW-NAX	-2.09	107.45	113.36

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402	BBI	CBC-CAT-OAU-CAV
3	A	402	BBI	CAR-CAT-OAU-CAV
3	A	402	BBI	OAU-CAV-CAW-NAX
3	A	402	BBI	CAL-CAM-CAN-CAP
3	A	402	BBI	CAW-CAV-OAU-CAT
3	A	402	BBI	CBB-CBA-NAX-CAY

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	BB1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/303 (89%)	0.09	22 (8%) 18 17	13, 27, 62, 95	1 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	TYR	4.8
1	A	155	PRO	3.9
1	A	169	GLN	3.8
1	A	164	LEU	3.5
1	A	163	GLY	3.4
1	A	392	GLY	3.4
1	A	394	LEU	3.4
1	A	170	GLN	3.3
1	A	391	THR	3.1
1	A	160	PRO	3.0
1	A	159	SER	3.0
1	A	172	ASP	2.8
1	A	167	ASN	2.6
1	A	162	SER	2.6
1	A	388	GLN	2.6
1	A	175	TYR	2.6
1	A	157	PHE	2.5
1	A	203	ASN	2.5
1	A	165	TYR	2.4
1	A	154	ILE	2.2
1	A	156	ASP	2.1
1	A	389	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BBI	A	402	31/31	0.61	0.29	78,94,121,162	19
2	ZN	A	401	1/1	1.00	0.01	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BBI A 402:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.