



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 08:22 PM UTC

PDB ID : 5O82 / pdb_00005o82
Title : Crystal Structure of R67A/E173A Mutant of alpha-L-arabinofuranosidase Ara51 from Clostridium thermocellum in complex with arabinofuranose
Authors : Lafite, P.; Daniellou, R.
Deposited on : 2017-06-12
Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	3% 83% 15% ..
1	B	501	5% 83% 15% ..
1	C	501	10% 78% 19% ..
1	D	501	4% 84% 14% ..
1	E	501	6% 81% 16% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	501	4% 80% 17% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Intracellular exo-alpha-(1->5)-L-arabinofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	0	1	0
			3954	2512	671	748	23			
1	B	496	Total	C	N	O	S	0	0	0
			3962	2519	673	748	22			
1	C	496	Total	C	N	O	S	0	0	0
			3965	2520	673	750	22			
1	D	496	Total	C	N	O	S	0	0	0
			3965	2520	673	750	22			
1	E	495	Total	C	N	O	S	0	0	0
			3956	2514	671	749	22			
1	F	495	Total	C	N	O	S	0	0	0
			3955	2514	671	748	22			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	ALA	ARG	engineered mutation	UNP A3DIH0
A	173	ALA	GLU	engineered mutation	UNP A3DIH0
B	67	ALA	ARG	engineered mutation	UNP A3DIH0
B	173	ALA	GLU	engineered mutation	UNP A3DIH0
C	67	ALA	ARG	engineered mutation	UNP A3DIH0
C	173	ALA	GLU	engineered mutation	UNP A3DIH0
D	67	ALA	ARG	engineered mutation	UNP A3DIH0
D	173	ALA	GLU	engineered mutation	UNP A3DIH0
E	67	ALA	ARG	engineered mutation	UNP A3DIH0
E	173	ALA	GLU	engineered mutation	UNP A3DIH0
F	67	ALA	ARG	engineered mutation	UNP A3DIH0
F	173	ALA	GLU	engineered mutation	UNP A3DIH0

- Molecule 2 is alpha-L-arabinofuranose (CCD ID: AHR) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			10	5	5		
2	C	1	Total	C	O	0	0
			10	5	5		
2	D	1	Total	C	O	0	0
			10	5	5		
2	E	1	Total	C	O	0	0
			10	5	5		
2	F	1	Total	C	O	0	0
			10	5	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	40	Total	O	0	0
			40	40		
3	B	50	Total	O	0	0
			50	50		
3	C	47	Total	O	0	0
			47	47		
3	D	42	Total	O	0	0
			42	42		
3	E	44	Total	O	0	0
			44	44		
3	F	36	Total	O	0	0
			36	36		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Intracellular exo- α -(1->5)-L-arabinofuranosidase




- Molecule 1: Intracellular exo- α -(1->5)-L-arabinofuranosidase





- Molecule 1: Intracellular exo- α -(1->5)-L-arabinofuranosidase









● Molecule 1: Intracellular exo-alpha-(1->5)-L-arabinofuranosidase




● Molecule 1: Intracellular exo-alpha-(1->5)-L-arabinofuranosidase



● Molecule 1: Intracellular exo-alpha-(1->5)-L-arabinofuranosidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	170.55Å 170.55Å 265.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.85 – 2.39 47.85 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.85-2.39) 99.8 (47.85-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.38 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.11.1	Depositor
R, R_{free}	0.187 , 0.237 0.191 , 0.239	Depositor DCC
R_{free} test set	7712 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	24076	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AHR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/4044	0.59	0/5480
1	B	0.39	0/4052	0.64	3/5488 (0.1%)
1	C	0.44	0/4055	0.64	3/5492 (0.1%)
1	D	0.42	1/4055 (0.0%)	0.60	2/5492 (0.0%)
1	E	0.40	0/4046	0.59	0/5481
1	F	0.50	0/4045	0.62	3/5480 (0.1%)
All	All	0.43	1/24297 (0.0%)	0.61	11/32913 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	223	GLN	C-O	-5.29	1.17	1.24

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	431	ASN	CA-C-N	6.30	133.04	121.70
1	C	431	ASN	C-N-CA	6.30	133.04	121.70
1	B	476	ASN	CA-C-N	6.24	132.93	121.70
1	B	476	ASN	C-N-CA	6.24	132.93	121.70
1	F	471	GLU	N-CA-C	-6.22	105.85	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3954	0	3837	53	0
1	B	3962	0	3859	53	0
1	C	3965	0	3861	84	0
1	D	3965	0	3861	53	0
1	E	3956	0	3848	56	0
1	F	3955	0	3845	70	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
2	E	10	0	0	0	0
2	F	10	0	0	0	0
3	A	40	0	0	0	0
3	B	50	0	0	0	0
3	C	47	0	0	0	0
3	D	42	0	0	0	0
3	E	44	0	0	0	0
3	F	36	0	0	0	0
All	All	24076	0	23111	365	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 365 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:444:GLY:HA3	1:B:445:MET:HG2	1.40	1.00
1:B:107:ILE:CD1	1:B:123:MET:HE1	1.96	0.96
1:C:478:ASP:HA	1:C:479:LYS:HB3	1.45	0.96
1:C:10:LYS:NZ	1:C:445:MET:SD	2.39	0.96
1:B:107:ILE:HD13	1:B:123:MET:HE1	1.47	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	490/501 (98%)	465 (95%)	19 (4%)	6 (1%)	10	16
1	B	490/501 (98%)	461 (94%)	21 (4%)	8 (2%)	7	11
1	C	490/501 (98%)	448 (91%)	29 (6%)	13 (3%)	4	4
1	D	490/501 (98%)	450 (92%)	33 (7%)	7 (1%)	9	13
1	E	489/501 (98%)	454 (93%)	25 (5%)	10 (2%)	6	7
1	F	489/501 (98%)	463 (95%)	21 (4%)	5 (1%)	12	20
All	All	2938/3006 (98%)	2741 (93%)	148 (5%)	49 (2%)	7	10

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	364	ARG
1	A	484	ASP
1	B	364	ARG
1	B	478	ASP
1	C	10	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	425/438 (97%)	425 (100%)	0	100	100
1	B	426/438 (97%)	426 (100%)	0	100	100
1	C	427/438 (98%)	427 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	427/438 (98%)	427 (100%)	0	100	100
1	E	426/438 (97%)	422 (99%)	4 (1%)	70	85
1	F	425/438 (97%)	420 (99%)	5 (1%)	63	81
All	All	2556/2628 (97%)	2547 (100%)	9 (0%)	84	92

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	484	ASP
1	F	487	LEU
1	E	478	ASP
1	F	44	LYS
1	F	450	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	468	ASN
1	F	223	GLN
1	F	431	ASN
1	F	365	ASN
1	F	179	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AHR	E	600	-	10,10,10	1.52	2 (20%)	13,14,14	1.52	3 (23%)
2	AHR	A	600	-	10,10,10	1.73	2 (20%)	13,14,14	1.45	2 (15%)
2	AHR	B	600	-	10,10,10	1.78	3 (30%)	13,14,14	1.33	2 (15%)
2	AHR	D	600	-	10,10,10	1.80	2 (20%)	13,14,14	1.27	2 (15%)
2	AHR	F	600	-	10,10,10	1.85	2 (20%)	13,14,14	1.37	1 (7%)
2	AHR	C	600	-	10,10,10	1.82	3 (30%)	13,14,14	1.36	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AHR	E	600	-	-	0/2/18/18	0/1/1/1
2	AHR	A	600	-	-	2/2/18/18	0/1/1/1
2	AHR	B	600	-	-	2/2/18/18	0/1/1/1
2	AHR	D	600	-	-	2/2/18/18	0/1/1/1
2	AHR	F	600	-	-	1/2/18/18	0/1/1/1
2	AHR	C	600	-	-	2/2/18/18	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	600	AHR	C2-C3	-3.56	1.43	1.53
2	A	600	AHR	C2-C3	-3.48	1.43	1.53
2	F	600	AHR	C2-C3	-3.44	1.44	1.53
2	F	600	AHR	C1-C2	-3.33	1.48	1.52
2	C	600	AHR	C2-C3	-3.20	1.44	1.53

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	AHR	O1-C1-O4	-3.56	106.59	111.12
2	F	600	AHR	C1-C2-C3	3.21	106.24	102.29
2	C	600	AHR	O4-C4-C5	2.92	115.39	109.22
2	E	600	AHR	C1-C2-C3	2.67	105.57	102.29
2	C	600	AHR	C1-C2-C3	2.63	105.52	102.29

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	AHR	O4-C4-C5-O5
2	B	600	AHR	O4-C4-C5-O5
2	C	600	AHR	O4-C4-C5-O5
2	C	600	AHR	C3-C4-C5-O5
2	D	600	AHR	O4-C4-C5-O5

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	495/501 (98%)	0.03	16 (3%)	50	46	17, 44, 75, 99	1 (0%)
1	B	496/501 (99%)	-0.01	23 (4%)	37	33	31, 43, 77, 100	0
1	C	496/501 (99%)	0.37	52 (10%)	11	8	32, 46, 89, 115	0
1	D	496/501 (99%)	0.16	19 (3%)	44	40	30, 48, 82, 100	0
1	E	495/501 (98%)	0.13	28 (5%)	29	25	30, 45, 84, 116	0
1	F	495/501 (98%)	0.18	22 (4%)	39	34	32, 49, 87, 113	0
All	All	2973/3006 (98%)	0.14	160 (5%)	31	28	17, 45, 83, 116	1 (0%)

The worst 5 of 160 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	502	GLY	6.6
1	D	367	GLY	5.6
1	B	478	ASP	5.0
1	C	12	TYR	4.7
1	E	367	GLY	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	AHR	E	600	10/10	0.77	0.21	36,43,48,50	10
2	AHR	F	600	10/10	0.79	0.20	41,47,51,54	10
2	AHR	C	600	10/10	0.80	0.20	34,44,50,53	10
2	AHR	A	600	10/10	0.83	0.20	35,42,47,49	10
2	AHR	D	600	10/10	0.84	0.17	38,48,56,59	10
2	AHR	B	600	10/10	0.85	0.17	38,40,48,49	10

6.5 Other polymers [i](#)

There are no such residues in this entry.