



Full wwPDB EM Validation Report ⓘ

Mar 10, 2026 – 04:13 AM UTC

PDB ID : 5O8O / pdb_00005o8o
EMDB ID : EMD-3761
Title : N. crassa Tom40 model based on cryo-EM structure of the TOM core complex at 6.8 Å
Authors : Bausewein, T.; Mills, D.J.; Nussberger, S.; Nitschke, B.; Kuehlbrandt, W.
Deposited on : 2017-06-13
Resolution : 6.80 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	503 (6.30 - 7.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	16% 69% 25% • 5%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5062 atoms, of which 2500 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitochondrial import receptor subunit tom40.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	332	5062	1617	2500	443	493	9	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitochondrial import receptor subunit tom40



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	92144	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	JEOL 3200FSC	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.189	Depositor
Minimum map value	-0.071	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.0568	Depositor
Map size (Å)	203.84, 203.84, 203.84	wwPDB
Map dimensions	182, 182, 182	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.12, 1.12, 1.12	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2613	1.56	82/3531 (2.3%)

There are no bond length outliers.

All (82) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	38	LEU	CA-C-N	10.18	135.17	120.82
1	A	38	LEU	C-N-CA	10.18	135.17	120.82
1	A	128	ARG	CA-C-O	9.61	131.22	119.31
1	A	254	ALA	CA-C-N	8.61	128.65	119.78
1	A	254	ALA	C-N-CA	8.61	128.65	119.78
1	A	128	ARG	O-C-N	-8.01	111.41	122.46
1	A	92	ASN	CA-C-N	7.86	127.53	119.82
1	A	92	ASN	C-N-CA	7.86	127.53	119.82
1	A	62	PHE	CA-C-N	-7.33	110.68	121.24
1	A	62	PHE	C-N-CA	-7.33	110.68	121.24
1	A	138	GLY	O-C-N	-7.30	115.71	123.45
1	A	42	GLY	CA-C-N	-7.18	113.51	122.84
1	A	42	GLY	C-N-CA	-7.18	113.51	122.84
1	A	277	PHE	CA-C-N	-7.03	110.91	120.82
1	A	277	PHE	C-N-CA	-7.03	110.91	120.82
1	A	312	ASP	CA-C-N	-6.97	114.02	123.14
1	A	312	ASP	C-N-CA	-6.97	114.02	123.14
1	A	247	VAL	CA-C-N	-6.88	110.58	121.87
1	A	247	VAL	C-N-CA	-6.88	110.58	121.87
1	A	191	LEU	CA-C-N	-6.75	113.15	123.14
1	A	191	LEU	C-N-CA	-6.75	113.15	123.14
1	A	203	GLY	CA-C-N	6.67	132.35	120.88
1	A	203	GLY	C-N-CA	6.67	132.35	120.88
1	A	139	GLY	O-C-N	6.64	128.63	122.19
1	A	184	THR	O-C-N	-6.59	117.20	121.88
1	A	112	ASP	O-C-N	6.37	130.53	123.33
1	A	37	GLY	O-C-N	6.29	129.97	122.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	347	ILE	CA-C-N	6.21	126.13	119.85
1	A	347	ILE	C-N-CA	6.21	126.13	119.85
1	A	64	GLY	O-C-N	-6.20	112.82	122.18
1	A	40	ASN	CA-C-N	6.14	125.76	119.56
1	A	40	ASN	C-N-CA	6.14	125.76	119.56
1	A	232	ASN	CA-C-N	-6.05	113.08	122.59
1	A	232	ASN	C-N-CA	-6.05	113.08	122.59
1	A	67	ALA	CA-C-N	-6.00	114.30	122.95
1	A	67	ALA	C-N-CA	-6.00	114.30	122.95
1	A	86	ALA	O-C-N	5.90	129.69	123.03
1	A	208	ILE	CA-C-N	-5.88	113.09	121.50
1	A	208	ILE	C-N-CA	-5.88	113.09	121.50
1	A	201	THR	CA-C-O	5.84	126.10	119.27
1	A	68	ASP	CA-C-N	-5.82	115.60	123.10
1	A	68	ASP	C-N-CA	-5.82	115.60	123.10
1	A	210	TYR	CA-C-N	-5.81	114.23	122.94
1	A	210	TYR	C-N-CA	-5.81	114.23	122.94
1	A	67	ALA	CA-C-O	-5.77	114.28	120.92
1	A	66	ARG	CA-C-O	5.75	127.84	121.11
1	A	145	GLN	O-C-N	-5.73	115.56	123.12
1	A	281	THR	CA-C-N	-5.63	112.69	122.37
1	A	281	THR	C-N-CA	-5.63	112.69	122.37
1	A	284	ALA	CA-C-N	-5.46	112.19	121.85
1	A	284	ALA	C-N-CA	-5.46	112.19	121.85
1	A	72	ALA	CA-C-N	5.45	131.64	122.76
1	A	72	ALA	C-N-CA	5.45	131.64	122.76
1	A	222	SER	CA-C-N	-5.44	112.22	121.85
1	A	222	SER	C-N-CA	-5.44	112.22	121.85
1	A	154	ASP	CA-C-O	5.41	126.06	119.78
1	A	326	VAL	CA-C-N	-5.38	114.02	122.73
1	A	326	VAL	C-N-CA	-5.38	114.02	122.73
1	A	269	THR	CA-C-N	-5.34	114.28	122.62
1	A	269	THR	C-N-CA	-5.34	114.28	122.62
1	A	77	PRO	O-C-N	5.27	129.01	122.22
1	A	244	GLN	CA-C-N	-5.27	112.64	121.75
1	A	244	GLN	C-N-CA	-5.27	112.64	121.75
1	A	219	TRP	CA-C-N	-5.25	115.81	122.37
1	A	219	TRP	C-N-CA	-5.25	115.81	122.37
1	A	43	THR	CA-C-O	5.17	125.92	120.54
1	A	187	LEU	CA-C-O	-5.13	115.35	121.56
1	A	327	SER	CA-C-N	-5.12	114.58	122.68
1	A	327	SER	C-N-CA	-5.12	114.58	122.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	THR	CA-C-N	-5.07	114.01	122.64
1	A	201	THR	C-N-CA	-5.07	114.01	122.64
1	A	207	ALA	CA-C-N	-5.07	116.35	123.10
1	A	207	ALA	C-N-CA	-5.07	116.35	123.10
1	A	160	LYS	CA-C-N	-5.07	113.65	122.37
1	A	160	LYS	C-N-CA	-5.07	113.65	122.37
1	A	183	VAL	N-CA-C	-5.04	107.52	113.42
1	A	235	PHE	CA-C-N	-5.04	113.71	122.37
1	A	235	PHE	C-N-CA	-5.04	113.71	122.37
1	A	128	ARG	CA-C-N	5.03	129.77	123.03
1	A	128	ARG	C-N-CA	5.03	129.77	123.03
1	A	73	PHE	CA-C-N	-5.03	113.73	120.87
1	A	73	PHE	C-N-CA	-5.03	113.73	120.87

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2562	2500	2499	35	0
All	All	2562	2500	2499	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:PHE:CD1	1:A:62:PHE:C	2.65	0.74
1:A:35:GLN:OE1	1:A:35:GLN:N	2.21	0.73
1:A:91:LEU:O	1:A:91:LEU:HG	1.98	0.61
1:A:62:PHE:C	1:A:62:PHE:HD1	2.08	0.60
1:A:292:LEU:C	1:A:292:LEU:HD12	2.31	0.56
1:A:99:LEU:C	1:A:99:LEU:HD12	2.30	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:LEU:C	1:A:171:LEU:HD12	2.33	0.53
1:A:45:GLU:OE1	1:A:45:GLU:N	2.36	0.52
1:A:240:THR:OG1	1:A:243:VAL:N	2.43	0.51
1:A:291:LYS:NZ	1:A:293:SER:OG	2.44	0.50
1:A:254:ALA:N	1:A:264:THR:OG1	2.44	0.50
1:A:161:ALA:HA	1:A:174:ILE:O	2.13	0.48
1:A:165:SER:OG	1:A:172:THR:OG1	2.30	0.48
1:A:242:ARG:NE	1:A:242:ARG:HA	2.30	0.46
1:A:214:TYR:HB2	1:A:221:ALA:HB3	1.98	0.46
1:A:106:PHE:H	1:A:124:ARG:HB2	1.81	0.44
1:A:110:ASN:O	1:A:119:THR:HA	2.17	0.43
1:A:62:PHE:HD1	1:A:62:PHE:O	2.02	0.43
1:A:157:ALA:HA	1:A:178:ASP:O	2.19	0.43
1:A:45:GLU:OE2	1:A:160:LYS:NZ	2.50	0.43
1:A:107:ALA:HA	1:A:122:ASN:O	2.19	0.42
1:A:243:VAL:HA	1:A:274:LYS:O	2.20	0.42
1:A:84:GLN:HB2	1:A:95:ALA:HB3	2.00	0.42
1:A:248:ASP:OD1	1:A:248:ASP:C	2.61	0.42
1:A:51:VAL:HG22	1:A:236:TRP:HB2	2.02	0.41
1:A:163:ASN:N	1:A:164:PRO:CD	2.82	0.41
1:A:296:LEU:O	1:A:308:THR:HA	2.20	0.41
1:A:108:GLN:O	1:A:121:PHE:HA	2.21	0.41
1:A:122:ASN:HA	1:A:133:THR:O	2.20	0.41
1:A:144:ALA:O	1:A:162:ILE:HA	2.21	0.41
1:A:297:GLU:HA	1:A:307:LEU:O	2.20	0.41
1:A:312:ASP:HB2	1:A:321:LYS:HB3	2.02	0.40
1:A:254:ALA:HA	1:A:255:PRO:HD3	1.93	0.40
1:A:78:LEU:HD12	1:A:78:LEU:N	2.37	0.40
1:A:124:ARG:NH1	1:A:124:ARG:HB3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/349 (95%)	321 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/280 (95%)	265 (100%)	1 (0%)	84	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	GLN
1	A	202	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3761. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 91



Y Index: 91




Z Index: 91


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 83



Y Index: 78




Z Index: 102


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 85 nm^3 ; this corresponds to an approximate mass of 77 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.147 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.147 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.80	-	-
Author-provided FSC curve	6.77	8.80	7.60
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-3761 and PDB model 5O8O. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0568 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0568).

9.4 Atom inclusion [i](#)



At the recommended contour level, 63% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0568) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5920	0.1580
A	0.5940	0.1580

