



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:33 PM UTC

PDB ID : 7O8Z / pdb_00007o8z
Title : Anomalous bromide substructure of NmHR under continuous illumination determined at 13.7 keV with serial crystallography
Authors : Mous, S.; Gotthard, G.; Ehrenberg, D.; Sen, S.; James, D.; Johnson, P.; Weinert, T.; Nass, K.; Furrer, A.; Kekilli, D.; Ma, P.; Bruenle, S.; Casadei, C.; Martiel, I.; Dworkowski, F.; Gashi, D.; Skopintsev, P.; Wranik, M.; Knopp, G.; Panepucci, E.; Panneels, V.; Cirelli, C.; Ozerov, D.; Schertler, G.; Wang, M.; Milne, C.; Standfuss, J.; Schapiro, I.; Heberle, J.; Nogly, P.
Deposited on : 2021-04-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.49

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BR	A	303[B]	-	-	-	X
3	BR	A	306[B]	-	-	X	-
3	BR	A	310[B]	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2312 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chloride pumping rhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	2099	1402	317	361	19	0	8	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	initiating methionine	UNP W8VZW3
A	-14	ALA	-	expression tag	UNP W8VZW3
A	-13	SER	-	expression tag	UNP W8VZW3
A	-12	MET	-	expression tag	UNP W8VZW3
A	-11	THR	-	expression tag	UNP W8VZW3
A	-10	GLY	-	expression tag	UNP W8VZW3
A	-9	GLY	-	expression tag	UNP W8VZW3
A	-8	GLN	-	expression tag	UNP W8VZW3
A	-7	GLN	-	expression tag	UNP W8VZW3
A	-6	MET	-	expression tag	UNP W8VZW3
A	-5	GLY	-	expression tag	UNP W8VZW3
A	-4	ARG	-	expression tag	UNP W8VZW3
A	-3	ASP	-	expression tag	UNP W8VZW3
A	-2	PRO	-	expression tag	UNP W8VZW3
A	-1	ASN	-	expression tag	UNP W8VZW3
A	0	SER	-	expression tag	UNP W8VZW3
A	273	LEU	-	expression tag	UNP W8VZW3
A	274	GLU	-	expression tag	UNP W8VZW3
A	275	HIS	-	expression tag	UNP W8VZW3
A	276	HIS	-	expression tag	UNP W8VZW3
A	277	HIS	-	expression tag	UNP W8VZW3
A	278	HIS	-	expression tag	UNP W8VZW3
A	279	HIS	-	expression tag	UNP W8VZW3
A	280	HIS	-	expression tag	UNP W8VZW3

- Molecule 2 is RETINAL (CCD ID: RET) (formula: C₂₀H₂₈O).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 20 20	0	0

- Molecule 3 is BROMIDE ION (CCD ID: BR) (formula: Br).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	9	Total Br 9 9	0	9

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (CCD ID: OLC) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			25	21	4		
4	A	1	Total	C	O	0	0
			13	9	4		

- Molecule 5 is OLEIC ACID (CCD ID: OLA) (formula: $C_{18}H_{34}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			20	18	2		
5	A	1	Total	C	O	0	0
			20	18	2		
5	A	1	Total	C	O	0	0
			18	16	2		
5	A	1	Total	C	O	0	0
			13	11	2		
5	A	1	Total	C	O	0	0
			13	11	2		
5	A	1	Total	C		0	0
			14	14			

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	48	Total	O	0	6
			48	48		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chloride pumping rhodopsin

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.48Å 51.18Å 78.33Å 90.00° 131.78° 90.00°	Depositor
Resolution (Å)	52.24 – 1.80 52.24 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (52.24-1.80) 99.1 (52.24-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.174 , 0.195 0.179 , 0.202	Depositor DCC
R_{free} test set	1456 reflections (1.45%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 68.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.045 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2312	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, BR, OLA, RET

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/2173	0.38	0/2960

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2099	0	2151	15	0
2	A	20	0	27	3	0
3	A	9	0	0	5	1
4	A	38	0	55	2	1
5	A	98	0	146	0	0
6	A	48	0	0	1	0
All	All	2312	0	2379	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:ASN:ND2	3:A:302[A]:BR:BR	2.58	0.92
4:A:311:OLC:O25	6:A:401[A]:HOH:O	1.98	0.81
1:A:49:ILE:HG23	1:A:112:ILE:HD13	1.64	0.79
2:A:301:RET:H8	2:A:301:RET:H161	1.69	0.75
1:A:103:ILE:HD12	1:A:136:ILE:HD11	1.76	0.67
1:A:168:ASN:CG	1:A:198[A]:MET:HE1	2.27	0.59
1:A:165:VAL:HG13	4:A:311:OLC:H8A	1.88	0.56
1:A:33:LEU:HD22	1:A:59:VAL:HG22	1.89	0.53
1:A:49:ILE:CG2	1:A:112:ILE:HD13	2.36	0.53
1:A:71:MET:HE3	1:A:89:THR:O	2.09	0.52
1:A:167:MET:HE1	1:A:201:TRP:CZ3	2.45	0.52
1:A:45:PRO:HD2	3:A:306[B]:BR:BR	2.66	0.51
1:A:189:THR:OG1	3:A:308[B]:BR:BR	2.77	0.48
2:A:301:RET:H161	2:A:301:RET:C8	2.39	0.48
1:A:54:SER:O	1:A:58:MET:HG2	2.18	0.44
1:A:44:ALA:HB1	3:A:306[B]:BR:BR	2.74	0.42
1:A:71:MET:HE1	1:A:94:TYR:CD2	2.55	0.42
2:A:301:RET:H181	2:A:301:RET:H7	1.81	0.41
1:A:143:GLN:HB3	3:A:304[B]:BR:BR	2.76	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:307[B]:BR:BR	4:A:311:OLC:O25[4_455]	1.64	0.56

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/296 (90%)	266 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/242 (93%)	223 (100%)	1 (0%)	84	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	153	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	68	GLN
1	A	86	GLN
1	A	152	GLN
1	A	259	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 9 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	RET	A	301	1	20,20,21	0.74	1 (5%)	27,27,28	1.39	5 (18%)
5	OLA	A	317	-	12,12,19	0.94	1 (8%)	12,12,19	1.14	1 (8%)
5	OLA	A	313	-	19,19,19	0.77	1 (5%)	19,19,19	0.97	1 (5%)
5	OLA	A	315	-	17,17,19	0.83	1 (5%)	17,17,19	0.91	0
5	OLA	A	314	-	19,19,19	0.84	1 (5%)	19,19,19	0.87	1 (5%)
5	OLA	A	316	-	12,12,19	0.95	1 (8%)	12,12,19	1.10	0
5	OLA	A	318	-	13,13,19	0.81	1 (7%)	12,12,19	0.53	0
4	OLC	A	311	-	24,24,24	0.82	2 (8%)	25,25,25	1.03	1 (4%)
4	OLC	A	312	-	12,12,24	1.12	2 (16%)	13,13,25	1.13	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	301	1	-	0/13/30/31	0/1/1/1
5	OLA	A	317	-	-	6/10/10/17	-
5	OLA	A	313	-	-	8/17/17/17	-
5	OLA	A	315	-	-	11/15/15/17	-
5	OLA	A	314	-	-	12/17/17/17	-
5	OLA	A	316	-	-	7/10/10/17	-
5	OLA	A	318	-	-	5/11/11/17	-
4	OLC	A	311	-	-	10/24/24/24	-
4	OLC	A	312	-	-	6/12/12/24	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	314	OLA	C10-C9	2.82	1.47	1.31
5	A	318	OLA	C10-C9	2.78	1.47	1.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	315	OLA	C10-C9	2.73	1.47	1.31
5	A	313	OLA	C10-C9	2.69	1.47	1.31
4	A	312	OLC	O20-C1	2.52	1.40	1.33
5	A	317	OLA	C9-C10	2.50	1.47	1.29
5	A	316	OLA	C9-C10	2.48	1.47	1.29
4	A	311	OLC	O20-C1	2.41	1.40	1.33
4	A	312	OLC	O20-C21	-2.17	1.40	1.45
4	A	311	OLC	O20-C21	-2.16	1.40	1.45
2	A	301	RET	C14-C13	2.08	1.35	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	RET	C3-C4-C5	-3.10	108.52	114.06
2	A	301	RET	C10-C11-C12	-2.98	114.56	123.20
4	A	311	OLC	O20-C1-C2	2.94	120.79	111.83
4	A	312	OLC	O20-C1-C2	2.87	120.59	111.83
2	A	301	RET	C7-C8-C9	-2.51	122.52	126.23
2	A	301	RET	C20-C13-C12	2.17	121.41	118.09
5	A	313	OLA	O2-C1-C2	2.15	120.79	114.00
2	A	301	RET	C2-C1-C6	2.12	113.52	110.44
5	A	317	OLA	C3-C2-C1	-2.10	109.03	114.51
5	A	314	OLA	O2-C1-C2	2.04	120.43	114.00

There are no chirality outliers.

All (65) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	311	OLC	C21-C22-C24-O25
4	A	312	OLC	C21-C22-C24-O25
4	A	311	OLC	O23-C22-C24-O25
5	A	313	OLA	C1-C2-C3-C4
5	A	314	OLA	C1-C2-C3-C4
5	A	316	OLA	C1-C2-C3-C4
4	A	312	OLC	O20-C21-C22-O23
5	A	316	OLA	C5-C6-C7-C8
4	A	312	OLC	O23-C22-C24-O25
4	A	311	OLC	C4-C5-C6-C7
4	A	311	OLC	C2-C1-O20-C21
4	A	312	OLC	C2-C3-C4-C5
4	A	311	OLC	O19-C1-O20-C21
5	A	314	OLA	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	314	OLA	C12-C13-C14-C15
4	A	311	OLC	C6-C7-C8-C9
4	A	311	OLC	C11-C12-C13-C14
5	A	314	OLA	C3-C4-C5-C6
5	A	315	OLA	C11-C12-C13-C14
5	A	316	OLA	C4-C5-C6-C7
5	A	315	OLA	C4-C5-C6-C7
5	A	317	OLA	C5-C6-C7-C8
4	A	312	OLC	O20-C21-C22-C24
5	A	314	OLA	C11-C10-C9-C8
5	A	315	OLA	C11-C10-C9-C8
5	A	318	OLA	C11-C10-C9-C8
5	A	314	OLA	C15-C16-C17-C18
5	A	316	OLA	C3-C4-C5-C6
5	A	313	OLA	C11-C10-C9-C8
5	A	313	OLA	C3-C4-C5-C6
5	A	314	OLA	C5-C6-C7-C8
5	A	316	OLA	C11-C10-C9-C8
5	A	317	OLA	C11-C10-C9-C8
4	A	312	OLC	C3-C4-C5-C6
5	A	315	OLA	C3-C4-C5-C6
5	A	315	OLA	C12-C13-C14-C15
5	A	318	OLA	C12-C13-C14-C15
5	A	315	OLA	C6-C7-C8-C9
5	A	316	OLA	C6-C7-C8-C9
5	A	314	OLA	C4-C5-C6-C7
5	A	318	OLA	C5-C6-C7-C8
5	A	314	OLA	C11-C12-C13-C14
5	A	313	OLA	C5-C6-C7-C8
5	A	315	OLA	C9-C10-C11-C12
5	A	313	OLA	C9-C10-C11-C12
4	A	311	OLC	C2-C3-C4-C5
5	A	317	OLA	O1-C1-C2-C3
5	A	318	OLA	C9-C10-C11-C12
5	A	317	OLA	O2-C1-C2-C3
5	A	313	OLA	O1-C1-C2-C3
5	A	314	OLA	C9-C10-C11-C12
5	A	315	OLA	O2-C1-C2-C3
4	A	311	OLC	C15-C16-C17-C18
5	A	315	OLA	O1-C1-C2-C3
5	A	315	OLA	C2-C3-C4-C5
4	A	311	OLC	C7-C8-C9-C10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	316	OLA	C7-C8-C9-C10
5	A	317	OLA	C7-C8-C9-C10
5	A	315	OLA	C1-C2-C3-C4
5	A	313	OLA	O2-C1-C2-C3
5	A	314	OLA	O1-C1-C2-C3
5	A	318	OLA	C3-C4-C5-C6
5	A	314	OLA	O2-C1-C2-C3
5	A	317	OLA	C4-C5-C6-C7
5	A	313	OLA	C7-C8-C9-C10


There are no ring outliers.


2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	RET	3	0
4	A	311	OLC	2	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/296 (88%)	1.09	41 (15%) 5 4	21, 45, 71, 90	8 (3%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	90	PHE	4.8
1	A	212	ALA	4.4
1	A	88	LEU	4.4
1	A	262	LEU	4.3
1	A	97	VAL	4.1
1	A	94	TYR	3.6
1	A	254	GLY	3.6
1	A	98	ASN	3.5
1	A	256	VAL	3.4
1	A	151	ALA	3.3
1	A	180	ALA	3.3
1	A	89	THR	3.2
1	A	251	ALA	3.1
1	A	150	ILE	3.0
1	A	122[A]	PHE	3.0
1	A	261	ALA	3.0
1	A	176	PHE	2.9
1	A	213	PHE	2.9
1	A	259	GLN	2.9
1	A	252	ALA	2.8
1	A	1	MET	2.8
1	A	108	LEU	2.8
1	A	211	PRO	2.8
1	A	148	ASP	2.6
1	A	152	GLN	2.5
1	A	78	TYR	2.5
1	A	241	MET	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	118	GLY	2.4
1	A	147	VAL	2.4
1	A	87	ASP	2.3
1	A	86	GLN	2.3
1	A	181	THR	2.3
1	A	195	TRP	2.3
1	A	130	LEU	2.2
1	A	91	SER	2.2
1	A	260	GLN	2.2
1	A	247	ILE	2.2
1	A	209	LEU	2.2
1	A	183	LEU	2.1
1	A	155	ILE	2.1
1	A	2	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BR	A	310[B]	1/1	-0.27	0.58	59,59,59,59	1
3	BR	A	309[B]	1/1	0.72	0.19	52,52,52,52	1
5	OLA	A	316	13/20	0.72	0.22	80,87,108,118	0
3	BR	A	303[B]	1/1	0.75	0.54	65,65,65,65	1
5	OLA	A	314	20/20	0.78	0.23	71,81,92,111	0
5	OLA	A	318	14/20	0.78	0.23	63,72,76,80	0
5	OLA	A	317	13/20	0.79	0.21	72,74,85,88	0
5	OLA	A	315	18/20	0.79	0.24	63,74,79,84	0
5	OLA	A	313	20/20	0.84	0.22	63,79,90,93	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	OLC	A	312	13/25	0.84	0.18	70,74,80,86	0
4	OLC	A	311	25/25	0.86	0.19	59,67,73,75	0
3	BR	A	308[B]	1/1	0.91	0.13	55,55,55,55	1
3	BR	A	307[B]	1/1	0.92	0.13	59,59,59,59	1
3	BR	A	304[B]	1/1	0.92	0.23	49,49,49,49	1
3	BR	A	305[B]	1/1	0.93	0.19	53,53,53,53	1
3	BR	A	302[A]	1/1	0.95	0.15	48,48,48,48	1
3	BR	A	306[B]	1/1	0.95	0.10	51,51,51,51	1
2	RET	A	301	20/21	0.96	0.09	34,37,43,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around OLA A 314:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 318:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLA A 317:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 315:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLA A 313:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 312:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 311:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RET A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.