



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:00 AM UTC

PDB ID : 1O94 / pdb_00001o94
Title : Ternary complex between trimethylamine dehydrogenase and electron transferring flavoprotein
Authors : Leys, D.; Basran, J.; Talfournier, F.; Sutcliffe, M.J.; Scrutton, N.S.
Deposited on : 2002-12-11
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	729	90%9%
1	B	729	% 88%11%
2	C	264	% 68%16%12%
2	E	264	3% 73%11%11%
3	D	320	6% 48%9%41%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	320	% 51% 7% 41% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SF4	B	1732	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 19925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIMETHYLAMINE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	729	Total	C	N	O	S	0	0	0
			5692	3589	996	1079	28			
1	B	729	Total	C	N	O	S	0	0	0
			5676	3583	994	1071	28			


- Molecule 2 is a protein called ELECTRON TRANSFER FLAVOPROTEIN BETA-SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	233	Total	C	N	O	S	0	0	0
			1749	1097	301	341	10			
2	E	236	Total	C	N	O	S	0	0	0
			1751	1102	299	340	10			

- Molecule 3 is a protein called ELECTRON TRANSFER FLAVOPROTEIN ALPHA-SUBUNIT.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	189	Total	C	N	O		0	0	0
			1354	857	230	267				
3	F	189	Total	C	N	O		0	0	0
			1376	870	232	274				

- Molecule 4 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
4	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Fe	S	0	0
			8	4	4		
6	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 7 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			23	10	5	7		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	718	Total	O	0	0
			718	718		
8	B	831	Total	O	0	0
			831	831		
8	C	178	Total	O	0	0
			178	178		
8	D	73	Total	O	0	0
			73	73		
8	E	198	Total	O	0	0
			198	198		
8	F	151	Total	O	0	0
			151	151		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TRIMETHYLAMINE DEHYDROGENASE



• Molecule 1: TRIMETHYLAMINE DEHYDROGENASE




• Molecule 2: ELECTRON TRANSFER FLAVOPROTEIN BETA-SUBUNIT



PHE
LYS
GLY
ALA


• Molecule 2: ELECTRON TRANSFER FLAVOPROTEIN BETA-SUBUNIT

Chain E: 3% 73% 11% 11%




• Molecule 3: ELECTRON TRANSFER FLAVOPROTEIN ALPHA-SUBUNIT

Chain D: 6% 48% 9% 41%



• Molecule 3: ELECTRON TRANSFER FLAVOPROTEIN ALPHA-SUBUNIT

Chain F: 51% 7% 41%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.93Å 211.54Å 125.97Å 90.00° 100.03° 90.00°	Depositor
Resolution (Å)	19.88 – 2.00 19.88 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.88-2.00) 94.7 (19.88-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.1.08	Depositor
R, R_{free}	0.172 , 0.212 0.175 , 0.212	Depositor DCC
R_{free} test set	9811 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.550	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19925	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ADP, SF4, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.25	13/5834 (0.2%)	1.16	12/7918 (0.2%)
1	B	1.32	15/5818 (0.3%)	1.18	12/7902 (0.2%)
2	C	1.45	6/1772 (0.3%)	1.32	16/2402 (0.7%)
2	E	1.20	5/1775 (0.3%)	1.23	10/2408 (0.4%)
3	D	0.88	0/1378	1.06	1/1884 (0.1%)
3	F	1.06	1/1400 (0.1%)	1.23	7/1913 (0.4%)
All	All	1.25	40/17977 (0.2%)	1.19	58/24427 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	C	0	1
3	D	0	1
3	F	0	1
All	All	0	4

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	54	ASP	C-N	35.93	1.83	1.33
1	A	568	MET	SD-CE	-10.87	1.52	1.79
1	B	253	MET	SD-CE	10.15	2.04	1.79
2	E	1	MET	SD-CE	-8.88	1.57	1.79
1	A	139	MET	SD-CE	-8.55	1.58	1.79

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	54	ASP	CA-C-N	18.07	156.06	121.54
2	C	54	ASP	C-N-CA	18.07	156.06	121.54
2	C	156	ASN	N-CA-C	11.13	126.38	113.02
3	F	49	VAL	N-CA-CB	10.76	117.28	110.50
3	F	48	PHE	CA-C-N	-9.11	114.50	120.24

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	167	TYR	Peptide
2	C	156	ASN	Peptide
3	D	109	LYS	Peptide
3	F	110	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5692	0	5444	46	1
1	B	5676	0	5423	62	1
2	C	1749	0	1722	48	1
2	E	1751	0	1716	46	1
3	D	1354	0	1311	26	0
3	F	1376	0	1358	18	0
4	A	31	0	18	0	0
4	B	31	0	18	1	0
5	A	27	0	12	0	0
5	B	27	0	12	0	0
6	A	8	0	0	0	0
6	B	8	0	0	2	0
7	C	23	0	12	2	0
7	E	23	0	12	1	0
8	A	718	0	0	22	0
8	B	831	0	0	31	0
8	C	178	0	0	10	0
8	D	73	0	0	2	0
8	E	198	0	0	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	F	151	0	0	3	0
All	All	19925	0	17058	230	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 230 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:110:LYS:CE	2:E:110:LYS:NZ	1.69	1.55
1:B:253:MET:CE	1:B:253:MET:SD	2.04	1.45
1:B:229:TYR:CD1	8:B:2325:HOH:O	1.75	1.35
2:C:54:ASP:C	2:C:56:ASP:N	1.84	1.35
1:A:11:GLU:HG2	8:A:2015:HOH:O	1.31	1.26

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:GLU:OE2	2:E:156:ASN:OD1[4_545]	1.97	0.23
1:B:11:GLU:OE2	2:C:156:ASN:OD1[4_556]	2.05	0.15

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	727/729 (100%)	702 (97%)	25 (3%)	0	100	100
1	B	727/729 (100%)	700 (96%)	27 (4%)	0	100	100
2	C	229/264 (87%)	222 (97%)	4 (2%)	3 (1%)	9	5
2	E	234/264 (89%)	230 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	187/320 (58%)	179 (96%)	7 (4%)	1 (0%)	24	21
3	F	187/320 (58%)	184 (98%)	3 (2%)	0	100	100
All	All	2291/2626 (87%)	2217 (97%)	70 (3%)	4 (0%)	43	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	56	ASP
2	C	157	LYS
3	D	31	SER
2	C	145	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	584/602 (97%)	573 (98%)	11 (2%)	50	56
1	B	578/602 (96%)	564 (98%)	14 (2%)	43	47
2	C	180/216 (83%)	169 (94%)	11 (6%)	17	14
2	E	177/216 (82%)	170 (96%)	7 (4%)	28	27
3	D	143/258 (55%)	133 (93%)	10 (7%)	14	10
3	F	150/258 (58%)	140 (93%)	10 (7%)	15	11
All	All	1812/2152 (84%)	1749 (96%)	63 (4%)	32	32

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	89	ASP
3	F	49	VAL
2	C	231	ARG
3	F	37	VAL
3	F	67	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	44	GLN
3	F	124	GLN
3	D	173	ASN
3	F	173	ASN
2	E	183	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	AMP	C	1236	-	25,25,25	1.27	4 (16%)	37,38,38	2.44	14 (37%)
6	SF4	A	1732	1	0,12,12	-	-	-	-	-
5	ADP	B	1731	-	28,29,29	1.27	5 (17%)	43,45,45	1.92	10 (23%)
5	ADP	A	1731	-	28,29,29	1.66	2 (7%)	43,45,45	1.74	8 (18%)
4	FMN	B	1730	1	33,33,33	1.49	7 (21%)	48,50,50	1.54	9 (18%)
7	AMP	E	1237	-	25,25,25	1.51	5 (20%)	37,38,38	2.46	15 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SF4	B	1732	1	0,12,12	-	-	-		
4	FMN	A	1730	1	33,33,33	1.36	6 (18%)	48,50,50	1.59	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	AMP	C	1236	-	-	4/10/26/26	0/3/3/3
6	SF4	A	1732	1	-	-	0/6/5/5
5	ADP	B	1731	-	-	2/16/32/32	0/3/3/3
5	ADP	A	1731	-	-	2/16/32/32	0/3/3/3
4	FMN	B	1730	1	-	1/18/18/18	0/3/3/3
7	AMP	E	1237	-	-	4/10/26/26	0/3/3/3
6	SF4	B	1732	1	-	-	0/6/5/5
4	FMN	A	1730	1	-	1/18/18/18	0/3/3/3

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1731	ADP	PA-O3A	7.18	1.67	1.59
7	E	1237	AMP	C2-N3	4.31	1.41	1.33
4	A	1730	FMN	C4A-N5	3.25	1.37	1.30
7	C	1236	AMP	C2-N3	3.21	1.39	1.33
7	E	1237	AMP	C2-N1	3.09	1.39	1.33

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	1236	AMP	N3-C2-N1	-6.80	118.29	128.58
7	C	1236	AMP	O2P-P-O5'	6.56	123.76	106.67
7	E	1237	AMP	N3-C2-N1	-6.36	118.96	128.58
5	B	1731	ADP	N9-C8-N7	-5.44	106.22	113.94
7	E	1237	AMP	O2P-P-O5'	5.22	120.29	106.67

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	C	1236	AMP	C5'-O5'-P-O2P
7	C	1236	AMP	C5'-O5'-P-O3P
7	E	1237	AMP	C5'-O5'-P-O3P
7	C	1236	AMP	C5'-O5'-P-O1P
7	E	1237	AMP	C5'-O5'-P-O1P


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	1236	AMP	2	0
4	B	1730	FMN	1	0
7	E	1237	AMP	1	0
6	B	1732	SF4	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand AMP C 1236





Ligand ADP B 1731






Ligand FMN B 1730







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	729/729 (100%)	-0.44	2 (0%) 90 89	21, 28, 40, 47	0
1	B	729/729 (100%)	-0.53	4 (0%) 87 87	18, 26, 38, 53	0
2	C	233/264 (88%)	0.04	3 (1%) 75 74	25, 35, 56, 66	0
2	E	236/264 (89%)	-0.01	7 (2%) 52 51	25, 33, 50, 58	0
3	D	189/320 (59%)	0.68	20 (10%) 11 10	32, 50, 66, 76	0
3	F	189/320 (59%)	0.02	3 (1%) 70 70	27, 36, 52, 63	0
All	All	2305/2626 (87%)	-0.25	39 (1%) 69 68	18, 30, 53, 76	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	22	ASP	4.1
3	D	174	VAL	3.5
2	E	54	ASP	3.4
2	E	23	GLY	3.4
3	F	32	GLY	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	SF4	B	1732	8/8	0.89	0.15	28,29,43,49	0
4	FMN	B	1730	31/31	0.96	0.07	19,23,26,28	0
4	FMN	A	1730	31/31	0.96	0.07	18,23,25,27	0
7	AMP	C	1236	23/23	0.96	0.06	26,30,37,38	0
7	AMP	E	1237	23/23	0.97	0.07	23,28,33,35	0
5	ADP	A	1731	27/27	0.99	0.04	23,28,30,31	0
5	ADP	B	1731	27/27	0.99	0.04	20,24,28,29	0
6	SF4	A	1732	8/8	0.99	0.07	29,31,32,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FMN B 1730:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FMN A 1730:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP C 1236:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP E 1237:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP A 1731:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP B 1731:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.