



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:52 PM UTC

PDB ID : 4O9D / pdb_00004o9d
Title : Structure of Dos1 propeller
Authors : Kucsu, C.; Schalch, T.; Joshua-Tor, L.
Deposited on : 2014-01-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	7% 77% 14% 9%
1	B	428	7% 76% 14% • 8%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rik1-associated factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	392	Total	C	N	O	S	0	0	0
			3047	1936	500	595	16			
1	A	391	Total	C	N	O	S	0	0	0
			3037	1930	496	594	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	211	GLY	-	expression tag	UNP O74910
B	212	ALA	-	expression tag	UNP O74910
A	211	GLY	-	expression tag	UNP O74910
A	212	ALA	-	expression tag	UNP O74910

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	186	Total O 186 186	0	0
4	A	163	Total O 163 163	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rik1-associated factor 1



• Molecule 1: Rik1-associated factor 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	121.15Å 102.79Å 95.10Å 90.00° 122.36° 90.00°	Depositor
Resolution (Å)	45.87 – 2.00 45.87 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (45.87-2.00) 99.3 (45.87-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.00Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.213 , 0.250 0.213 , 0.250	Depositor DCC
R_{free} test set	1734 reflections (2.63%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6470	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3103	0.80	1/4199 (0.0%)
1	B	0.66	2/3114 (0.1%)	0.84	4/4214 (0.1%)
All	All	0.63	2/6217 (0.0%)	0.82	5/8413 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	467	ASN	C-O	-5.95	1.16	1.23
1	B	466	SER	C-O	-5.41	1.17	1.24

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	624	ASN	N-CA-C	7.76	122.13	113.21
1	B	238	GLU	N-CA-C	-6.79	104.30	112.59
1	B	555	GLN	CA-C-N	-6.01	113.57	119.76
1	B	555	GLN	C-N-CA	-6.01	113.57	119.76
1	B	270	GLY	N-CA-C	-5.20	106.12	112.68

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3037	0	2919	38	0
1	B	3047	0	2924	43	0
2	A	20	0	30	0	0
2	B	16	0	24	0	0
3	B	1	0	0	0	0
4	A	163	0	0	1	0
4	B	186	0	0	2	0
All	All	6470	0	5897	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:467:ASN:HB2	1:B:486:THR:O	1.81	0.81
1:B:238:GLU:HG3	1:B:631:ILE:HD11	1.70	0.72
1:A:467:ASN:HB3	1:A:486:THR:O	1.95	0.67
1:A:294:VAL:HG12	1:A:294:VAL:O	1.97	0.63
1:A:474:ASP:HB2	1:A:481:CYS:SG	2.40	0.61
1:B:241:GLY:HA2	1:B:262:GLN:HG2	1.82	0.60
1:B:240:SER:HB3	1:B:267:ASN:HB2	1.84	0.60
1:A:469:ARG:HH12	1:A:483:ASN:HD21	1.50	0.57
1:A:458:PHE:CE2	1:A:473:LEU:HD13	2.40	0.56
1:B:290:LYS:HE3	1:B:320:ASP:HA	1.88	0.55
1:B:305:LEU:HB3	1:B:316:THR:HG22	1.89	0.55
1:B:617:SER:C	1:B:619:GLY:H	2.15	0.55
1:B:452:HIS:CG	1:B:453:PRO:HD2	2.43	0.54
1:B:608:CYS:SG	1:B:612:ASP:HB2	2.48	0.53
1:B:240:SER:CB	1:B:267:ASN:HB2	2.38	0.53
1:A:292:ASN:N	1:A:296:GLU:O	2.36	0.53
1:A:469:ARG:NH2	1:A:483:ASN:OD1	2.42	0.53
1:A:222:MET:HA	1:A:228:PHE:CE2	2.44	0.53
1:B:294:VAL:HG12	1:B:294:VAL:O	2.09	0.53
1:A:405:SER:OG	1:A:406:SER:N	2.40	0.53
1:A:294:VAL:O	1:A:294:VAL:CG1	2.56	0.52
1:B:368:THR:HG22	1:B:375:ILE:HD13	1.92	0.51
1:B:588:MET:SD	1:B:606:VAL:HG12	2.50	0.51
1:A:414:ASP:OD1	1:A:414:ASP:C	2.53	0.51
1:A:291:LEU:HD23	1:A:297:LYS:CA	2.41	0.50
1:B:417:HIS:O	1:B:418:THR:C	2.55	0.50
1:A:551:MET:HE1	1:A:604:LEU:HD11	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:297:LYS:NZ	4:B:867:HOH:O	2.45	0.49
1:B:250:PRO:HD2	1:B:311:ASN:OD1	2.13	0.48
1:B:475:THR:O	1:B:475:THR:CG2	2.61	0.48
1:A:292:ASN:ND2	1:A:296:GLU:HB2	2.29	0.48
1:B:270:GLY:HA2	1:B:273:TRP:HE1	1.78	0.48
1:A:240:SER:HB3	1:A:271:ASN:HB3	1.96	0.48
1:B:327:ASP:OD2	1:B:331:ASN:HB2	2.14	0.47
1:A:255:PHE:CZ	1:A:275:GLY:HA3	2.49	0.47
1:B:240:SER:HB3	1:B:267:ASN:CB	2.44	0.47
1:B:291:LEU:HA	1:B:297:LYS:HA	1.96	0.47
1:A:225:GLU:OE2	1:A:555:GLN:HG2	2.15	0.47
1:A:327:ASP:OD1	1:A:327:ASP:C	2.57	0.47
1:B:452:HIS:CD2	1:B:453:PRO:HD2	2.50	0.46
1:B:291:LEU:HD13	1:B:291:LEU:N	2.30	0.46
1:A:238:GLU:HG3	1:A:631:ILE:HD11	1.97	0.46
1:B:401:PHE:HB2	1:B:475:THR:HG22	1.98	0.46
1:B:475:THR:O	1:B:475:THR:HG22	2.14	0.46
1:B:439:PHE:CD1	1:B:473:LEU:CD2	2.99	0.45
1:B:617:SER:C	1:B:619:GLY:N	2.73	0.45
1:B:232:PRO:HB3	1:B:626:PHE:CZ	2.52	0.45
1:B:262:GLN:OE1	1:B:262:GLN:HA	2.16	0.45
1:A:402:LEU:O	1:A:404:ARG:HD2	2.16	0.45
1:A:452:HIS:CG	1:A:453:PRO:HD2	2.52	0.45
1:A:594:TYR:CD1	1:A:595:GLY:N	2.86	0.44
1:A:442:ALA:HB1	4:A:922:HOH:O	2.17	0.44
1:B:240:SER:CB	1:B:267:ASN:CB	2.96	0.44
1:A:239:THR:CG2	1:A:241:GLY:O	2.66	0.44
1:B:267:ASN:ND2	4:B:970:HOH:O	2.35	0.44
1:A:401:PHE:HA	1:A:475:THR:HG21	2.00	0.44
1:B:285:LEU:HD22	1:B:330:GLY:HA2	1.99	0.44
1:B:506:SER:HB3	1:B:549:ILE:HG21	1.99	0.44
1:A:222:MET:HA	1:A:228:PHE:CD2	2.53	0.43
1:A:595:GLY:HA2	1:A:605:THR:O	2.18	0.43
1:A:365:LEU:O	1:A:378:SER:HA	2.19	0.43
1:B:402:LEU:HA	1:B:455:GLN:OE1	2.19	0.43
1:B:469:ARG:NH1	1:B:485:THR:HG22	2.33	0.43
1:A:617:SER:O	1:A:618:LEU:HB2	2.19	0.43
1:B:561:PHE:CE2	1:B:563:THR:CG2	3.02	0.43
1:A:221:LEU:HD12	1:A:221:LEU:O	2.19	0.42
1:A:408:LYS:HB3	1:A:427:LEU:HD11	2.01	0.42
1:A:291:LEU:HD23	1:A:297:LYS:HA	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:288:HIS:HB3	1:B:320:ASP:CG	2.44	0.42
1:B:501:ILE:O	1:B:501:ILE:HG22	2.18	0.42
1:B:497:SER:HA	1:B:554:TRP:CE2	2.54	0.42
1:A:290:LYS:HE3	1:A:320:ASP:HA	2.00	0.42
1:A:534:MET:HE1	1:A:609:VAL:HG21	2.02	0.42
1:A:501:ILE:HG22	1:A:502:LEU:HD23	2.02	0.41
1:B:320:ASP:O	1:B:321:ASN:HB2	2.19	0.41
1:A:459:VAL:HG13	1:A:496:ILE:HD11	2.03	0.41
1:A:509:ASP:CG	1:A:511:GLN:HG2	2.47	0.40
1:B:235:HIS:HE1	1:B:612:ASP:OD1	2.04	0.40
1:B:509:ASP:CG	1:B:511:GLN:HG2	2.47	0.40
1:B:556:PRO:O	1:B:557:LYS:HB2	2.22	0.40
1:A:310:GLY:O	1:A:311:ASN:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/428 (90%)	370 (97%)	13 (3%)	0	100	100
1	B	384/428 (90%)	372 (97%)	12 (3%)	0	100	100
All	All	767/856 (90%)	742 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/365 (92%)	332 (99%)	3 (1%)	70	78
1	B	334/365 (92%)	327 (98%)	7 (2%)	47	52
All	All	669/730 (92%)	659 (98%)	10 (2%)	57	64

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	245	CYS
1	B	268	ARG
1	B	291	LEU
1	B	293	GLN
1	B	305	LEU
1	B	329	GLU
1	B	466	SER
1	A	305	LEU
1	A	420	ASN
1	A	469	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	278	HIS
1	B	583	GLN
1	A	311	ASN
1	A	331	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	704	-	3,3,3	0.49	0	2,2,2	0.64	0
2	EDO	B	701	-	3,3,3	0.48	0	2,2,2	0.20	0
2	EDO	A	703	-	3,3,3	0.45	0	2,2,2	0.28	0
2	EDO	A	705	-	3,3,3	0.45	0	2,2,2	0.34	0
2	EDO	A	702	-	3,3,3	0.47	0	2,2,2	0.30	0
2	EDO	A	701	-	3,3,3	0.57	0	2,2,2	0.12	0
2	EDO	B	702	-	3,3,3	0.42	0	2,2,2	0.46	0
2	EDO	A	704	-	3,3,3	0.47	0	2,2,2	0.04	0
2	EDO	B	703	-	3,3,3	0.42	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	704	-	-	1/1/1/1	-
2	EDO	B	701	-	-	0/1/1/1	-
2	EDO	A	703	-	-	1/1/1/1	-
2	EDO	A	705	-	-	1/1/1/1	-
2	EDO	A	702	-	-	1/1/1/1	-
2	EDO	A	701	-	-	0/1/1/1	-
2	EDO	B	702	-	-	1/1/1/1	-
2	EDO	A	704	-	-	0/1/1/1	-
2	EDO	B	703	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	702	EDO	O1-C1-C2-O2
2	A	702	EDO	O1-C1-C2-O2
2	A	705	EDO	O1-C1-C2-O2
2	B	703	EDO	O1-C1-C2-O2
2	B	704	EDO	O1-C1-C2-O2
2	A	703	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/428 (91%)	0.63	28 (7%) 21 20	20, 32, 52, 61	0
1	B	392/428 (91%)	0.56	29 (7%) 20 19	19, 30, 54, 68	0
All	All	783/856 (91%)	0.59	57 (7%) 21 19	19, 31, 53, 68	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	261	ALA	5.8
1	B	291	LEU	5.8
1	A	221	LEU	4.8
1	B	294	VAL	4.7
1	A	535	ILE	4.4
1	B	264	GLY	4.3
1	B	417	HIS	4.2
1	A	240	SER	4.1
1	B	267	ASN	4.0
1	B	623	GLY	4.0
1	A	623	GLY	3.8
1	A	291	LEU	3.8
1	B	262	GLN	3.8
1	A	465	LEU	3.6
1	A	618	LEU	3.6
1	B	263	ASP	3.6
1	B	265	ALA	3.5
1	B	295	GLY	3.4
1	B	266	TYR	3.4
1	B	268	ARG	3.3
1	B	418	THR	3.3
1	B	270	GLY	3.3
1	B	619	GLY	3.3
1	B	269	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	239	THR	3.2
1	A	416	PHE	3.2
1	B	296	GLU	3.2
1	A	224	GLY	3.1
1	B	292	ASN	3.1
1	A	417	HIS	3.0
1	B	290	LYS	2.9
1	A	632	ILE	2.8
1	B	293	GLN	2.7
1	A	611	GLY	2.7
1	A	467	ASN	2.7
1	B	251	LEU	2.6
1	A	270	GLY	2.6
1	A	547	ALA	2.6
1	B	229	PHE	2.6
1	A	294	VAL	2.6
1	A	418	THR	2.6
1	A	293	GLN	2.5
1	A	420	ASN	2.4
1	A	631	ILE	2.4
1	A	482	LEU	2.4
1	A	292	ASN	2.4
1	B	298	GLU	2.3
1	B	240	SER	2.3
1	A	236	PHE	2.3
1	A	415	SER	2.2
1	A	466	SER	2.2
1	B	239	THR	2.2
1	B	297	LYS	2.2
1	B	299	TYR	2.1
1	A	522	LYS	2.1
1	A	350	ASP	2.1
1	B	521	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	703	4/4	0.80	0.21	30,33,35,36	0
2	EDO	A	705	4/4	0.81	0.17	50,51,54,55	0
2	EDO	A	702	4/4	0.85	0.12	35,38,46,57	0
2	EDO	A	704	4/4	0.85	0.15	35,41,42,47	0
2	EDO	B	704	4/4	0.85	0.19	31,35,36,40	0
2	EDO	B	702	4/4	0.87	0.13	42,48,49,51	0
2	EDO	B	701	4/4	0.90	0.11	35,38,39,50	0
2	EDO	A	703	4/4	0.93	0.15	31,37,41,42	0
2	EDO	A	701	4/4	0.94	0.09	33,34,35,36	0
3	CL	B	705	1/1	0.97	0.07	34,34,34,34	0

6.5 Other polymers

There are no such residues in this entry.