



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 01:08 PM UTC

PDB ID : 6O9E / pdb_00006o9e
Title : Structure of HIV-1 Reverse Transcriptase in complex with DNA and INDOPY-1
Authors : Ruiz, F.X.; Hoang, A.; Das, K.; Arnold, E.
Deposited on : 2019-03-13
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	0% 74% 25%
1	C	555	0% 75% 24% •
2	B	429	2% 80% 17% •
2	D	429	3% 78% 17% •
3	E	38	32% 55% 5% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	38	
4	G	2	
4	H	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLC	H	1	-	-	X	-
5	SO4	B	502	-	-	X	-
8	PEG	B	506	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 17591 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase p66.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	553	Total	C	N	O	S	0	1	0
			4510	2920	751	832	7			
1	C	553	Total	C	N	O	S	0	0	0
			4504	2916	750	831	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	SER	CYS	engineered mutation	UNP P03366
A	498	ASN	ASP	engineered mutation	UNP P03366
C	280	SER	CYS	engineered mutation	UNP P03366
C	498	ASN	ASP	engineered mutation	UNP P03366

- Molecule 2 is a protein called Reverse transcriptase p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	415	Total	C	N	O	S	0	1	0
			3434	2239	567	621	7			
2	D	410	Total	C	N	O	S	0	2	0
			3409	2225	562	615	7			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366
D	0	GLY	-	expression tag	UNP P03366
D	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is a DNA chain called DNA (38-MER).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	35	Total	C	N	O	P	0	0	0
			720	341	129	215	35			
3	E	35	Total	C	N	O	P	0	0	0
			720	341	129	215	35			

- Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	G	2	Total	C	O	0	0	0
			23	12	11			
4	H	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is AMMONIUM ION (CCD ID: NH4) (formula: H_4N).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	H	N	0	0
			5	4	1		
6	C	1	Total	H	N	0	0
			5	4	1		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 9 is 5-methyl-1-(4-nitrophenyl)-2-oxo-2,5-dihydro-1H-pyrido[3,2-b]indole-3-carbonitrile (CCD ID: IY1) (formula: C₁₉H₁₂N₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	F	1	Total	C	N	O	0	0
			26	19	4	3		
9	E	1	Total	C	N	O	0	0
			26	19	4	3		

- Molecule 10 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	E	1	Total	C	O	0	0
			13	8	5		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	32	Total	O	0	0
			32	32		
11	B	35	Total	O	0	0
			35	35		
11	C	21	Total	O	0	0
			21	21		
11	D	15	Total	O	0	0
			15	15		
11	F	3	Total	O	0	0
			3	3		
11	E	5	Total	O	0	0
			5	5		


- Molecule 2: Reverse transcriptase p51




- Molecule 2: Reverse transcriptase p51




- Molecule 3: DNA (38-MER)



- Molecule 3: DNA (38-MER)



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain H:



GLU1
FRU2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.76Å 127.66Å 131.61Å 90.00° 101.50° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 50.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.40) 99.8 (50.00-2.40)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.40Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.203 , 0.242 0.210 , 0.249	Depositor DCC
R_{free} test set	2000 reflections (1.76%)	wwPDB-VP
Wilson B-factor (Å ²)	73.6	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 88.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17591	wwPDB-VP
Average B, all atoms (Å ²)	126.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, NH4, SO4, GLC, OMC, PG4, GOL, IY1, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.09	0/4631	0.27	0/6291
1	C	0.10	0/4622	0.30	2/6279 (0.0%)
2	B	0.09	0/3534	0.26	0/4800
2	D	0.10	0/3515	0.27	0/4775
3	E	0.18	0/759	0.37	0/1170
3	F	0.84	6/759 (0.8%)	0.46	0/1170
All	All	0.20	6/17820 (0.0%)	0.29	2/24485 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	-1	DT	C3'-C2'	-15.31	1.22	1.52
3	F	-1	DT	C5-C6	7.28	1.48	1.34
3	F	-1	DT	C2-N3	6.12	1.50	1.37
3	F	-1	DT	N1-C2	5.59	1.49	1.38
3	F	-1	DT	O4'-C1'	-5.42	1.31	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	419	THR	CA-C-N	5.16	123.43	119.66
1	C	419	THR	C-N-CA	5.16	123.43	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4510	0	4575	122	0
1	C	4504	0	4567	100	0
2	B	3434	0	3465	62	0
2	D	3409	0	3438	63	0
3	E	720	0	398	27	0
3	F	720	0	398	16	0
4	G	23	0	21	5	0
4	H	23	0	21	9	0
5	A	10	0	0	0	0
5	B	10	0	0	3	0
5	C	10	0	0	0	0
6	A	1	4	0	0	0
6	C	1	4	0	0	0
7	B	12	0	16	1	0
7	C	6	0	8	1	0
8	B	14	0	20	6	0
9	E	26	0	0	1	0
9	F	26	0	0	2	0
10	E	13	0	18	2	0
11	A	32	0	0	0	0
11	B	35	0	0	1	0
11	C	21	0	0	1	0
11	D	15	0	0	1	0
11	E	5	0	0	1	0
11	F	3	0	0	0	0
All	All	17583	8	16945	373	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 373 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:70:LYS:HE3	2:B:226:PRO:HD2	1.38	1.03
2:D:257:ILE:HD12	2:D:293:ILE:HD11	1.44	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:THR:HG22	1:A:143:ARG:HG2	1.46	0.96
3:E:4:OMC:H5	3:E:30:DG:H1	1.16	0.91
1:C:246:LEU:HD13	1:C:307:ARG:HD3	1.54	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	552/555 (100%)	534 (97%)	18 (3%)	0	100	100
1	C	551/555 (99%)	536 (97%)	14 (2%)	1 (0%)	43	58
2	B	411/429 (96%)	398 (97%)	13 (3%)	0	100	100
2	D	408/429 (95%)	392 (96%)	16 (4%)	0	100	100
All	All	1922/1968 (98%)	1860 (97%)	61 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	543	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/495 (100%)	493 (99%)	3 (1%)	78	89
1	C	495/495 (100%)	492 (99%)	3 (1%)	78	89
2	B	377/390 (97%)	376 (100%)	1 (0%)	86	93
2	D	374/390 (96%)	371 (99%)	3 (1%)	73	86
All	All	1742/1770 (98%)	1732 (99%)	10 (1%)	78	89

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	266	TRP
2	D	303	LEU
2	D	427	TYR
2	B	428	GLN
1	C	109	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	524	GLN
2	D	147	ASN
2	D	96	HIS
1	C	147	ASN
1	C	520	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OMC	F	4	3	19,22,23	3.09	7 (36%)	25,31,34	0.74	0
3	OMC	E	4	3	19,22,23	3.04	7 (36%)	25,31,34	1.00	2 (8%)
3	OMC	F	2	3	19,22,23	3.08	7 (36%)	25,31,34	0.77	0
3	OMC	E	2	3	19,22,23	3.07	7 (36%)	25,31,34	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	F	4	3	-	1/9/27/28	0/2/2/2
3	OMC	E	4	3	-	3/9/27/28	0/2/2/2
3	OMC	F	2	3	-	1/9/27/28	0/2/2/2
3	OMC	E	2	3	-	1/9/27/28	0/2/2/2

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	4	OMC	C2-N3	6.62	1.49	1.36
3	F	2	OMC	C2-N3	6.60	1.49	1.36
3	F	4	OMC	C2-N3	6.57	1.49	1.36
3	E	2	OMC	C2-N3	6.54	1.49	1.36
3	F	4	OMC	C6-C5	6.50	1.50	1.35

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	4	OMC	C6-C5-C4	2.25	121.22	117.53
3	E	4	OMC	O2-C2-N3	-2.10	119.02	122.33

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	2	OMC	C1'-C2'-O2'-CM2
3	E	2	OMC	C1'-C2'-O2'-CM2
3	F	4	OMC	C1'-C2'-O2'-CM2
3	E	4	OMC	O4'-C1'-N1-C2
3	E	4	OMC	O4'-C1'-N1-C6

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	4	OMC	2	0
3	E	4	OMC	3	0
3	F	2	OMC	4	0
3	E	2	OMC	2	0

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	G	1	4	11,11,12	0.28	0	15,15,17	1.19	1 (6%)
4	FRU	G	2	4	11,12,12	0.51	0	10,18,18	0.99	1 (10%)
4	GLC	H	1	4	11,11,12	0.29	0	15,15,17	1.27	3 (20%)
4	FRU	H	2	4	11,12,12	0.49	0	10,18,18	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	G	1	4	-	0/2/19/22	0/1/1/1
4	FRU	G	2	4	-	0/5/24/24	0/1/1/1
4	GLC	H	1	4	-	2/2/19/22	0/1/1/1
4	FRU	H	2	4	-	5/5/24/24	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	1	GLC	C1-O5-C5	3.27	116.58	112.19
4	H	1	GLC	C1-O5-C5	2.74	115.86	112.19
4	H	1	GLC	C1-C2-C3	-2.26	106.35	109.64
4	G	2	FRU	O1-C1-C2	-2.05	107.15	111.67
4	H	1	GLC	O5-C5-C6	2.03	111.62	107.66

There are no chirality outliers.

5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	1	GLC	O5-C5-C6-O6
4	H	1	GLC	C4-C5-C6-O6
4	H	2	FRU	C4-C5-C6-O6
4	H	2	FRU	O5-C5-C6-O6
4	H	2	FRU	O1-C1-C2-O5

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	1	GLC	4	0
4	H	1	GLC	6	0
4	G	2	FRU	5	0
4	H	2	FRU	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 16 ligands modelled in this entry, 2 are modelled with single atom - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	IY1	F	101	-	29,29,29	0.51	0	35,43,43	1.19	2 (5%)
8	PEG	B	507	-	6,6,6	0.11	0	5,5,5	0.08	0
10	PG4	E	101	-	12,12,12	0.54	0	11,11,11	0.23	0
5	SO4	A	601	-	4,4,4	0.23	0	6,6,6	0.07	0
5	SO4	C	601	-	4,4,4	0.23	0	6,6,6	0.07	0
7	GOL	C	603	-	5,5,5	0.35	0	5,5,5	0.27	0
9	IY1	E	102	-	29,29,29	0.50	0	35,43,43	1.17	2 (5%)
7	GOL	B	505	-	5,5,5	0.37	0	5,5,5	0.30	0
5	SO4	B	503	-	4,4,4	0.26	0	6,6,6	0.12	0
5	SO4	C	602	-	4,4,4	0.38	0	6,6,6	0.08	0
5	SO4	B	502	-	4,4,4	0.42	0	6,6,6	0.08	0
7	GOL	B	504	-	5,5,5	0.07	0	5,5,5	0.29	0
8	PEG	B	506	-	6,6,6	0.13	0	5,5,5	0.07	0
5	SO4	A	602	-	4,4,4	0.23	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	IY1	F	101	-	-	0/6/10/10	0/4/4/4
10	PG4	E	101	-	-	6/10/10/10	-
9	IY1	E	102	-	-	0/6/10/10	0/4/4/4
7	GOL	C	603	-	-	4/4/4/4	-
7	GOL	B	505	-	-	0/4/4/4	-
8	PEG	B	507	-	-	1/4/4/4	-
7	GOL	B	504	-	-	4/4/4/4	-
8	PEG	B	506	-	-	2/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	F	101	IY1	C9-C8-N	5.93	132.53	128.97
9	E	102	IY1	C9-C8-N	5.79	132.45	128.97
9	F	101	IY1	C6-C7-N1	2.56	132.68	126.88
9	E	102	IY1	C6-C7-N1	2.52	132.57	126.88

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	504	GOL	O1-C1-C2-C3
7	B	504	GOL	C1-C2-C3-O3
7	B	504	GOL	O2-C2-C3-O3
7	C	603	GOL	O1-C1-C2-C3
7	C	603	GOL	C1-C2-C3-O3


There are no ring outliers.

8 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	F	101	IY1	2	0
8	B	507	PEG	1	0
10	E	101	PG4	2	0
7	C	603	GOL	1	0
9	E	102	IY1	1	0
5	B	502	SO4	3	0
7	B	504	GOL	1	0
8	B	506	PEG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand IY1 F 101




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	553/555 (99%)	0.10	4 (0%) 84 81	39, 114, 216, 292	1 (0%)
1	C	553/555 (99%)	0.14	8 (1%) 73 69	56, 137, 230, 292	0
2	B	415/429 (96%)	-0.01	7 (1%) 69 65	60, 101, 176, 249	0
2	D	410/429 (95%)	0.20	13 (3%) 50 46	51, 123, 207, 256	2 (0%)
3	E	33/38 (86%)	-0.39	0 100 100	81, 123, 187, 247	0
3	F	33/38 (86%)	-0.45	0 100 100	91, 145, 177, 223	0
All	All	1997/2044 (97%)	0.09	32 (1%) 70 66	39, 117, 217, 292	3 (0%)

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	226	PRO	4.0
2	D	266	TRP	3.8
2	B	266	TRP	3.5
1	C	544	GLY	3.1
2	B	228	LEU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OMC	F	2	21/22	0.93	0.07	135,146,151,153	0
3	OMC	F	4	21/22	0.93	0.09	114,128,141,154	0
3	OMC	E	4	21/22	0.93	0.11	71,84,97,118	0
3	OMC	E	2	21/22	0.95	0.08	92,106,115,118	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	FRU	G	2	12/12	0.81	0.18	138,151,156,161	0
4	FRU	H	2	12/12	0.83	0.11	135,157,166,170	0
4	GLC	G	1	11/12	0.85	0.12	102,114,127,132	0
4	GLC	H	1	11/12	0.88	0.12	132,137,151,157	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NH4	A	603	1/1	0.50	0.28	130,156,156,156	0
5	SO4	C	602	5/5	0.61	0.08	188,191,192,196	0
5	SO4	A	601	5/5	0.63	0.11	166,171,173,176	0
9	IY1	F	101	26/26	0.64	0.14	169,182,210,216	0
6	NH4	C	604	1/1	0.73	0.26	104,125,125,125	0
5	SO4	B	503	5/5	0.73	0.10	190,190,193,194	0
5	SO4	C	601	5/5	0.76	0.10	159,161,162,163	0
10	PG4	E	101	13/13	0.81	0.15	125,144,149,150	0
9	IY1	E	102	26/26	0.85	0.11	144,158,178,183	0
7	GOL	B	504	6/6	0.85	0.13	91,113,121,126	0
7	GOL	C	603	6/6	0.87	0.10	115,119,121,124	0
8	PEG	B	507	7/7	0.88	0.42	168,187,201,203	0
5	SO4	B	502	5/5	0.89	0.11	137,137,138,150	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	B	505	6/6	0.90	0.22	89,95,106,107	0
8	PEG	B	506	7/7	0.91	0.12	92,100,103,109	0
5	SO4	A	602	5/5	0.92	0.09	178,182,184,184	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IY1 F 101:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IY1 E 102:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.