



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:13 PM UTC

PDB ID : 2OA1 / pdb_00002oa1
Title : Crystal Structure of RebH, a FAD-dependent halogenase from *Lechevalieria aerocolonigenes*, the L-Tryptophan with FAD complex
Authors : Bitto, E.; Bingman, C.A.; Singh, S.; Phillips Jr., G.N.
Deposited on : 2006-12-14
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	550	
1	B	550	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9627 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan halogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4234	2691	737	787	19			
1	B	527	Total	C	N	O	S	0	0	0
			4234	2691	737	787	19			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q8KHZ8
A	-18	GLY	-	expression tag	UNP Q8KHZ8
A	-17	SER	-	expression tag	UNP Q8KHZ8
A	-16	SER	-	expression tag	UNP Q8KHZ8
A	-15	HIS	-	expression tag	UNP Q8KHZ8
A	-14	HIS	-	expression tag	UNP Q8KHZ8
A	-13	HIS	-	expression tag	UNP Q8KHZ8
A	-12	HIS	-	expression tag	UNP Q8KHZ8
A	-11	HIS	-	expression tag	UNP Q8KHZ8
A	-10	HIS	-	expression tag	UNP Q8KHZ8
A	-9	SER	-	expression tag	UNP Q8KHZ8
A	-8	SER	-	expression tag	UNP Q8KHZ8
A	-7	GLY	-	expression tag	UNP Q8KHZ8
A	-6	LEU	-	expression tag	UNP Q8KHZ8
A	-5	VAL	-	expression tag	UNP Q8KHZ8
A	-4	PRO	-	expression tag	UNP Q8KHZ8
A	-3	ARG	-	expression tag	UNP Q8KHZ8
A	-2	GLY	-	expression tag	UNP Q8KHZ8
A	-1	SER	-	expression tag	UNP Q8KHZ8
A	0	HIS	-	expression tag	UNP Q8KHZ8
B	-19	MET	-	expression tag	UNP Q8KHZ8
B	-18	GLY	-	expression tag	UNP Q8KHZ8
B	-17	SER	-	expression tag	UNP Q8KHZ8
B	-16	SER	-	expression tag	UNP Q8KHZ8
B	-15	HIS	-	expression tag	UNP Q8KHZ8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q8KHZ8
B	-13	HIS	-	expression tag	UNP Q8KHZ8
B	-12	HIS	-	expression tag	UNP Q8KHZ8
B	-11	HIS	-	expression tag	UNP Q8KHZ8
B	-10	HIS	-	expression tag	UNP Q8KHZ8
B	-9	SER	-	expression tag	UNP Q8KHZ8
B	-8	SER	-	expression tag	UNP Q8KHZ8
B	-7	GLY	-	expression tag	UNP Q8KHZ8
B	-6	LEU	-	expression tag	UNP Q8KHZ8
B	-5	VAL	-	expression tag	UNP Q8KHZ8
B	-4	PRO	-	expression tag	UNP Q8KHZ8
B	-3	ARG	-	expression tag	UNP Q8KHZ8
B	-2	GLY	-	expression tag	UNP Q8KHZ8
B	-1	SER	-	expression tag	UNP Q8KHZ8
B	0	HIS	-	expression tag	UNP Q8KHZ8

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0

- Molecule 3 is TRYPTOPHAN (CCD ID: TRP) (formula: C₁₁H₁₂N₂O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 15	C 11	N 2	O 2	0	0
3	A	1	Total 15	C 11	N 2	O 2	0	0

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is ADENOSINE (CCD ID: ADN) (formula: $\text{C}_{10}\text{H}_{13}\text{N}_5\text{O}_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			19	10	5	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	513	Total	O	0	0
			513	513		
6	B	543	Total	O	0	0
			543	543		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan halogenase



• Molecule 1: Tryptophan halogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	114.49Å 114.49Å 231.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.95 – 2.15 19.95 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.95-2.15) 98.7 (19.95-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.152 , 0.194 (Not available) , 0.245	Depositor DCC
R_{free} test set	4630 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9627	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ADN, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	2/4351 (0.0%)	0.91	1/5905 (0.0%)
1	B	0.87	0/4351	0.93	2/5905 (0.0%)
All	All	0.86	2/8702 (0.0%)	0.92	3/11810 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	120	ILE	CA-CB	6.12	1.59	1.54
1	A	235	ILE	CA-CB	5.05	1.60	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330	PHE	CB-CA-C	-5.87	99.59	109.50
1	B	65	ILE	N-CA-C	5.33	112.82	107.55
1	B	293	GLY	N-CA-C	-5.21	103.22	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4234	0	4038	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4234	0	4038	30	0
2	A	1	0	0	1	0
3	A	30	0	18	0	0
4	A	53	0	31	4	0
5	B	19	0	13	4	0
6	A	513	0	0	9	1
6	B	543	0	0	6	0
All	All	9627	0	8138	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:231:ARG:HH12	5:B:2005:ADN:H8	1.16	1.06
1:B:231:ARG:NH1	5:B:2005:ADN:H8	1.80	0.95
1:A:49:GLU:HG3	1:A:173:ALA:HB2	1.53	0.90
1:B:12:GLY:HA2	5:B:2005:ADN:H1'	1.58	0.85
1:B:49:GLU:HG3	1:B:173:ALA:HB2	1.59	0.84
1:A:71:MET:HE3	6:A:2385:HOH:O	1.84	0.76
1:B:231:ARG:NH1	5:B:2005:ADN:C8	2.52	0.72
1:A:13:GLY:O	6:A:2358:HOH:O	2.11	0.68
1:A:437:TYR:HB2	1:A:443:ILE:HD11	1.79	0.65
1:B:390:ARG:HD2	6:B:2274:HOH:O	1.96	0.64
1:A:19:MET:HE1	1:A:365:TYR:HD2	1.63	0.64
1:A:405:ALA:HB2	1:A:430:MET:HE2	1.81	0.61
1:A:12:GLY:HA2	4:A:2004:FAD:H1B	1.83	0.60
1:A:250:LEU:HD13	1:A:353:VAL:HG23	1.84	0.59
1:A:390:ARG:NH1	6:A:2468:HOH:O	2.36	0.59
1:B:437:TYR:HB2	1:B:443:ILE:HD11	1.83	0.58
1:A:71:MET:CE	6:A:2385:HOH:O	2.48	0.58
1:B:198:GLU:OE1	1:B:212:ARG:NH2	2.37	0.58
1:B:528:HIS:C	6:B:2324:HOH:O	2.47	0.57
4:A:2004:FAD:H5'1	6:A:2116:HOH:O	2.05	0.56
1:B:19:MET:HE1	1:B:365:TYR:HD2	1.70	0.55
1:A:243:PHE:CE2	1:A:331:ARG:HG2	2.41	0.55
1:B:255:ALA:HA	1:B:296:TYR:O	2.09	0.53
1:B:378:LYS:NZ	6:B:2066:HOH:O	2.41	0.53
1:A:45:LEU:HD11	1:A:324:PRO:HB2	1.90	0.52
1:A:360:GLY:HA3	4:A:2004:FAD:H1'2	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:489:LEU:HA	1:B:492:MET:HE3	1.92	0.52
1:A:425:ARG:HD3	6:A:2307:HOH:O	2.10	0.51
1:B:129:ARG:HD2	6:B:2299:HOH:O	2.11	0.50
1:A:243:PHE:CZ	1:A:331:ARG:HG2	2.49	0.48
1:B:84:PHE:O	1:B:106:HIS:HA	2.14	0.47
1:B:18:TRP:CZ2	1:B:181:ARG:HG3	2.50	0.47
1:A:71:MET:HG2	6:A:2385:HOH:O	2.14	0.47
4:A:2004:FAD:H51A	6:A:2481:HOH:O	2.15	0.46
1:A:49:GLU:CG	1:A:173:ALA:HB2	2.37	0.46
1:B:45:LEU:HD23	6:B:2514:HOH:O	2.14	0.46
1:B:71:MET:HE2	1:B:71:MET:HB2	1.80	0.46
1:A:255:ALA:HA	1:A:296:TYR:O	2.16	0.46
1:B:146:GLU:O	1:B:150:LEU:HG	2.16	0.46
1:B:250:LEU:HD13	1:B:353:VAL:HG23	1.99	0.45
1:B:521:HIS:NE2	1:B:525:ARG:HD2	2.32	0.44
1:A:438:ARG:HA	1:A:482:PRO:HA	2.00	0.44
1:B:405:ALA:HB2	1:B:430:MET:HE2	2.00	0.43
1:A:355:PRO:HA	2:A:2001:CL:CL	2.56	0.43
1:B:178:ASP:OD2	1:B:182:ARG:NH1	2.52	0.43
1:B:163:LYS:HE3	6:B:2258:HOH:O	2.19	0.42
1:B:241:GLU:HA	1:B:242:PRO:HD3	1.94	0.42
1:B:164:VAL:HG21	1:B:517:LEU:HD11	2.00	0.42
1:B:459:GLU:O	1:B:463:ARG:HG3	2.20	0.42
1:A:116:TYR:OH	1:A:445:ALA:HB1	2.20	0.41
1:B:246:MET:HG3	1:B:333:GLY:HA2	2.02	0.41
1:B:405:ALA:HB2	1:B:430:MET:CE	2.50	0.41
1:B:182:ARG:O	1:B:186:GLU:HB2	2.21	0.41
1:A:53:PRO:HD3	1:A:111:PHE:CE1	2.55	0.41
1:A:528:HIS:C	6:A:2108:HOH:O	2.64	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2207:HOH:O	6:A:2207:HOH:O[4_565]	2.03	0.17

5.3 Torsion angles

5.3.1 Protein backbone

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/463 (96%)	438 (99%)	6 (1%)	59	66
1	B	444/463 (96%)	437 (98%)	7 (2%)	55	61
All	All	888/926 (96%)	875 (98%)	13 (2%)	57	64

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	49	GLU
1	A	115	LYS
1	A	256	VAL
1	A	322	THR
1	A	452	GLN
1	B	2	SER
1	B	198	GLU
1	B	211	VAL
1	B	233	LEU
1	B	237	LYS
1	B	256	VAL
1	B	390	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	ASN
1	A	452	GLN
1	A	494	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	29	GLN
1	B	38	GLN
1	B	252	ASN
1	B	421	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ADN	B	2005	-	21,21,21	1.57	3 (14%)	31,31,31	2.32	12 (38%)
3	TRP	A	2003	-	15,16,16	1.66	3 (20%)	18,22,22	0.71	0
3	TRP	A	2002	-	15,16,16	1.71	3 (20%)	18,22,22	0.66	0
4	FAD	A	2004	-	58,58,58	1.30	7 (12%)	85,89,89	1.73	18 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADN	B	2005	-	-	2/6/22/22	0/3/3/3
3	TRP	A	2003	-	-	2/8/8/8	0/2/2/2
3	TRP	A	2002	-	-	2/8/8/8	0/2/2/2
4	FAD	A	2004	-	-	11/34/50/50	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	2005	ADN	C5-C4	4.90	1.47	1.39
4	A	2004	FAD	C4X-N5	4.67	1.40	1.30
3	A	2002	TRP	CB-CG	4.21	1.58	1.50
3	A	2003	TRP	CB-CG	3.75	1.57	1.50
4	A	2004	FAD	P-O3P	3.23	1.63	1.59
4	A	2004	FAD	C10-N1	3.17	1.39	1.33
4	A	2004	FAD	PA-O3P	3.11	1.62	1.59
5	B	2005	ADN	C5-C6	2.95	1.49	1.41
4	A	2004	FAD	C2A-N1A	2.69	1.38	1.33
4	A	2004	FAD	C2A-N3A	2.69	1.38	1.33
3	A	2002	TRP	CZ3-CE3	2.69	1.43	1.38
5	B	2005	ADN	C8-N7	2.51	1.36	1.31
4	A	2004	FAD	C8A-N7A	2.46	1.36	1.31
3	A	2003	TRP	CH2-CZ3	2.43	1.43	1.38
3	A	2003	TRP	CZ2-CE2	2.12	1.43	1.39
3	A	2002	TRP	CH2-CZ3	2.06	1.42	1.38

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2004	FAD	N3A-C2A-N1A	-5.94	119.59	128.58
5	B	2005	ADN	C5-C4-N3	-5.60	119.00	126.72
4	A	2004	FAD	N9A-C8A-N7A	-5.50	106.13	113.94
5	B	2005	ADN	N3-C4-N9	4.72	135.19	127.17
4	A	2004	FAD	C5A-N7A-C8A	4.62	110.71	103.45
4	A	2004	FAD	C5A-C4A-N3A	-4.41	120.65	126.72
5	B	2005	ADN	O4'-C1'-N9	4.02	115.82	108.09
5	B	2005	ADN	C2'-C1'-N9	-3.89	103.65	113.30
4	A	2004	FAD	C2A-N3A-C4A	3.52	120.42	111.83
5	B	2005	ADN	C2-N3-C4	3.43	120.22	111.83
4	A	2004	FAD	C4A-N9A-C8A	3.32	109.23	105.74
4	A	2004	FAD	C4A-C5A-N7A	-3.24	106.88	110.58
5	B	2005	ADN	N3-C2-N1	-3.07	123.94	128.58
5	B	2005	ADN	C4-C5-N7	-3.05	107.09	110.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2004	FAD	N3A-C4A-N9A	3.02	132.30	127.17
5	B	2005	ADN	C4-N9-C8	2.98	108.86	105.74
5	B	2005	ADN	O3'-C3'-C2'	-2.96	102.34	111.82
4	A	2004	FAD	C10-C4X-N5	-2.79	119.12	124.81
4	A	2004	FAD	C4X-C10-N10	2.65	120.27	116.48
4	A	2004	FAD	O4B-C1B-N9A	2.59	113.06	108.09
4	A	2004	FAD	C4-C4X-N5	2.57	121.75	118.21
5	B	2005	ADN	C2-N1-C6	2.40	122.67	118.73
5	B	2005	ADN	O4'-C1'-C2'	-2.36	101.57	106.62
4	A	2004	FAD	C10-N1-C2	2.33	121.90	116.85
5	B	2005	ADN	C5-N7-C8	2.32	107.09	103.45
4	A	2004	FAD	O3'-C3'-C2'	2.19	113.90	108.93
4	A	2004	FAD	C9A-C5X-N5	-2.14	120.19	122.45
4	A	2004	FAD	C4X-C4-N3	2.11	118.62	113.25
4	A	2004	FAD	C5X-C9A-N10	2.10	119.86	117.97
4	A	2004	FAD	C4X-C10-N1	-2.05	119.57	124.59

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	2002	TRP	O-C-CA-N
4	A	2004	FAD	C2'-C3'-C4'-O4'
4	A	2004	FAD	O3'-C3'-C4'-O4'
4	A	2004	FAD	C5'-O5'-P-O1P
4	A	2004	FAD	C5'-O5'-P-O3P
5	B	2005	ADN	O4'-C4'-C5'-O5'
5	B	2005	ADN	C3'-C4'-C5'-O5'
4	A	2004	FAD	O3'-C3'-C4'-C5'
4	A	2004	FAD	C2'-C3'-C4'-C5'
3	A	2002	TRP	OXT-C-CA-N
3	A	2003	TRP	OXT-C-CA-N
4	A	2004	FAD	N10-C1'-C2'-O2'
4	A	2004	FAD	C5'-O5'-P-O2P
4	A	2004	FAD	O4B-C4B-C5B-O5B
4	A	2004	FAD	C3B-C4B-C5B-O5B
3	A	2003	TRP	O-C-CA-N
4	A	2004	FAD	C4'-C5'-O5'-P


There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	2005	ADN	4	0
4	A	2004	FAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	527/550 (95%)	0.46	30 (5%)	29 33	30, 36, 44, 58	0
1	B	527/550 (95%)	0.37	25 (4%)	36 41	30, 36, 44, 57	0
All	All	1054/1100 (95%)	0.41	55 (5%)	33 36	30, 36, 44, 58	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	336	ARG	5.4
1	A	2	SER	5.2
1	A	42	ILE	4.9
1	A	335	ASN	4.7
1	A	251	LEU	4.7
1	A	334	ARG	3.9
1	B	337	ARG	3.6
1	A	528	HIS	3.5
1	B	2	SER	3.5
1	B	120	ILE	3.4
1	A	199	HIS	3.3
1	A	3	GLY	3.2
1	B	528	HIS	3.0
1	B	42	ILE	2.9
1	A	333	GLY	2.8
1	B	336	ARG	2.7
1	A	331	ARG	2.7
1	B	5	ILE	2.7
1	B	41	ASP	2.7
1	A	252	ASN	2.7
1	A	44	THR	2.6
1	A	330	PHE	2.6
1	A	120	ILE	2.6
1	B	334	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	197	VAL	2.5
1	A	501	GLU	2.5
1	B	45	LEU	2.5
1	B	453	LEU	2.5
1	B	333	GLY	2.5
1	B	472	TYR	2.5
1	B	6	ASP	2.5
1	A	41	ASP	2.4
1	B	310	ARG	2.4
1	A	198	GLU	2.4
1	A	473	CYS	2.4
1	B	43	PRO	2.4
1	A	527	GLN	2.4
1	A	250	LEU	2.3
1	B	117	HIS	2.3
1	B	122	LEU	2.3
1	A	195	ASP	2.3
1	B	335	ASN	2.3
1	A	45	LEU	2.3
1	B	471	TYR	2.3
1	B	497	GLU	2.2
1	A	328	ILE	2.2
1	A	332	VAL	2.2
1	A	254	SER	2.1
1	B	71	MET	2.1
1	B	265	ASP	2.1
1	A	43	PRO	2.1
1	A	46	GLY	2.1
1	B	140	ASP	2.0
1	B	11	VAL	2.0
1	A	102	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ADN	B	2005	19/19	0.70	0.17	48,50,54,56	19
3	TRP	A	2002	15/15	0.84	0.14	37,40,43,43	0
4	FAD	A	2004	53/53	0.86	0.15	30,45,55,56	53
3	TRP	A	2003	15/15	0.90	0.10	29,31,32,34	0
2	CL	A	2001	1/1	0.96	0.29	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around FAD A 2004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.