



Full wwPDB EM Validation Report ⓘ

Mar 26, 2026 – 11:26 PM UTC

PDB ID : 9OA6 / pdb_00009oa6
EMDB ID : EMD-70274
Title : Cryo-EM structure of the Zdhhc5-GOLGA7 complex
Authors : Kahlson, M.A.; Dixon, S.J.; Butterwick, J.A.; Wang, H.
Deposited on : 2025-04-19
Resolution : 3.90 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8855 (3.40 - 4.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	153	
2	A	718	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2909 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Golgin subfamily A member 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	B	120	Total	C	N	O	S	0	0
			971	617	164	185	5		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	MET	-	initiating methionine	UNP Q7Z5G4
B	-11	ASP	-	expression tag	UNP Q7Z5G4
B	-10	TYR	-	expression tag	UNP Q7Z5G4
B	-9	LYS	-	expression tag	UNP Q7Z5G4
B	-8	ASP	-	expression tag	UNP Q7Z5G4
B	-7	ASP	-	expression tag	UNP Q7Z5G4
B	-6	ASP	-	expression tag	UNP Q7Z5G4
B	-5	ASP	-	expression tag	UNP Q7Z5G4
B	-4	LYS	-	expression tag	UNP Q7Z5G4
B	-3	SER	-	expression tag	UNP Q7Z5G4
B	-2	GLY	-	expression tag	UNP Q7Z5G4
B	-1	ARG	-	expression tag	UNP Q7Z5G4
B	0	ALA	-	expression tag	UNP Q7Z5G4
B	138	ALA	-	expression tag	UNP Q7Z5G4
B	139	ALA	-	expression tag	UNP Q7Z5G4
B	140	ALA	-	expression tag	UNP Q7Z5G4

- Molecule 2 is a protein called Palmitoyltransferase ZDHHC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	A	249	Total	C	N	O	S	0	0
			1936	1252	334	330	20		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	716	ALA	-	expression tag	UNP Q8VDZ4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	717	ALA	-	expression tag	UNP Q8VDZ4
A	718	ALA	-	expression tag	UNP Q8VDZ4

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
3	A	2	Total	Zn	0
			2	2	

LEU	ASP	THR	THR
THR	ASN	PRO	THR
PRO	LEU	SER	ASP
SER	ARG	SER	PRO
ASP	HIS	THR	ASP
PRO	ILE	PRO	ASP
ASP	VAL	ASP	PHE
GLU	ALA	GLU	GLU
SER	LEU	SER	VAL
VAL	GLN	VAL	GLN
GLN	GLU	GLU	ALA
ARG	ARG	GLY	GLY
PRO	LYS	PRO	GLY
LEU	LEU	GLU	LEU
PRO	LEU	PRO	ASP
ASP	ARG	ASP	PRO
PRO	GLN	PRO	PRO
SER	SER	LEU	GLY
PRO	PRO	GLY	GLY
GLY	PRO	ARG	TYR
LEU	LEU	GLY	ALA
GLY	ARG	PRO	SER
PRO	GLU	GLU	THR
GLU	GLU	SER	ALA
ALA	PRO	ALA	ARG
ARG	GLY	LEU	ARG
LEU	LEU	GLY	GLY
GLY	LEU	GLY	ALA
ALA	GLN	GLU	GLU
GLU	THR	ARG	HIS
HIS	PRO	GLY	PRO
PRO	SER	ARG	ARG
LEU	GLY	LEU	LEU
LEU	HIS	LEU	PRO
PRO	ALA	THR	GLY
THR	ARG	GLY	PRO
PRO	THR	SER	HIS
ASP	ASP	PRO	GLU
PRO	ASP	PRO	THR
PRO	SER	PRO	PRO
VAL	ARG	VAL	VAL
ARG	SER	ARG	ARG
SER	SER	SER	SER
TYR	TYR	VAL	TYR
ILE	ILE	ALA	ILE

LEU	PRO
GLN	LEU
PRO	SER
LEU	LYS
THR	THR
PRO	PRO
LYS	LEU
ASP	GLY
GLU	ARG
VAL	PRO
VAL	ALA
LEU	GLN
LEU	PRO
LYS	ARG
THR	PHE
THR	GLY
TYR	LYS
SER	PRO
LYS	ASP
SER	GLY
ASN	LEU
GLY	ARG
GLN	SER
PRO	ARG
LYS	GLY
SER	LEU
ILE	GLY
GLY	SER
SER	PRO
ALA	GLU
SER	PRO
PRO	GLY
GLY	THR
PRO	THR
GLY	ALA
GLN	PRO
PRO	GLN
LEU	TYR
GLY	LEU
LEU	GLY
SER	ARG
SER	SER
PRO	ILE
THR	SER
ARG	TYR
GLY	ARG
GLY	THR
VAL	SER
LYS	LYS
VAL	VAL
SER	SER
GLY	GLY
VAL	VAL
GLY	GLY
GLY	GLY
THR	THR
THR	THR
TYR	TYR
GLU	GLU
VAL	VAL
ILE	ALA

SER
VAL
ALA
ALA
ALA

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	93643	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	80	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.184	Depositor
Minimum map value	-0.631	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.011	Depositor
Recommended contour level	0.17	Depositor
Map size ($\text{\AA}$)	319.488, 319.488, 319.488	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.832, 0.832, 0.832	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.15	0/987	0.43	0/1330
2	A	0.17	0/1989	0.51	0/2702
All	All	0.16	0/2976	0.48	0/4032

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	971	0	974	27	0
2	A	1936	0	1897	52	0
3	A	2	0	0	0	0
All	All	2909	0	2871	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:68:MET:HE3	2:A:69:ASP:H	1.52	0.74
2:A:90:LEU:HD22	2:A:105:TRP:HD1	1.52	0.72
1:B:57:ALA:HB2	1:B:89:VAL:HG11	1.70	0.71
2:A:68:MET:HE2	2:A:144:ARG:HH21	1.62	0.64
2:A:215:ARG:HH12	2:A:220:GLN:HG2	1.61	0.63
2:A:134:CYS:HB3	2:A:137:VAL:HG22	1.82	0.61
2:A:90:LEU:HB3	2:A:105:TRP:HB3	1.84	0.59
1:B:76:TYR:HB2	2:A:64:MET:HE1	1.83	0.59
2:A:72:ILE:HD13	2:A:127:VAL:HG12	1.84	0.59
1:B:101:ASN:HA	1:B:105:TYR:HD1	1.68	0.57
2:A:40:SER:HB3	2:A:45:PRO:HA	1.87	0.56
2:A:215:ARG:NH2	2:A:220:GLN:HA	2.20	0.56
2:A:90:LEU:HD22	2:A:105:TRP:CD1	2.37	0.56
1:B:56:GLU:HA	1:B:59:LYS:HE2	1.88	0.56
2:A:130:PHE:HE1	2:A:140:CYS:HB2	1.71	0.56
2:A:143:ARG:HH12	2:A:249:ALA:HB2	1.72	0.55
1:B:44:GLU:HG3	1:B:48:ARG:HE	1.71	0.55
2:A:71:GLY:HA3	2:A:128:GLU:HG2	1.89	0.54
2:A:63:SER:HB2	2:A:67:PHE:HE1	1.73	0.53
1:B:18:TYR:HB3	2:A:96:ILE:HG22	1.88	0.53
2:A:105:TRP:HZ2	2:A:110:ARG:NH1	2.07	0.53
1:B:100:GLN:HB3	1:B:105:TYR:HE1	1.73	0.52
2:A:116:ARG:O	2:A:129:GLU:HA	2.09	0.52
1:B:90:LEU:HD11	2:A:250:PRO:HG2	1.91	0.52
1:B:34:LEU:HB3	1:B:38:ILE:HG22	1.92	0.51
2:A:234:ASN:HB3	2:A:239:ASN:OD1	2.11	0.51
2:A:68:MET:HE2	2:A:144:ARG:NH2	2.24	0.50
2:A:47:VAL:HB	2:A:48:PRO:HD3	1.92	0.50
2:A:82:LYS:HA	2:A:82:LYS:HE2	1.94	0.50
2:A:104:LYS:HB3	2:A:113:ARG:NE	2.26	0.50
2:A:238:ASN:O	2:A:242:ARG:HG2	2.11	0.50
1:B:84:THR:HG23	1:B:87:GLU:H	1.78	0.49
2:A:43:VAL:HG22	2:A:44:SER:H	1.77	0.49
1:B:29:LYS:HD3	1:B:30:PHE:N	2.27	0.48
2:A:36:CYS:N	2:A:37:PRO:HD3	2.27	0.48
1:B:29:LYS:HA	1:B:29:LYS:HE2	1.95	0.48
2:A:177:GLU:HG3	2:A:179:SER:HB3	1.95	0.48
1:B:74:THR:HG22	1:B:74:THR:O	2.14	0.48
2:A:253:LEU:HD23	2:A:254:GLY:N	2.29	0.47
1:B:16:ARG:HH21	1:B:24:CYS:HA	1.79	0.47
1:B:16:ARG:HA	1:B:26:PHE:HA	1.96	0.47
1:B:95:LYS:HA	1:B:95:LYS:HD2	1.76	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:120:CYS:HB3	2:A:123:CYS:HB3	1.98	0.46
1:B:14:ILE:HB	1:B:123:ILE:HG22	1.97	0.46
2:A:62:PHE:O	2:A:66:THR:HG23	2.16	0.46
1:B:85:HIS:O	1:B:89:VAL:HG23	2.17	0.45
1:B:23:ARG:HH12	2:A:242:ARG:HH12	1.63	0.45
2:A:215:ARG:CZ	2:A:220:GLN:HA	2.46	0.45
2:A:120:CYS:SG	2:A:123:CYS:HB3	2.55	0.45
2:A:34:PHE:HB3	2:A:182:ARG:NE	2.31	0.44
2:A:36:CYS:HB2	2:A:39:LEU:HD22	1.99	0.44
2:A:215:ARG:NH1	2:A:220:GLN:HG2	2.32	0.44
2:A:91:TYR:C	2:A:92:LYS:HD3	2.43	0.44
1:B:94:SER:HA	1:B:97:ILE:HG12	2.00	0.44
2:A:43:VAL:HG13	2:A:44:SER:N	2.33	0.43
1:B:99:GLU:OE2	1:B:100:GLN:HG2	2.18	0.43
1:B:54:TYR:HE1	1:B:93:VAL:HG11	1.84	0.43
2:A:62:PHE:HB2	2:A:152:PHE:CD1	2.54	0.43
1:B:90:LEU:HA	1:B:93:VAL:HG12	2.00	0.43
2:A:31:PHE:HA	2:A:35:THR:OG1	2.18	0.42
2:A:147:ARG:HH22	2:A:246:SER:HB3	1.84	0.42
2:A:251:ARG:HG2	2:A:253:LEU:O	2.19	0.42
1:B:118:ARG:HH22	2:A:80:GLU:CD	2.28	0.42
2:A:50:TYR:CE2	2:A:166:PHE:HE2	2.38	0.41
2:A:127:VAL:HG23	2:A:128:GLU:O	2.20	0.41
2:A:70:PRO:HB3	2:A:145:ASN:ND2	2.35	0.41
2:A:153:LEU:HD23	2:A:153:LEU:HA	1.94	0.41
1:B:94:SER:O	1:B:98:GLN:HG2	2.20	0.41
2:A:240:VAL:HA	2:A:243:VAL:HG22	2.01	0.41
2:A:95:GLU:HA	2:A:100:GLN:HA	2.02	0.41
2:A:137:VAL:HG23	2:A:139:ASN:H	1.85	0.41
2:A:237:CYS:HA	2:A:240:VAL:HG22	2.03	0.41
1:B:53:LEU:HD23	1:B:53:LEU:HA	1.93	0.40
1:B:64:SER:HA	1:B:67:GLU:HG2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	118/153 (77%)	116 (98%)	2 (2%)	0	100	100
2	A	247/718 (34%)	201 (81%)	42 (17%)	4 (2%)	7	36
All	All	365/871 (42%)	317 (87%)	44 (12%)	4 (1%)	14	43

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	43	VAL
2	A	121	SER
2	A	181	VAL
2	A	217	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	106/132 (80%)	106 (100%)	0	100	100
2	A	206/614 (34%)	206 (100%)	0	100	100
All	All	312/746 (42%)	312 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	15	GLN
1	B	27	GLN
2	A	218	ASN
2	A	238	ASN
2	A	239	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-70274. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 212




Y Index: 196




Z Index: 177


6.3.2 Raw map



X Index: 212



Y Index: 197




Z Index: 177


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.17. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 15 nm^3 ; this corresponds to an approximate mass of 14 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.256 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.90	-	-
Author-provided FSC curve	3.85	5.18	3.94
Unmasked-calculated*	5.30	6.91	5.50

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 5.30 differs from the reported value 3.9 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-70274 and PDB model 9OA6. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.17 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.17).

9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 52% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.17) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5200	0.3140
A	0.5080	0.3090
B	0.5430	0.3220

