



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 01:15 PM UTC

PDB ID : 4OBP / pdb_00004obp
Title : MAP4K4 in complex with inhibitor (compound 29), 6-(2-FLUOROPYRIDIN-4-YL)PYRIDO[3,2-D]PYRIMIDIN-4-AMINE
Authors : Harris, S.F.; Wu, P.
Deposited on : 2014-01-07
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	
1	B	332	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase kinase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2300	1466	402	419	13			
1	B	290	Total	C	N	O	S	0	1	0
			2364	1504	420	427	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP O95819
A	1	SER	-	expression tag	UNP O95819
A	329	GLY	-	expression tag	UNP O95819
A	330	ASN	-	expression tag	UNP O95819
A	331	SER	-	expression tag	UNP O95819
B	0	GLY	-	expression tag	UNP O95819
B	1	SER	-	expression tag	UNP O95819
B	329	GLY	-	expression tag	UNP O95819
B	330	ASN	-	expression tag	UNP O95819
B	331	SER	-	expression tag	UNP O95819

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is 6-(2-fluoropyridin-4-yl)pyrido[3,2-d]pyrimidin-4-amine (CCD ID: 2QU) (formula: $C_{12}H_8FN_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	0	0
			18	12	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	169	Total 169	O 169	0	0
5	B	152	Total 152	O 152	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 4



- Molecule 1: Mitogen-activated protein kinase kinase kinase kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.30Å 88.69Å 91.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.27 – 2.27 60.27 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.0 (60.27-2.27) 99.1 (60.27-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.27Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.11.2	Depositor
R, R_{free}	0.203 , 0.247 0.207 , 0.255	Depositor DCC
R_{free} test set	1546 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtriage
Anisotropy	0.420	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5016	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MES, 2QU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/2352	1.33	11/3178 (0.3%)
1	B	0.81	1/2420 (0.0%)	1.31	7/3270 (0.2%)
All	All	0.81	1/4772 (0.0%)	1.32	18/6448 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	72	MET	SD-CE	-6.05	1.64	1.79

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	THR	CB-CA-C	7.00	121.09	109.89
1	B	193	TYR	N-CA-C	6.47	119.15	111.71
1	A	193	TYR	N-CA-C	5.96	118.57	111.71
1	A	41	LYS	N-CA-C	-5.76	100.44	109.25
1	B	105	MET	CA-C-N	5.43	128.56	120.90
1	B	105	MET	C-N-CA	5.43	128.56	120.90
1	A	203	ASP	N-CA-C	5.38	117.84	111.33
1	B	174	VAL	CA-C-N	5.30	127.64	120.38
1	B	174	VAL	C-N-CA	5.30	127.64	120.38
1	B	104	VAL	N-CA-C	5.29	115.58	108.17
1	A	276	GLN	N-CA-CB	5.25	118.86	110.73
1	A	65	GLU	CA-C-N	5.22	127.14	120.56
1	A	65	GLU	C-N-CA	5.22	127.14	120.56
1	A	244	LEU	N-CA-C	5.20	117.03	111.36
1	A	186	ASN	CA-CB-CG	5.15	117.75	112.60
1	B	244	LEU	N-CA-C	5.13	116.95	111.36
1	A	211	ASP	CA-CB-CG	5.08	117.69	112.60
1	A	191	THR	CB-CA-C	5.06	117.36	109.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	0	2296	19	0
1	B	2364	0	2363	23	0
2	A	1	0	0	0	0
3	A	12	0	13	0	0
4	A	18	0	8	2	0
5	A	169	0	0	4	0
5	B	152	0	0	3	0
All	All	5016	0	4680	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:GLU:OE1	1:B:181:THR:HG21	1.68	0.93
1:B:18:LEU:HD11	1:B:66:ILE:HD11	1.61	0.81
1:B:44:HIS:ND1	1:B:47:THR:HG22	2.08	0.69
1:A:189:ILE:HG13	1:B:238:PRO:HB2	1.76	0.67
1:A:70:ILE:HD11	1:A:101:LEU:HD21	1.82	0.60
1:B:73:LEU:HD23	1:B:175:SER:HB3	1.84	0.60
1:A:73:LEU:HD23	1:A:103:LEU:HD11	1.84	0.59
1:A:276:GLN:HG2	5:A:576:HOH:O	2.04	0.57
5:A:635:HOH:O	1:B:240:ARG:HD2	2.04	0.56
1:A:73:LEU:HG	1:A:85:TYR:HB2	1.87	0.55
1:B:47:THR:HG23	1:B:49:GLN:H	1.72	0.55
1:A:36:TYR:CD1	4:A:403:2QU:H6	2.43	0.54
1:A:243:PHE:CZ	1:B:206:PRO:HB3	2.44	0.53
1:A:36:TYR:CG	4:A:403:2QU:H2	2.44	0.52
1:A:168:LYS:NZ	5:A:632:HOH:O	2.42	0.52
1:B:213:ARG:NH1	5:B:523:HOH:O	2.41	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:ARG:HH21	1:A:297:GLN:HG3	1.79	0.48
1:A:251:PRO:HD2	1:A:270:LEU:HD13	1.96	0.47
1:A:15:LEU:HG	1:A:90:ILE:HD11	1.95	0.47
1:B:62:GLU:O	1:B:66:ILE:HG23	2.17	0.45
1:B:63:GLU:HG2	1:B:90:ILE:HD13	1.99	0.45
1:B:18:LEU:HG	1:B:90:ILE:HD12	1.99	0.44
1:B:69:GLU:HA	1:B:72:MET:HE3	2.00	0.43
1:B:181:THR:HG23	1:B:182:VAL:HG13	2.00	0.43
1:A:189:ILE:HD13	1:B:242:LEU:HD22	1.98	0.43
1:A:255:SER:OG	1:A:257:LYS:HG2	2.18	0.43
1:A:242:LEU:HD22	1:B:189:ILE:HD13	1.99	0.43
1:A:38:GLN:HG3	1:A:40:TYR:HE1	1.83	0.42
1:A:189:ILE:CG1	1:B:238:PRO:HB2	2.48	0.42
1:A:240:ARG:NH2	5:A:647:HOH:O	2.52	0.42
1:B:18:LEU:HD21	1:B:66:ILE:HG13	2.02	0.42
1:B:168:LYS:NZ	5:B:522:HOH:O	2.53	0.42
1:A:38:GLN:HB3	1:A:55:VAL:O	2.20	0.42
1:B:247:ARG:NH2	5:B:551:HOH:O	2.53	0.42
1:B:251:PRO:HD2	1:B:270:LEU:HD13	2.01	0.42
1:B:255:SER:OG	1:B:257:LYS:HG2	2.19	0.42
1:B:43:ARG:HH11	1:B:48:GLY:HA2	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/332 (84%)	267 (96%)	12 (4%)	0	100	100
1	B	287/332 (86%)	279 (97%)	8 (3%)	0	100	100
All	All	566/664 (85%)	546 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/290 (87%)	236 (94%)	15 (6%)	17	23
1	B	258/290 (89%)	243 (94%)	15 (6%)	18	24
All	All	509/580 (88%)	479 (94%)	30 (6%)	18	23

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ILE
1	A	56	MET
1	A	60	GLU
1	A	66	ILE
1	A	68	LEU
1	A	72	MET
1	A	80	ARG
1	A	163	GLU
1	A	186	ASN
1	A	209	THR
1	A	240	ARG
1	A	276	GLN
1	A	295	GLU
1	A	304	ASP
1	A	308	ARG
1	B	18	LEU
1	B	61	ASP
1	B	66	ILE
1	B	75	LYS
1	B	80	ARG
1	B	93	SER
1	B	121	LYS
1	B	126	LYS
1	B	163	GLU
1	B	164	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	178	LEU
1	B	186	ASN
1	B	210	TYR
1	B	295	GLU
1	B	308	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	71	ASN
1	A	164	ASN
1	A	294	ASN
1	B	38	GLN
1	B	97	HIS
1	B	297	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MES	A	402	-	12,12,12	0.77	0	15,16,16	0.33	0
4	2QU	A	403	-	20,20,20	0.65	1 (5%)	26,28,28	1.34	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	A	402	-	-	0/6/14/14	0/1/1/1
4	2QU	A	403	-	-	0/4/4/4	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	403	2QU	C16-N15	2.35	1.37	1.31

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	2QU	C10-C5-N4	4.62	123.89	120.33
4	A	403	2QU	C3-N4-C5	3.92	121.20	118.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	2QU	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/332 (85%)	0.29	14 (4%) 35 36	23, 42, 87, 125	0
1	B	290/332 (87%)	0.56	21 (7%) 21 22	18, 49, 90, 110	1 (0%)
All	All	575/664 (86%)	0.43	35 (6%) 27 28	18, 46, 90, 125	1 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	LEU	3.4
1	A	13	ILE	3.3
1	A	176	ALA	3.0
1	B	210	TYR	2.8
1	B	15	LEU	2.8
1	A	188	PHE	2.8
1	B	24	ILE	2.6
1	B	23	GLY	2.6
1	B	66	ILE	2.6
1	A	58	VAL	2.6
1	B	58	VAL	2.6
1	A	66	ILE	2.6
1	B	212[A]	TYR	2.6
1	A	173	GLY	2.5
1	B	184	ARG	2.4
1	B	181	THR	2.4
1	A	33	ASN	2.4
1	B	45	VAL	2.4
1	B	38	GLN	2.4
1	B	17	SER	2.4
1	B	178	LEU	2.3
1	A	35	THR	2.3
1	A	307	ASP	2.2
1	B	180	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	21	PRO	2.2
1	B	94	PRO	2.2
1	A	209	THR	2.2
1	A	186	ASN	2.2
1	B	49	GLN	2.2
1	B	18	LEU	2.1
1	A	101	LEU	2.1
1	B	95	PRO	2.1
1	B	39	VAL	2.1
1	A	64	GLU	2.0
1	B	91	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MES	A	402	12/12	0.73	0.19	65,67,74,76	0
2	MG	A	401	1/1	0.94	0.06	53,53,53,53	0
4	2QU	A	403	18/18	0.96	0.06	30,36,38,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.