



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:52 AM UTC

PDB ID : 4OBR / pdb_00004obr
Title : Crystal structure of human alpha-L-iduronidase complex with alpha-L-iduronic acid
Authors : Bie, H.Y.; Yin, J.; He, X.; Kermode, A.R.; James, M.N.G.
Deposited on : 2014-01-07
Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	4% 90% 5%
1	B	627	3% 92% 5%
2	C	2	50% 50%
2	E	2	50% 50%
3	D	5	20% 80%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	9	 A horizontal bar chart showing the quality of chain F. The bar is divided into two segments: a green segment on the left representing 22% and a yellow segment on the right representing 78%. The percentages are labeled below the corresponding segments.

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 10073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-L-iduronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	595	Total	C	N	O	S	0	2	0
			4746	3036	854	842	14			
1	B	606	Total	C	N	O	S	0	5	0
			4854	3099	881	860	14			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLN	HIS	SEE REMARK 999	UNP P35475
A	63	PRO	GLN	SEE REMARK 999	UNP P35475
A	105	GLN	ARG	SEE REMARK 999	UNP P35475
B	33	GLN	HIS	SEE REMARK 999	UNP P35475
B	63	PRO	GLN	SEE REMARK 999	UNP P35475
B	105	GLN	ARG	SEE REMARK 999	UNP P35475

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	9	Total	C	N	O	0	0	0
			105	58	2	45			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			10	4	6		
7	B	1	Total	C	O	0	0
			10	4	6		
7	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 8 is alpha-L-idopyranuronic acid (CCD ID: IDR) (formula: $C_6H_{10}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			13	6	7		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Cl	0	0
			1	1		
9	B	1	Total	Cl	0	0
			1	1		

- Molecule 10 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	37	Total	O	0	0
			37	37		
11	B	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-L-iduronidase



• Molecule 1: Alpha-L-iduronidase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose







- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  20% 80%



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  22% 78%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	258.94Å 258.94Å 71.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.94 – 2.46 48.94 – 2.46	Depositor EDS
% Data completeness (in resolution range)	97.8 (48.94-2.46) 97.8 (48.94-2.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.202 , 0.239 (Not available) , 0.224	Depositor DCC
R_{free} test set	3175 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.697	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 20.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10073	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MAN, IDR, TLA, SO4, NAG, GOL, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/4891	0.73	0/6682
1	B	0.53	0/5002	0.75	2/6832 (0.0%)
All	All	0.52	0/9893	0.74	2/13514 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	62	ASP	CA-C-N	5.13	126.25	119.84
1	B	62	ASP	C-N-CA	5.13	126.25	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4746	0	4640	13	0
1	B	4854	0	4754	16	0
2	C	28	0	25	0	0
2	E	28	0	25	0	0
3	D	61	0	52	0	0
4	F	105	0	88	0	0
5	A	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	14	0	13	0	0
6	A	18	0	24	0	0
6	B	36	0	48	6	0
7	A	10	0	4	1	0
7	B	20	0	8	0	0
8	A	13	0	9	0	0
8	B	13	0	9	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	5	0	0	0	0
11	A	37	0	0	0	0
11	B	69	0	0	0	0
All	All	10073	0	9712	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:PRO:O	1:A:531:ARG:NH1	2.33	0.61
1:A:127:LEU:HD22	1:A:171:HIS:CE1	2.43	0.54
1:B:501:PHE:HA	1:B:504:MET:HE3	1.91	0.52
1:B:75:ALA:HB2	6:B:917:GOL:H32	1.90	0.52
1:A:276:GLU:OE1	1:A:329:HIS:NE2	2.40	0.51
1:A:127:LEU:HD23	1:A:127:LEU:N	2.26	0.50
1:B:356:HIS:HB2	6:B:913:GOL:C3	2.43	0.49
1:A:401:LEU:HD21	1:A:424:ALA:HB2	1.95	0.48
1:B:530:LEU:HB3	1:B:534:SER:OG	2.13	0.48
1:B:314:ALA:HA	1:B:360:PHE:O	2.14	0.47
1:B:604:LEU:C	1:B:604:LEU:HD23	2.40	0.47
1:B:471:TYR:HA	1:B:539:HIS:O	2.15	0.47
1:B:555:ARG:NH2	6:B:917:GOL:O2	2.41	0.46
1:A:273:LEU:HD11	1:A:333:LEU:HD13	1.97	0.46
1:A:315:ASP:HB2	1:A:478:ASN:OD1	2.16	0.46
1:B:434:TRP:CH2	1:B:436:ALA:HB2	2.51	0.45
1:A:588:ASP:OD1	1:A:588:ASP:N	2.50	0.45
1:A:245:THR:HB	7:A:912:TLA:O4	2.17	0.45
1:B:127:LEU:CD2	1:B:171:HIS:CE1	3.01	0.44
1:A:383:ARG:HG2	1:A:387:LEU:HD23	2.01	0.43
1:B:356:HIS:HB2	6:B:913:GOL:H32	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:570:ASP:CG	1:B:603:ASN:HD21	2.28	0.42
1:B:75:ALA:HB2	6:B:917:GOL:C3	2.50	0.42
1:B:346:LEU:C	1:B:346:LEU:HD23	2.45	0.41
1:A:434:TRP:CH2	1:A:436:ALA:HB2	2.56	0.41
1:B:224:SER:OG	6:B:918:GOL:H32	2.21	0.41
1:A:584:GLN:OE1	1:A:619:ARG:NH2	2.47	0.41
1:B:188:PHE:CE1	1:B:193:MET:HG2	2.55	0.40
1:A:501:PHE:HA	1:A:504:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	589/627 (94%)	564 (96%)	25 (4%)	0	100	100
1	B	605/627 (96%)	584 (96%)	20 (3%)	1 (0%)	43	54
All	All	1194/1254 (95%)	1148 (96%)	45 (4%)	1 (0%)	48	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	247	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/527 (96%)	496 (98%)	9 (2%)	51	65
1	B	516/527 (98%)	508 (98%)	8 (2%)	55	69
All	All	1021/1054 (97%)	1004 (98%)	17 (2%)	53	67

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	LEU
1	A	245	THR
1	A	322	VAL
1	A	336	ASN
1	A	375	ARG
1	A	553	ARG
1	A	588	ASP
1	A	598	LYS
1	A	639	LEU
1	B	64	TYR
1	B	195	MET
1	B	227	THR
1	B	337	THR
1	B	379	VAL
1	B	417	THR
1	B	590	LYS
1	B	623	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	171	HIS
1	A	196	GLN
1	A	200	ASN
1	A	358	HIS
1	A	416	HIS
1	B	73	ASN
1	B	171	HIS
1	B	196	GLN
1	B	200	ASN
1	B	262	HIS
1	B	281	GLN
1	B	330	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	603	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	C	1	2,1	14,14,15	0.60	0	17,19,21	1.06	1 (5%)
2	NAG	C	2	2	14,14,15	0.57	0	17,19,21	0.80	0
3	NAG	D	1	1,3	14,14,15	0.62	0	17,19,21	1.05	1 (5%)
3	NAG	D	2	3	14,14,15	0.61	0	17,19,21	1.19	2 (11%)
3	BMA	D	3	3	11,11,12	0.38	0	15,15,17	0.46	0
3	MAN	D	4	3	11,11,12	0.44	0	15,15,17	0.86	1 (6%)
3	MAN	D	5	3	11,11,12	0.52	0	15,15,17	0.99	1 (6%)
2	NAG	E	1	2,1	14,14,15	0.52	0	17,19,21	1.54	3 (17%)
2	NAG	E	2	2	14,14,15	0.59	0	17,19,21	0.90	0
4	NAG	F	1	4,1	14,14,15	0.70	0	17,19,21	1.09	1 (5%)
4	NAG	F	2	4	14,14,15	0.76	1 (7%)	17,19,21	0.88	0
4	BMA	F	3	4	11,11,12	0.56	0	15,15,17	0.93	0
4	MAN	F	4	4	11,11,12	0.66	0	15,15,17	1.12	1 (6%)
4	MAN	F	5	4	11,11,12	0.45	0	15,15,17	1.11	1 (6%)
4	MAN	F	6	4	11,11,12	0.53	0	15,15,17	0.74	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MAN	F	7	4	11,11,12	0.48	0	15,15,17	1.04	1 (6%)
4	MAN	F	8	4	11,11,12	0.47	0	15,15,17	1.06	1 (6%)
4	MAN	F	9	4	11,11,12	0.68	0	15,15,17	1.06	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1
3	MAN	D	5	3	-	1/2/19/22	0/1/1/1
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
4	NAG	F	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
4	MAN	F	4	4	-	2/2/19/22	0/1/1/1
4	MAN	F	5	4	-	0/2/19/22	0/1/1/1
4	MAN	F	6	4	-	0/2/19/22	0/1/1/1
4	MAN	F	7	4	-	2/2/19/22	0/1/1/1
4	MAN	F	8	4	-	2/2/19/22	0/1/1/1
4	MAN	F	9	4	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	2	NAG	O5-C1	-2.01	1.40	1.43

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	NAG	O5-C1-C2	-3.88	105.29	111.29
2	E	1	NAG	C1-O5-C5	3.73	117.19	112.19
4	F	7	MAN	C1-O5-C5	3.52	116.90	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-O5-C5	2.74	115.86	112.19
4	F	4	MAN	C1-O5-C5	2.72	115.83	112.19
3	D	5	MAN	C1-O5-C5	2.71	115.81	112.19
2	C	1	NAG	O5-C1-C2	-2.60	107.27	111.29
4	F	8	MAN	C1-C2-C3	-2.56	105.92	109.64
4	F	9	MAN	C1-C2-C3	2.56	113.37	109.64
3	D	2	NAG	C1-O5-C5	2.53	115.58	112.19
4	F	5	MAN	O5-C1-C2	2.48	116.70	110.79
4	F	1	NAG	O5-C1-C2	-2.45	107.50	111.29
3	D	2	NAG	O4-C4-C3	-2.36	104.81	110.38
3	D	4	MAN	O5-C1-C2	2.20	116.04	110.79
2	E	1	NAG	O7-C7-C8	-2.13	118.25	122.05

There are no chirality outliers.


All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	F	8	MAN	O5-C5-C6-O6
4	F	7	MAN	C4-C5-C6-O6
4	F	8	MAN	C4-C5-C6-O6
4	F	9	MAN	O5-C5-C6-O6
4	F	7	MAN	O5-C5-C6-O6
3	D	5	MAN	O5-C5-C6-O6
4	F	4	MAN	O5-C5-C6-O6
4	F	4	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
3	D	1	NAG	C4-C5-C6-O6
4	F	9	MAN	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	901	1	14,14,15	0.46	0	17,19,21	0.77	0
6	GOL	B	914	-	5,5,5	0.27	0	5,5,5	0.32	0
6	GOL	B	915	-	5,5,5	0.38	0	5,5,5	0.24	0
6	GOL	B	916	-	5,5,5	0.43	0	5,5,5	0.42	0
8	IDR	B	921	-	13,13,13	0.98	0	18,19,19	2.18	6 (33%)
7	TLA	B	920	-	9,9,9	1.09	0	12,12,12	1.18	1 (8%)
7	TLA	B	919	-	9,9,9	1.30	0	12,12,12	1.02	1 (8%)
6	GOL	A	909	-	5,5,5	0.30	0	5,5,5	0.32	0
5	NAG	A	901	1	14,14,15	0.57	0	17,19,21	1.37	1 (5%)
6	GOL	B	918	-	5,5,5	0.30	0	5,5,5	0.45	0
8	IDR	A	913	-	13,13,13	0.87	0	18,19,19	2.06	4 (22%)
10	SO4	A	915	-	4,4,4	0.46	0	6,6,6	0.05	0
6	GOL	A	911	-	5,5,5	0.29	0	5,5,5	0.58	0
6	GOL	A	910	-	5,5,5	0.36	0	5,5,5	0.14	0
7	TLA	A	912	-	9,9,9	1.15	0	12,12,12	1.23	2 (16%)
6	GOL	B	913	-	5,5,5	0.35	0	5,5,5	0.37	0
6	GOL	B	917	-	5,5,5	0.34	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	901	1	-	0/6/23/26	0/1/1/1
6	GOL	B	914	-	-	0/4/4/4	-
6	GOL	B	915	-	-	2/4/4/4	-
6	GOL	B	916	-	-	2/4/4/4	-
8	IDR	B	921	-	-	0/4/24/24	0/1/1/1
7	TLA	B	920	-	-	6/12/12/12	-
7	TLA	B	919	-	-	2/12/12/12	-
6	GOL	A	909	-	-	0/4/4/4	-
5	NAG	A	901	1	-	0/6/23/26	0/1/1/1
6	GOL	B	918	-	-	2/4/4/4	-
8	IDR	A	913	-	-	0/4/24/24	0/1/1/1
6	GOL	A	911	-	-	1/4/4/4	-
6	GOL	A	910	-	-	4/4/4/4	-
7	TLA	A	912	-	-	0/12/12/12	-
6	GOL	B	913	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	917	-	-	4/4/4/4	-

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	921	IDR	O5-C5-C6	5.91	124.62	106.14
8	A	913	IDR	O5-C5-C6	5.78	124.22	106.14
5	A	901	NAG	C1-O5-C5	3.72	117.17	112.19
8	A	913	IDR	C1-C2-C3	3.32	117.13	110.36
8	B	921	IDR	C1-C2-C3	3.25	116.98	110.36
8	B	921	IDR	C1-O5-C5	3.05	116.70	112.22
8	A	913	IDR	C1-O5-C5	2.87	116.44	112.22
8	B	921	IDR	C4-C3-C2	2.66	115.50	110.83
8	A	913	IDR	C4-C3-C2	2.65	115.48	110.83
7	A	912	TLA	O41-C4-C3	2.39	119.94	113.31
8	B	921	IDR	O6B-C6-C5	2.28	121.86	113.64
7	A	912	TLA	O11-C1-C2	2.27	119.62	113.31
7	B	920	TLA	O11-C1-C2	2.18	119.36	113.31
8	B	921	IDR	O1-C1-O5	-2.09	104.21	110.41
7	B	919	TLA	O11-C1-C2	2.07	119.07	113.31

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	910	GOL	O1-C1-C2-O2
6	A	910	GOL	O1-C1-C2-C3
6	B	913	GOL	C1-C2-C3-O3
6	B	913	GOL	O2-C2-C3-O3
6	B	917	GOL	O1-C1-C2-C3
7	B	920	TLA	O1-C1-C2-O2
7	B	920	TLA	O11-C1-C2-O2
7	B	920	TLA	C2-C3-C4-O4
7	B	920	TLA	C2-C3-C4-O41
7	B	920	TLA	O3-C3-C4-O4
7	B	920	TLA	O3-C3-C4-O41
6	A	910	GOL	C1-C2-C3-O3
6	B	915	GOL	O1-C1-C2-C3
6	B	916	GOL	C1-C2-C3-O3
6	B	918	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	910	GOL	O2-C2-C3-O3
6	B	916	GOL	O2-C2-C3-O3
6	B	917	GOL	O1-C1-C2-O2
6	B	918	GOL	O2-C2-C3-O3
6	A	911	GOL	O2-C2-C3-O3
7	B	919	TLA	C2-C3-C4-O4
6	B	915	GOL	O1-C1-C2-O2
7	B	919	TLA	C2-C3-C4-O41
6	B	917	GOL	C1-C2-C3-O3
6	B	917	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	918	GOL	1	0
7	A	912	TLA	1	0
6	B	913	GOL	2	0
6	B	917	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	595/627 (94%)	0.24	26 (4%) 39 38	25, 39, 67, 92	2 (0%)
1	B	606/627 (96%)	-0.10	21 (3%) 47 47	14, 33, 58, 87	5 (0%)
All	All	1201/1254 (95%)	0.07	47 (3%) 43 43	14, 35, 64, 92	7 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	THR	4.2
1	A	64	TYR	4.2
1	B	589	GLY	4.1
1	B	252	ALA	3.8
1	A	639	LEU	3.6
1	B	340	ALA	3.5
1	A	337	THR	3.4
1	B	590	LYS	3.4
1	B	186	HIS	3.3
1	A	339	SER	3.3
1	B	64	TYR	3.2
1	B	338	THR	3.0
1	A	588	ASP	3.0
1	A	431	ALA	3.0
1	A	340	ALA	3.0
1	A	430	PRO	2.9
1	A	186	HIS	2.8
1	B	62	ASP	2.8
1	A	28	ALA	2.7
1	A	522	GLY	2.7
1	B	642	PRO	2.7
1	B	28	ALA	2.6
1	A	336	ASN	2.6
1	B	229	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	335	ALA	2.6
1	A	517	PRO	2.5
1	B	337	THR	2.5
1	B	189	ASP	2.4
1	B	341	PHE	2.4
1	B	336	ASN	2.4
1	A	429	GLY	2.4
1	A	65	VAL	2.4
1	A	546	LYS	2.4
1	B	521	GLY	2.3
1	A	521	GLY	2.3
1	B	339	SER	2.3
1	B	65	VAL	2.3
1	A	627	ALA	2.3
1	A	66	LEU	2.3
1	B	103	THR	2.2
1	A	55	PRO	2.2
1	A	466	GLY	2.1
1	A	626	TRP	2.1
1	B	426	ARG	2.1
1	A	451	ASN	2.0
1	B	428	GLN	2.0
1	A	523	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	F	5	11/12	0.57	0.23	71,85,89,93	0
2	NAG	E	2	14/15	0.64	0.26	79,84,91,95	0
4	MAN	F	7	11/12	0.65	0.22	80,84,87,90	0
2	NAG	C	2	14/15	0.74	0.17	78,82,88,88	0
4	MAN	F	6	11/12	0.74	0.21	84,93,95,95	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAN	D	4	11/12	0.74	0.16	76,82,87,87	0
4	MAN	F	9	11/12	0.78	0.15	71,74,77,77	0
3	MAN	D	5	11/12	0.81	0.14	68,73,75,76	0
4	MAN	F	4	11/12	0.83	0.14	63,70,78,79	0
2	NAG	E	1	14/15	0.89	0.10	47,53,61,70	0
3	BMA	D	3	11/12	0.91	0.09	57,64,71,72	0
4	MAN	F	8	11/12	0.91	0.11	53,56,59,65	0
3	NAG	D	1	14/15	0.91	0.11	44,46,49,52	0
2	NAG	C	1	14/15	0.92	0.10	46,53,60,70	0
4	NAG	F	2	14/15	0.93	0.09	32,35,39,40	0
3	NAG	D	2	14/15	0.94	0.10	47,50,53,57	0
4	BMA	F	3	11/12	0.94	0.08	42,47,53,55	0
4	NAG	F	1	14/15	0.95	0.08	33,34,36,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	B	918	6/6	0.71	0.20	67,69,72,76	0
5	NAG	A	901	14/15	0.77	0.17	71,75,76,77	0
6	GOL	A	911	6/6	0.78	0.19	47,49,51,53	0
8	IDR	A	913	13/13	0.83	0.13	49,54,58,59	0
7	TLA	B	920	10/10	0.84	0.15	78,84,86,88	0
6	GOL	B	916	6/6	0.84	0.16	46,49,50,52	0
7	TLA	A	912	10/10	0.85	0.12	55,58,59,62	0
10	SO4	A	915	5/5	0.85	0.13	82,82,85,88	0
8	IDR	B	921	13/13	0.87	0.11	42,53,58,60	0
6	GOL	A	910	6/6	0.88	0.20	63,64,65,66	0
6	GOL	B	914	6/6	0.89	0.13	40,41,42,44	0
6	GOL	A	909	6/6	0.90	0.14	52,55,56,58	0
6	GOL	B	915	6/6	0.90	0.14	49,51,52,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	B	913	6/6	0.90	0.12	44,44,44,45	0
5	NAG	B	901	14/15	0.93	0.09	44,49,52,55	0
7	TLA	B	919	10/10	0.93	0.09	37,40,46,46	0
6	GOL	B	917	6/6	0.94	0.11	36,36,37,37	0
9	CL	A	914	1/1	0.97	0.07	37,37,37,37	0
9	CL	B	922	1/1	0.99	0.03	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.