



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:15 AM UTC

PDB ID : 1OC2 / pdb_00001oc2
Title : The structure of NADH in the dTDP-D-glucose dehydratase (RmlB) enzyme
Authors : Beis, K.; Naismith, J.H.
Deposited on : 2003-02-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	3% 88% 10% ...
1	B	348	5% 89% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	1352	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6465 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DTDP-GLUCOSE 4,6-DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	S	0	13	0
			2787	1760	474	550	3			
1	B	347	Total	C	N	O	S	0	4	1
			2750	1737	471	539	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	313	GLU	GLN	conflict	UNP Q8GIP9
B	313	GLU	GLN	conflict	UNP Q8GIP9

- Molecule 2 is THYMIDINE-5'-DIPHOSPHO-BETA-D-XYLOSE (CCD ID: TDX) (formula: C₁₅H₂₄N₂O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			34	15	2	15	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	34	15	2	15	2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	44	21	7	14	2	0	0
3	B	1	44	21	7	14	2	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	404	Total	O	0	0
			404	404		
5	B	353	Total	O	0	0
			353	353		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DTDP-GLUCOSE 4,6-DEHYDRATASE



• Molecule 1: DTDP-GLUCOSE 4,6-DEHYDRATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.41Å 94.83Å 183.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.29 – 1.50 91.29 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (91.29-1.50) 98.3 (91.29-1.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.170 , 0.200 0.184 , 0.207	Depositor DCC
R_{free} test set	16689 reflections (9.89%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6465	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, TDX, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.10	4/2924 (0.1%)	1.00	7/3973 (0.2%)
1	B	1.03	3/2838 (0.1%)	0.94	0/3860
All	All	1.07	7/5762 (0.1%)	0.97	7/7833 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131[A]	ASP	CA-CB	7.88	1.67	1.53
1	A	131[B]	ASP	CA-CB	7.88	1.67	1.53
1	A	49[A]	GLU	CD-OE2	7.86	1.40	1.25
1	A	49[B]	GLU	CD-OE2	7.86	1.40	1.25
1	B	347	ILE	C-N	-6.38	1.24	1.33
1	B	131[A]	ASP	CA-CB	6.07	1.63	1.53
1	B	131[B]	ASP	CA-CB	6.07	1.63	1.53

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	GLN	CA-CB-CG	7.62	129.33	114.10
1	A	131[A]	ASP	N-CA-C	6.50	120.29	109.95
1	A	131[B]	ASP	N-CA-C	6.50	120.29	109.95
1	A	131[A]	ASP	CB-CA-C	-6.03	98.34	109.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131[B]	ASP	CB-CA-C	-6.03	98.34	109.37
1	A	130	GLY	CA-C-N	5.45	131.65	122.87
1	A	130	GLY	C-N-CA	5.45	131.65	122.87

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	49[A]	GLU	Sidechain
1	A	49[B]	GLU	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2787	0	2655	34	0
1	B	2750	0	2627	45	0
2	A	34	0	22	3	0
2	B	34	0	22	2	0
3	A	44	0	26	4	0
3	B	44	0	26	4	0
4	A	15	0	0	4	0
5	A	404	0	0	17	0
5	B	353	0	0	19	0
All	All	6465	0	5378	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265[A]:GLU:CD	5:A:2278:HOH:O	1.82	1.19
1:B:19[B]:ASN:ND2	1:B:230:THR:HG23	1.58	1.18
1:A:265[A]:GLU:OE2	5:A:2278:HOH:O	1.60	1.18

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1352:SO4:O1	5:A:2398:HOH:O	1.71	1.07
1:A:131[B]:ASP:OD1	5:A:2169:HOH:O	1.71	1.07
4:A:1352:SO4:S	5:A:2398:HOH:O	2.17	1.02
1:B:154[B]:ASN:OD1	5:B:2196:HOH:O	1.80	0.97
1:B:19[B]:ASN:HD22	1:B:230:THR:HG23	1.14	0.97
4:A:1352:SO4:O1	5:A:2401:HOH:O	1.82	0.96
1:A:131[B]:ASP:OD2	5:A:2170:HOH:O	1.84	0.95
1:B:154[B]:ASN:OD1	5:B:2194:HOH:O	1.89	0.90
4:A:1352:SO4:S	5:A:2401:HOH:O	2.32	0.88
1:B:19[B]:ASN:CG	5:B:2015:HOH:O	2.20	0.85
1:B:19[B]:ASN:HD22	1:B:230:THR:CG2	1.87	0.85
1:B:2:SER:OG	5:B:2001:HOH:O	1.95	0.84
1:B:19[B]:ASN:ND2	1:B:230:THR:CG2	2.42	0.82
1:B:19[B]:ASN:HD21	1:B:231:ASN:HD22	1.28	0.81
1:B:19[A]:ASN:HB3	5:B:2015:HOH:O	1.82	0.79
1:B:229:HIS:HD2	1:B:231:ASN:H	1.31	0.78
1:B:22:HIS:HE1	1:B:47:ASN:O	1.68	0.77
1:B:2:SER:HA	5:B:2002:HOH:O	1.85	0.76
1:B:19[B]:ASN:ND2	5:B:2015:HOH:O	2.21	0.73
1:B:131[B]:ASP:OD1	5:B:2165:HOH:O	2.07	0.73
1:B:19[B]:ASN:HD21	1:B:231:ASN:ND2	1.88	0.72
1:A:131[B]:ASP:CG	5:A:2169:HOH:O	2.24	0.70
1:A:327:ASP:HB2	5:A:2352:HOH:O	1.98	0.64
1:B:3:GLN:NE2	1:B:242:THR:O	2.31	0.63
2:A:1349:TDX:H5D	5:A:2211:HOH:O	1.98	0.61
1:B:131[B]:ASP:CG	5:B:2165:HOH:O	2.42	0.61
1:B:2:SER:CA	5:B:2002:HOH:O	2.45	0.60
3:B:1348:NAD:H4N	5:B:2218:HOH:O	2.02	0.59
1:A:81[A]:HIS:CE1	1:A:109:LEU:HD12	2.38	0.58
1:A:154:ASN:HA	1:B:154[B]:ASN:OD1	2.03	0.58
1:B:131[A]:ASP:CG	1:B:289:LEU:H	2.12	0.58
1:A:188:CYS:HB3	3:A:1350:NAD:H5N	1.84	0.57
1:B:283:ASP:OD1	1:B:287:HIS:HD2	1.87	0.57
1:A:197:HIS:HD2	1:A:199:GLU:H	1.54	0.56
1:B:188:CYS:HB3	3:B:1348:NAD:H5N	1.88	0.56
1:A:16:ILE:HD11	3:A:1350:NAD:C7N	2.37	0.55
1:B:313:GLU:HG3	5:B:2312:HOH:O	2.06	0.54
1:A:173:LYS:HE2	1:A:185:ILE:HD12	1.89	0.54
1:B:173:LYS:HE2	1:B:185:ILE:HD12	1.92	0.51
1:A:81[A]:HIS:ND1	1:A:109:LEU:CD1	2.74	0.51
1:A:305:THR:HG23	5:A:2329:HOH:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81[A]:HIS:CE1	1:A:109:LEU:CD1	2.94	0.50
3:B:1348:NAD:C3N	2:B:1349:TDX:H4'	2.42	0.50
1:B:155:TYR:OH	5:B:2199:HOH:O	2.18	0.50
1:B:122:HIS:HE1	1:B:169:ASP:OD1	1.95	0.50
1:A:127:GLU:OE1	2:A:1349:TDX:H5B	2.11	0.49
1:B:62:ASP:OD2	1:B:100:HIS:HE1	1.96	0.49
1:A:23:TYR:OH	1:A:28:HIS:HE1	1.95	0.49
1:B:131[A]:ASP:OD1	1:B:289:LEU:N	2.47	0.48
1:A:265[B]:GLU:HG2	5:A:2278:HOH:O	2.14	0.47
1:A:259:ASN:O	1:A:262[A]:GLU:HG2	2.15	0.47
1:B:122:HIS:CE1	1:B:169:ASP:OD1	2.67	0.47
1:A:275:LYS:O	5:A:2278:HOH:O	2.21	0.47
1:A:62:ASP:OD2	1:A:100:HIS:HE1	1.98	0.46
1:A:347:ILE:O	5:A:2385:HOH:O	2.21	0.46
1:B:173:LYS:CE	1:B:185:ILE:HD12	2.46	0.46
1:B:120:PHE:O	1:B:183:ALA:HA	2.17	0.45
1:A:32:HIS:HE1	5:A:2068:HOH:O	1.99	0.45
1:A:197:HIS:CD2	1:A:199:GLU:H	2.35	0.45
2:B:1349:TDX:H5D	5:B:2218:HOH:O	2.18	0.44
1:A:159:SER:O	1:A:163[A]:SER:HB2	2.18	0.44
1:B:19[B]:ASN:ND2	1:B:231:ASN:HD22	2.06	0.44
1:A:120:PHE:O	1:A:183:ALA:HA	2.17	0.44
1:B:16:ILE:HD11	3:B:1348:NAD:C7N	2.48	0.43
2:A:1349:TDX:H4'	3:A:1350:NAD:C3N	2.48	0.43
1:A:16:ILE:HD11	3:A:1350:NAD:N7N	2.34	0.43
1:A:173:LYS:CE	1:A:185:ILE:HD12	2.49	0.43
1:A:96[B]:SER:OG	1:A:97:PRO:HD3	2.19	0.42
1:B:32:HIS:HE1	5:B:2073:HOH:O	2.02	0.42
1:A:115:LYS:NZ	5:A:2151:HOH:O	2.53	0.42
1:A:201:PHE:O	1:A:205:GLN:HG2	2.20	0.41
1:B:81:HIS:HD2	1:B:122:HIS:HD2	1.68	0.41
1:B:258:LYS:HD3	5:B:2269:HOH:O	2.20	0.41
1:B:28:HIS:CE1	1:B:238:TRP:CE2	3.08	0.41
1:B:28:HIS:HD2	5:B:2001:HOH:O	2.03	0.41
1:B:186:SER:HA	1:B:250:TYR:O	2.21	0.41
1:B:16:ILE:HG22	1:B:82:TYR:CE2	2.56	0.40
1:B:263:VAL:HA	5:B:2269:HOH:O	2.21	0.40
1:A:163[A]:SER:HB3	1:B:171:ILE:HG22	2.02	0.40
1:A:191:ASN:HA	1:A:228:ILE:O	2.21	0.40
1:A:154:ASN:HA	1:B:154[B]:ASN:CG	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/348 (103%)	348 (98%)	9 (2%)	0	100	100
1	B	349/348 (100%)	341 (98%)	8 (2%)	0	100	100
All	All	706/696 (101%)	689 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/291 (104%)	291 (96%)	11 (4%)	31	6
1	B	293/291 (101%)	288 (98%)	5 (2%)	53	26
All	All	595/582 (102%)	579 (97%)	16 (3%)	50	12

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49[A]	GLU
1	A	49[B]	GLU
1	A	131[A]	ASP
1	A	131[B]	ASP
1	A	143	GLU
1	A	152[A]	GLU
1	A	152[B]	GLU
1	A	163[A]	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	163[B]	SER
1	A	191	ASN
1	A	245	ARG
1	B	3	GLN
1	B	131[A]	ASP
1	B	131[B]	ASP
1	B	191	ASN
1	B	246	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	32	HIS
1	A	100	HIS
1	A	122	HIS
1	A	197	HIS
1	A	231	ASN
1	A	260	ASN
1	A	280	HIS
1	B	22	HIS
1	B	28	HIS
1	B	32	HIS
1	B	87	HIS
1	B	100	HIS
1	B	122	HIS
1	B	191	ASN
1	B	197	HIS
1	B	229	HIS
1	B	231	ASN
1	B	233	HIS
1	B	260	ASN
1	B	280	HIS
1	B	287	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1353	-	4,4,4	0.54	0	6,6,6	0.71	0
3	NAD	B	1348	-	46,48,48	2.59	14 (30%)	64,73,73	3.44	16 (25%)
3	NAD	A	1350	-	46,48,48	2.24	10 (21%)	64,73,73	2.76	15 (23%)
4	SO4	A	1351	-	4,4,4	0.48	0	6,6,6	0.62	0
2	TDX	A	1349	-	35,36,36	1.38	5 (14%)	52,55,55	1.41	8 (15%)
4	SO4	A	1352	-	4,4,4	0.29	0	6,6,6	0.35	0
2	TDX	B	1349	-	35,36,36	1.19	4 (11%)	52,55,55	1.83	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	B	1348	-	-	5/30/62/62	0/5/5/5
2	TDX	B	1349	-	-	1/21/50/50	0/3/3/3
2	TDX	A	1349	-	-	0/21/50/50	0/3/3/3
3	NAD	A	1350	-	-	4/30/62/62	0/5/5/5

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1348	NAD	C4N-C3N	10.76	1.55	1.39
3	A	1350	NAD	C4N-C3N	10.11	1.54	1.39
3	B	1348	NAD	C5N-C4N	6.62	1.50	1.38
3	B	1348	NAD	C2N-N1N	5.64	1.41	1.35
3	A	1350	NAD	C5N-C4N	5.54	1.48	1.38
3	A	1350	NAD	C2N-N1N	5.27	1.40	1.35
3	B	1348	NAD	C6N-N1N	4.62	1.45	1.35
3	A	1350	NAD	C6N-N1N	4.05	1.44	1.35
2	A	1349	TDX	C4-N3	-3.76	1.31	1.38
3	B	1348	NAD	C5A-C4A	3.44	1.45	1.39
2	B	1349	TDX	PB-O3A	3.39	1.63	1.59
3	A	1350	NAD	C4A-N3A	3.25	1.40	1.34
2	A	1349	TDX	C2-N1	-2.98	1.33	1.38
3	B	1348	NAD	C4A-N9A	-2.98	1.31	1.37
2	B	1349	TDX	C4-N3	-2.97	1.33	1.38
2	A	1349	TDX	O2-C2	2.93	1.28	1.23
3	B	1348	NAD	C7N-N7N	2.88	1.38	1.33
2	A	1349	TDX	C2-N3	-2.88	1.33	1.38
3	B	1348	NAD	C6N-C5N	-2.77	1.32	1.38
3	B	1348	NAD	C4A-N3A	2.73	1.39	1.34
2	B	1349	TDX	O2-C2	2.72	1.27	1.23
3	B	1348	NAD	O4D-C4D	-2.71	1.39	1.45
3	A	1350	NAD	C5A-C6A	2.58	1.48	1.41
2	B	1349	TDX	C6-N1	-2.42	1.33	1.38
3	A	1350	NAD	C6N-C5N	-2.36	1.33	1.38
3	B	1348	NAD	C5A-C6A	2.35	1.47	1.41
3	A	1350	NAD	C8A-N9A	-2.34	1.33	1.37
3	B	1348	NAD	O3D-C3D	2.32	1.48	1.43
3	B	1348	NAD	C5A-N7A	-2.22	1.35	1.39
2	A	1349	TDX	C4'-C3'	-2.09	1.49	1.52
3	B	1348	NAD	C8A-N7A	2.06	1.35	1.31
3	A	1350	NAD	C5A-N7A	-2.04	1.35	1.39
3	A	1350	NAD	C7N-N7N	2.03	1.36	1.33

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1348	NAD	C5N-C4N-C3N	-16.65	104.00	120.36
3	B	1348	NAD	C6N-N1N-C2N	-13.67	110.25	121.88
3	A	1350	NAD	C5N-C4N-C3N	-13.60	107.00	120.36
2	B	1349	TDX	O4-C4-C5	-7.92	115.86	124.92
3	A	1350	NAD	O7N-C7N-C3N	-7.50	110.42	119.60
3	B	1348	NAD	C3N-C7N-N7N	7.48	126.96	117.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1350	NAD	C6N-N1N-C2N	-7.39	115.59	121.88
3	B	1348	NAD	O7N-C7N-C3N	-7.21	110.78	119.60
3	A	1350	NAD	C2N-C3N-C4N	6.33	125.62	118.26
3	B	1348	NAD	C6N-N1N-C1D	-5.97	108.00	119.73
2	B	1349	TDX	C5-C4-N3	5.05	119.71	115.32
3	A	1350	NAD	C6N-N1N-C1D	-4.77	110.36	119.73
3	B	1348	NAD	C5A-C4A-N3A	-4.59	120.39	126.72
3	A	1350	NAD	C3N-C7N-N7N	4.12	122.82	117.74
3	B	1348	NAD	N3A-C2A-N1A	-3.91	122.67	128.58
2	B	1349	TDX	C4-N3-C2	-3.86	122.27	127.34
3	A	1350	NAD	C2A-N1A-C6A	3.85	125.05	118.73
3	A	1350	NAD	N3A-C2A-N1A	-3.84	122.78	128.58
2	A	1349	TDX	N3-C2-N1	3.63	119.62	114.89
3	B	1348	NAD	C2A-N3A-C4A	3.60	120.62	111.83
3	B	1348	NAD	C6A-C5A-N7A	3.57	138.98	132.09
2	B	1349	TDX	C6-C5-C4	-3.13	115.44	118.02
3	B	1348	NAD	C4A-C5A-N7A	-2.99	107.16	110.58
3	A	1350	NAD	N6A-C6A-N1A	2.93	124.91	118.38
2	A	1349	TDX	C4'-C3'-C2'	-2.86	105.84	110.86
2	A	1349	TDX	C4-N3-C2	-2.86	123.60	127.34
3	B	1348	NAD	C5A-C6A-N6A	-2.84	116.25	123.29
2	A	1349	TDX	C5-C4-N3	2.80	117.76	115.32
2	B	1349	TDX	N3-C2-N1	2.69	118.39	114.89
3	A	1350	NAD	O7N-C7N-N7N	2.56	126.31	122.62
3	A	1350	NAD	C6A-C5A-C4A	-2.52	113.73	117.18
2	A	1349	TDX	C5-C6-N1	-2.44	120.67	123.31
3	B	1348	NAD	C5A-C4A-N9A	2.44	108.47	105.81
3	B	1348	NAD	C6A-C5A-C4A	-2.43	113.86	117.18
3	A	1350	NAD	C6N-C5N-C4N	2.38	122.88	119.45
2	A	1349	TDX	C2B-C1B-N1	2.37	119.74	113.81
3	A	1350	NAD	C5A-C6A-N6A	-2.37	117.42	123.29
3	B	1348	NAD	N3A-C4A-N9A	2.34	131.14	127.17
2	B	1349	TDX	O4B-C1B-N1	-2.34	103.72	107.86
2	A	1349	TDX	O4-C4-C5	-2.30	122.28	124.92
2	A	1349	TDX	O4'-C4'-C5'	-2.30	103.95	109.22
3	B	1348	NAD	C6N-C5N-C4N	2.23	122.66	119.45
2	B	1349	TDX	O3A-PA-O1A	-2.23	104.01	110.70
2	B	1349	TDX	O4-C4-N3	2.19	124.23	120.11
3	A	1350	NAD	O2N-PN-O1N	2.10	122.23	112.44
3	B	1348	NAD	C2N-C3N-C4N	2.10	120.70	118.26
3	A	1350	NAD	C6A-C5A-N7A	2.04	136.03	132.09

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1350	NAD	O4D-C1D-N1N-C2N
3	A	1350	NAD	O4D-C1D-N1N-C6N
3	A	1350	NAD	C2D-C1D-N1N-C6N
3	B	1348	NAD	C5D-O5D-PN-O1N
3	B	1348	NAD	O4D-C1D-N1N-C6N
3	B	1348	NAD	C2D-C1D-N1N-C6N
2	B	1349	TDX	C1'-O1'-PB-O3A
3	B	1348	NAD	C5D-O5D-PN-O2N
3	A	1350	NAD	PN-O3-PA-O2A
3	B	1348	NAD	PN-O3-PA-O2A

There are no ring outliers.


5 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1348	NAD	4	0
3	A	1350	NAD	4	0
2	A	1349	TDX	3	0
4	A	1352	SO4	4	0
2	B	1349	TDX	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand NAD A 1350



Ligand TDX A 1349





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	346/348 (99%)	-0.10	12 (3%)	47 51	7, 14, 23, 35	18 (5%)
1	B	347/348 (99%)	0.46	19 (5%)	30 33	6, 17, 28, 39	13 (3%)
All	All	693/696 (99%)	0.18	31 (4%)	38 42	6, 16, 27, 39	31 (4%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	141	HIS	7.8
1	B	121	HIS	7.3
1	B	262	GLU	5.7
1	B	332	GLU	5.5
1	B	85	GLU	5.4
1	B	269	GLU	5.4
1	B	248	GLU	5.1
1	B	265	GLU	4.7
1	B	2	SER	4.3
1	A	332	GLU	4.2
1	B	272	GLY	4.1
1	A	67	GLU	4.0
1	A	334	GLU	3.9
1	B	67	GLU	3.5
1	A	141	HIS	3.4
1	A	3	GLN	3.2
1	A	327	ASP	3.1
1	B	308	PHE	3.0
1	A	71	LYS	2.9
1	B	276	ASP	2.9
1	A	131[A]	ASP	2.8
1	B	277	ALA	2.6
1	A	275	LYS	2.6
1	B	131[A]	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	334	GLU	2.5
1	B	3	GLN	2.5
1	A	143	GLU	2.5
1	A	308	PHE	2.4
1	B	275	LYS	2.3
1	A	145	PRO	2.2
1	B	347	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	1352	5/5	0.69	0.24	90,90,90,90	0
3	NAD	A	1350	44/44	0.98	0.08	12,14,25,26	0
3	NAD	B	1348	44/44	0.98	0.07	16,18,31,32	0
4	SO4	A	1351	5/5	0.98	0.06	26,26,30,30	0
2	TDX	B	1349	34/34	0.98	0.07	18,20,22,25	0
4	SO4	A	1353	5/5	0.98	0.07	29,31,33,37	0
2	TDX	A	1349	34/34	0.99	0.06	12,14,17,19	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAD A 1350:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD B 1348:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TDX B 1349:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TDX A 1349:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.