



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 09:37 AM UTC

PDB ID : 5OCC / pdb_00005occ
Title : Crystal structure of CD32b (Fc Gamma Receptor IIb) in complex with Human IgG1 Fab fragment (6G08)
Authors : Tews, I.; Orr, C.
Deposited on : 2017-06-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	11% 71% 18% • 8%
2	H	222	81% 19%
3	L	217	% 77% 20% •
4	B	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	B	1	X	-	-	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4677 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Low affinity immunoglobulin gamma Fc region receptor II-b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1291	812	224	251	4			

- Molecule 2 is a protein called 6G08 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1624	1020	272	327	5			

- Molecule 3 is a protein called 6G08 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	216	Total	C	N	O	S	0	0	0
			1598	996	270	327	5			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	L	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	24	Total	O	0	0
			24	24		
8	H	50	Total	O	0	0
			50	50		
8	L	31	Total	O	0	0
			31	31		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Low affinity immunoglobulin gamma Fc region receptor II-b



- Molecule 2: 6G08 Fab heavy chain



- Molecule 3: 6G08 Fab Light Chain



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.23Å 75.02Å 134.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	134.95 – 2.50 67.47 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.2 (134.95-2.50) 99.3 (67.47-2.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.192 , 0.261 0.200 , 0.255	Depositor DCC
R_{free} test set	1281 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.352	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4677	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.24	4/1326 (0.3%)	1.14	3/1807 (0.2%)
2	H	1.51	8/1660 (0.5%)	1.26	5/2261 (0.2%)
3	L	1.46	7/1638 (0.4%)	1.37	10/2238 (0.4%)
All	All	1.42	19/4624 (0.4%)	1.27	18/6306 (0.3%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	ASP	N-CA	7.41	1.55	1.46
3	L	73	ALA	CA-C	-7.19	1.43	1.52
2	H	174	PHE	CA-C	-6.94	1.44	1.53
2	H	191	THR	C-O	-6.69	1.15	1.24
1	A	161	VAL	C-O	-6.46	1.17	1.24
3	L	7	PRO	CA-C	-6.36	1.46	1.52
2	H	171	VAL	C-O	-5.96	1.17	1.24
2	H	206	VAL	C-O	-5.70	1.17	1.24
2	H	211	SER	N-CA	5.70	1.53	1.46
2	H	114	GLY	C-O	-5.60	1.17	1.23
3	L	120	VAL	C-O	-5.53	1.18	1.24
3	L	6	GLN	C-O	-5.51	1.18	1.24
3	L	128	GLU	C-O	-5.40	1.17	1.24
2	H	95	TYR	C-O	-5.39	1.17	1.23
2	H	173	THR	N-CA	-5.31	1.40	1.46
1	A	79	SER	CA-C	5.27	1.59	1.52
3	L	120	VAL	CA-CB	-5.23	1.48	1.54
3	L	163	GLY	C-O	-5.07	1.17	1.24
1	A	151	ARG	C-O	-5.01	1.18	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	95	THR	N-CA-C	10.24	123.55	110.65
3	L	157	GLY	N-CA-C	-8.77	103.83	115.32
2	H	48	VAL	N-CA-CB	-8.05	105.03	111.64
3	L	215	GLU	N-CA-C	7.70	121.57	110.10
2	H	196	SER	N-CA-C	6.99	121.25	112.87
1	A	80	ASP	N-CA-C	6.76	125.20	110.80
3	L	105	GLY	N-CA-C	-6.55	104.96	112.29
2	H	174	PHE	CA-C-N	-5.93	113.57	119.92
2	H	174	PHE	C-N-CA	-5.93	113.57	119.92
3	L	54	THR	CB-CA-C	-5.89	100.39	109.29
3	L	214	THR	CA-C-N	5.65	128.79	120.82
3	L	214	THR	C-N-CA	5.65	128.79	120.82
3	L	168	LYS	CA-C-N	-5.48	114.94	120.31
3	L	168	LYS	C-N-CA	-5.48	114.94	120.31
1	A	118	THR	N-CA-C	5.34	116.92	108.96
2	H	174	PHE	N-CA-C	-5.19	102.67	110.24
3	L	197	SER	N-CA-C	5.04	117.05	109.23
3	L	177	TYR	N-CA-C	5.04	118.23	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1291	0	1227	14	0
2	H	1624	0	1591	11	0
3	L	1598	0	1543	19	0
4	B	28	0	25	0	0
5	A	14	0	13	0	0
6	A	12	0	16	0	0
7	L	5	0	0	0	0
8	A	24	0	0	1	0
8	H	50	0	0	0	0
8	L	31	0	0	0	0
All	All	4677	0	4415	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:125:PRO:HD3	3:L:137:LEU:HD23	1.53	0.88
3:L:125:PRO:CD	3:L:137:LEU:HD23	2.30	0.60
1:A:70:THR:HG22	1:A:98:SER:OG	2.03	0.59
1:A:115:THR:HG22	1:A:118:THR:OG1	2.04	0.57
3:L:82:SER:OG	3:L:175:ASN:HB3	2.09	0.52
2:H:20:LEU:HG	2:H:83:MET:HE2	1.92	0.51
3:L:56:ARG:HD2	3:L:60:VAL:HG12	1.93	0.51
3:L:117:ASN:O	3:L:117:ASN:ND2	2.45	0.50
2:H:2:VAL:HG13	2:H:27:PHE:CD1	2.47	0.50
2:H:178:LEU:HD12	2:H:179:GLN:N	2.27	0.50
1:A:67:VAL:O	1:A:100:ARG:HA	2.12	0.49
1:A:92:PRO:O	1:A:94:HIS:N	2.44	0.49
1:A:55:GLU:HB3	1:A:68:THR:HB	1.93	0.49
2:H:179:GLN:NE2	2:H:185:SER:OG	2.46	0.48
3:L:52:GLY:O	3:L:53:ASP:HB2	2.12	0.48
3:L:213:PRO:O	3:L:215:GLU:N	2.46	0.48
3:L:117:ASN:ND2	3:L:117:ASN:C	2.72	0.48
2:H:209:LYS:N	2:H:210:PRO:CD	2.77	0.48
1:A:92:PRO:O	1:A:95:THR:HG23	2.14	0.48
1:A:105:ASN:ND2	1:A:130:SER:OG	2.46	0.47
2:H:12:VAL:HG21	2:H:18:LEU:HD22	1.95	0.47
1:A:62:LEU:HD21	1:A:134:VAL:HB	1.96	0.47
3:L:190:TRP:CZ2	3:L:213:PRO:HA	2.49	0.47
3:L:17:ARG:HA	3:L:77:ILE:O	2.14	0.46
2:H:155:PRO:O	2:H:208:HIS:HE1	1.98	0.46
3:L:117:ASN:HB2	3:L:118:PRO:HD2	1.98	0.46
1:A:96:GLN:HB3	1:A:97:PRO:CD	2.47	0.45
1:A:181:PHE:CZ	1:A:183:ILE:HD11	2.51	0.45
2:H:12:VAL:HG12	2:H:13:GLN:O	2.18	0.44
1:A:115:THR:HG23	1:A:117:GLN:N	2.33	0.44
3:L:145:TYR:C	3:L:145:TYR:CD1	2.95	0.44
1:A:84:TRP:CZ3	1:A:95:THR:HA	2.54	0.42
1:A:151:ARG:NH2	8:A:401:HOH:O	2.30	0.42
3:L:113:GLN:HB3	3:L:114:PRO:CD	2.50	0.42
2:H:162:TRP:CH2	2:H:204:CYS:HB3	2.55	0.42
3:L:172:GLN:OE1	3:L:178:ALA:HB2	2.20	0.41
3:L:41:LEU:HD23	3:L:86:ALA:HB2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:161:LYS:HE3	3:L:162:ALA:N	2.36	0.41
3:L:154:LYS:HE2	3:L:199:GLN:HE21	1.85	0.41
3:L:20:ILE:HG12	3:L:106:THR:HG21	2.03	0.41
1:A:82:ILE:O	1:A:95:THR:HB	2.21	0.41
3:L:174:ASN:O	3:L:175:ASN:HB2	2.21	0.41
2:H:153:TYR:CZ	2:H:158:VAL:HG13	2.56	0.41
2:H:104:ALA:HB3	2:H:107:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/176 (90%)	140 (89%)	12 (8%)	6 (4%)	2	3
2	H	220/222 (99%)	215 (98%)	4 (2%)	1 (0%)	24	43
3	L	214/217 (99%)	201 (94%)	12 (6%)	1 (0%)	24	43
All	All	592/615 (96%)	556 (94%)	28 (5%)	8 (1%)	9	17

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	89	ASN
1	A	92	PRO
1	A	96	GLN
1	A	106	ASN
1	A	188	HIS
2	H	29	LEU
3	L	34	ASP
1	A	79	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/161 (93%)	142 (95%)	8 (5%)	20	42
2	H	181/181 (100%)	168 (93%)	13 (7%)	13	28
3	L	178/179 (99%)	169 (95%)	9 (5%)	21	43
All	All	509/521 (98%)	479 (94%)	30 (6%)	18	37

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	ILE
1	A	63	GLN
1	A	104	ASN
1	A	124	VAL
1	A	127	THR
1	A	144	GLU
1	A	182	SER
1	A	189	SER
2	H	5	LEU
2	H	6	GLU
2	H	19	ARG
2	H	62	ASP
2	H	63	SER
2	H	64	VAL
2	H	136	SER
2	H	143	THR
2	H	158	VAL
2	H	177	VAL
2	H	194	SER
2	H	200	GLN
2	H	201	THR
3	L	1	GLN
3	L	13	THR
3	L	47	LYS
3	L	72	SER
3	L	98	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	117	ASN
3	L	161	LYS
3	L	197	SER
3	L	215	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	GLN
1	A	104	ASN
1	A	105	ASN
1	A	199	ASN
2	H	172	HIS
2	H	179	GLN
2	H	208	HIS
3	L	39	GLN
3	L	117	ASN
3	L	175	ASN
3	L	193	HIS
3	L	202	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	1	1,4	14,14,15	0.98	1 (7%)	17,19,21	1.98	4 (23%)
4	NAG	B	2	4	14,14,15	1.19	1 (7%)	17,19,21	2.94	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	1,4	1/1/5/7	2/6/23/26	0/1/1/1
4	NAG	B	2	4	-	5/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	2	NAG	C1-C2	3.75	1.57	1.52
4	B	1	NAG	C1-C2	2.79	1.56	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2	NAG	C2-N2-C7	8.21	133.90	122.90
4	B	2	NAG	C8-C7-N2	5.70	125.57	116.12
4	B	1	NAG	O5-C1-C2	5.48	119.77	111.29
4	B	1	NAG	C1-O5-C5	4.25	117.88	112.19
4	B	2	NAG	O7-C7-N2	-3.48	115.83	121.98
4	B	2	NAG	O3-C3-C2	2.51	114.62	109.40
4	B	2	NAG	C1-C2-N2	-2.39	106.66	110.43
4	B	1	NAG	C1-C2-N2	2.17	113.85	110.43
4	B	2	NAG	C4-C3-C2	-2.12	107.91	111.02
4	B	1	NAG	O4-C4-C5	-2.08	104.21	109.32

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	1	NAG	C1


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2	NAG	C3-C2-N2-C7
4	B	2	NAG	O5-C5-C6-O6
4	B	2	NAG	C4-C5-C6-O6
4	B	2	NAG	C8-C7-N2-C2
4	B	2	NAG	O7-C7-N2-C2
4	B	1	NAG	O5-C5-C6-O6
4	B	1	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PO4	L	301	-	4,4,4	1.22	1 (25%)	6,6,6	1.15	1 (16%)
6	GOL	A	304	-	5,5,5	0.20	0	5,5,5	0.40	0
5	NAG	A	301	1	14,14,15	0.81	0	17,19,21	2.31	6 (35%)
6	GOL	A	305	-	5,5,5	0.53	0	5,5,5	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	304	-	-	0/4/4/4	-
5	NAG	A	301	1	-	4/6/23/26	0/1/1/1
6	GOL	A	305	-	-	4/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	301	PO4	P-O1	2.33	1.56	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	301	NAG	C1-C2-N2	5.40	118.94	110.43
5	A	301	NAG	O3-C3-C2	3.68	117.05	109.40
5	A	301	NAG	C8-C7-N2	3.60	122.09	116.12
5	A	301	NAG	O5-C1-C2	-3.30	106.19	111.29
5	A	301	NAG	O7-C7-N2	-3.26	116.22	121.98
5	A	301	NAG	O4-C4-C3	-2.24	105.09	110.38
7	L	301	PO4	O4-P-O3	2.22	114.83	107.91

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	305	GOL	O1-C1-C2-C3
5	A	301	NAG	O5-C5-C6-O6
5	A	301	NAG	C8-C7-N2-C2
5	A	301	NAG	O7-C7-N2-C2
6	A	305	GOL	O1-C1-C2-O2
6	A	305	GOL	C1-C2-C3-O3
5	A	301	NAG	C4-C5-C6-O6
6	A	305	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/176 (92%)	0.73	20 (12%) 8 7	36, 72, 120, 133	0
2	H	222/222 (100%)	-0.26	0 100 100	22, 39, 59, 80	0
3	L	216/217 (99%)	-0.23	3 (1%) 73 70	20, 41, 61, 87	0
All	All	600/615 (97%)	0.02	23 (3%) 44 39	20, 44, 101, 133	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	218	SER	3.7
1	A	217	PRO	3.4
3	L	1	GLN	3.4
1	A	91	ILE	3.1
1	A	92	PRO	2.9
3	L	117	ASN	2.9
1	A	116	GLY	2.8
1	A	97	PRO	2.8
1	A	80	ASP	2.7
1	A	120	LEU	2.7
1	A	188	HIS	2.6
1	A	90	LEU	2.5
1	A	112	THR	2.5
3	L	216	CYS	2.4
1	A	94	HIS	2.4
1	A	93	THR	2.3
1	A	79	SER	2.3
1	A	119	SER	2.2
1	A	148	ILE	2.2
1	A	100	ARG	2.2
1	A	145	GLY	2.2
1	A	125	HIS	2.1
1	A	71	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	1	14/15	-	-	93,98,105,106	0
4	NAG	B	2	14/15	-	-	89,101,115,115	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	301	14/15	0.70	0.18	95,101,104,106	0
7	PO4	L	301	5/5	0.87	0.11	42,48,52,69	0
6	GOL	A	305	6/6	0.89	0.12	70,73,76,78	0
6	GOL	A	304	6/6	0.90	0.13	58,59,65,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.