



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:32 AM UTC

PDB ID : 2ODO / pdb_00002odo
Title : Crystal structure of Pseudomonas Fluorescens alanine racemase
Authors : Tsuge, H.; Ohnishi, K.; Yokoigawa, K.
Deposited on : 2006-12-25
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 87% 11% 2% 2%
1	B	357	 86% 11% 3% 3%
1	C	357	 87% 10% 2% 2%
1	D	357	 89% 10% 2% 2%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 11074 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alanine racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	356	Total 2750	C 1749	N 484	O 500	P 1	S 16	0	0	0
1	B	356	Total 2750	C 1749	N 484	O 500	P 1	S 16	0	0	0
1	C	356	Total 2750	C 1749	N 484	O 500	P 1	S 16	0	0	0
1	D	356	Total 2750	C 1749	N 484	O 500	P 1	S 16	0	0	0

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	22	Total	O	0	0
			22	22		
3	C	8	Total	O	0	0
			8	8		
3	D	7	Total	O	0	0
			7	7		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alanine racemase

Chain A: 





- Molecule 1: Alanine racemase

Chain B: 





- Molecule 1: Alanine racemase

Chain C: 



- Molecule 1: Alanine racemase

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	186.74Å 186.74Å 103.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.32 – 2.80 46.32 – 2.80	Depositor EDS
% Data completeness (in resolution range)	93.5 (46.32-2.80) 93.6 (46.32-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.214 , 0.247 (Not available) , 0.208	Depositor DCC
R_{free} test set	2076 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	61.0	Xtrriage
Anisotropy	0.005	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.007 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11074	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, ACT, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2773	0.79	3/3774 (0.1%)
1	B	0.43	0/2773	0.78	3/3774 (0.1%)
1	C	0.43	0/2773	0.79	4/3774 (0.1%)
1	D	0.42	0/2773	0.77	3/3774 (0.1%)
All	All	0.43	0/11092	0.78	13/15096 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	VAL	CA-C-N	6.62	133.61	121.70
1	A	286	VAL	C-N-CA	6.62	133.61	121.70
1	C	286	VAL	CA-C-N	6.59	133.56	121.70
1	C	286	VAL	C-N-CA	6.59	133.56	121.70
1	B	286	VAL	CA-C-N	6.58	133.54	121.70
1	B	286	VAL	C-N-CA	6.58	133.54	121.70
1	D	286	VAL	CA-C-N	6.42	133.26	121.70
1	D	286	VAL	C-N-CA	6.42	133.26	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	286	VAL	N-CA-C	-6.12	96.60	109.34
1	D	286	VAL	N-CA-C	-5.99	96.88	109.34
1	B	286	VAL	N-CA-C	-5.84	97.19	109.34
1	C	286	VAL	N-CA-C	-5.55	97.79	109.34
1	C	285	PRO	N-CA-C	5.21	118.75	110.21

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	285	PRO	Peptide
1	B	285	PRO	Peptide
1	C	285	PRO	Peptide
1	D	285	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	0	2759	21	0
1	B	2750	0	2760	26	0
1	C	2750	0	2760	27	0
1	D	2750	0	2760	17	0
2	A	4	0	3	1	0
2	B	4	0	3	0	0
2	C	4	0	3	1	0
2	D	4	0	3	1	0
3	A	21	0	0	0	0
3	B	22	0	0	0	0
3	C	8	0	0	0	0
3	D	7	0	0	0	0
All	All	11074	0	11051	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:ARG:HG2	1:B:201:ARG:HH21	1.46	0.80
1:B:201:ARG:HH21	1:B:201:ARG:CG	2.10	0.64
1:B:287:PHE:H	1:B:292:ARG:HA	1.66	0.60
2:C:3400:ACT:H1	1:D:33:LLP:H4'1	1.84	0.59
2:A:1400:ACT:H1	1:B:33:LLP:H4'1	1.84	0.58
1:C:201:ARG:HH21	1:C:201:ARG:HG2	1.68	0.57
1:C:201:ARG:HH21	1:C:201:ARG:CG	2.18	0.56
1:B:158:HIS:HD2	1:B:208:ARG:HH21	1.54	0.55
1:A:287:PHE:H	1:A:292:ARG:HA	1.72	0.55
1:A:182:ALA:O	1:A:183:ASP:CB	2.55	0.55
1:C:90:ILE:HG23	1:C:95:PHE:HB2	1.90	0.54
1:A:265:ILE:HG23	1:A:305:CYS:SG	2.49	0.53
1:B:265:ILE:HG23	1:B:305:CYS:SG	2.49	0.53
1:C:257:PHE:CE2	1:C:263:MET:HE1	2.44	0.53
1:D:287:PHE:H	1:D:292:ARG:HA	1.74	0.52
1:B:182:ALA:O	1:B:183:ASP:CB	2.57	0.52
1:C:287:PHE:H	1:C:292:ARG:HA	1.73	0.52
1:D:182:ALA:O	1:D:183:ASP:CB	2.57	0.52
1:C:257:PHE:HE2	1:C:263:MET:HE1	1.75	0.51
1:C:259:THR:HG22	1:C:263:MET:HE2	1.94	0.50
1:C:182:ALA:O	1:C:183:ASP:CB	2.60	0.49
1:A:193:SER:HB2	1:A:194:PRO:HD3	1.93	0.49
1:B:67:ARG:NH1	1:B:72:ARG:HA	2.28	0.49
1:B:356:GLY:N	1:D:313:GLN:HE22	2.10	0.49
1:C:19:LEU:HD11	1:C:228:ARG:HB3	1.95	0.48
1:A:158:HIS:HD2	1:A:208:ARG:HH21	1.61	0.48
1:A:158:HIS:HE1	1:B:253:TYR:OH	1.97	0.48
1:A:201:ARG:HH21	1:A:201:ARG:CG	2.27	0.48
1:A:257:PHE:CE2	1:A:263:MET:HE1	2.48	0.47
1:A:182:ALA:O	1:A:183:ASP:HB3	2.14	0.47
1:C:33:LLP:H4'1	2:D:2400:ACT:H1	1.96	0.47
1:A:253:TYR:OH	1:B:158:HIS:HE1	1.98	0.47
1:A:191:ARG:HB2	1:A:207:VAL:HG12	1.97	0.46
1:B:96:TRP:CD1	1:B:118:HIS:HB2	2.51	0.46
1:A:33:LLP:HD3	1:B:302:ASP:OD2	2.17	0.45
1:C:259:THR:HG22	1:C:263:MET:CE	2.47	0.45
1:C:253:TYR:OH	1:D:158:HIS:HE1	1.99	0.45
1:A:339:ILE:HD12	1:A:341:TYR:CD1	2.51	0.44
1:A:121:LEU:HD13	1:A:144:LEU:HD11	2.00	0.44
1:C:193:SER:HB2	1:C:194:PRO:HD3	1.98	0.44
1:D:158:HIS:HD2	1:D:208:ARG:HH21	1.66	0.44
1:C:87:LEU:N	1:C:88:PRO:HD2	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:CYS:HB3	1:B:267:VAL:HB	2.00	0.43
1:C:37:TYR:CE2	1:D:301:MET:HE1	2.53	0.43
1:C:158:HIS:HE1	1:D:253:TYR:OH	2.01	0.43
1:D:20:ALA:O	1:D:24:THR:HG22	2.18	0.43
1:B:193:SER:HB2	1:B:194:PRO:HD3	2.01	0.43
1:C:20:ALA:O	1:C:24:THR:HG22	2.19	0.42
1:D:121:LEU:HD13	1:D:144:LEU:HD11	2.02	0.42
1:D:257:PHE:CE2	1:D:263:MET:HE1	2.53	0.42
1:C:67:ARG:NH1	1:C:72:ARG:HA	2.35	0.42
1:A:223:GLN:HG3	1:A:226:ALA:H	1.85	0.42
1:A:241:CYS:HB3	1:A:267:VAL:HB	2.02	0.42
1:A:261:LYS:HB2	1:A:262:PRO:HD2	2.02	0.42
1:B:261:LYS:HB2	1:B:262:PRO:HD2	2.01	0.42
1:B:87:LEU:HA	1:B:90:ILE:HD12	2.02	0.42
1:D:21:ARG:HH21	1:D:53:ASP:CG	2.28	0.42
1:A:257:PHE:HE2	1:A:263:MET:HE1	1.85	0.42
1:B:90:ILE:HG23	1:B:95:PHE:HB2	2.02	0.42
1:B:284:THR:HA	1:B:285:PRO:HD3	1.83	0.42
1:D:241:CYS:HB3	1:D:267:VAL:HB	2.01	0.42
1:A:280:ALA:HA	1:A:281:PRO:HD3	1.91	0.42
1:D:191:ARG:HB2	1:D:207:VAL:HG12	2.01	0.41
1:B:272:TYR:HA	1:B:276:TYR:O	2.21	0.41
1:B:59:CYS:SG	1:B:61:GLU:HG2	2.61	0.41
1:B:286:VAL:O	1:B:293:SER:N	2.53	0.41
1:C:201:ARG:CG	1:C:201:ARG:NH2	2.80	0.41
1:C:263:MET:HE3	1:C:265:ILE:HD11	2.03	0.41
1:B:257:PHE:CE2	1:B:263:MET:HE1	2.55	0.41
1:D:272:TYR:HA	1:D:276:TYR:O	2.21	0.41
1:A:272:TYR:HA	1:A:276:TYR:O	2.21	0.41
1:C:121:LEU:HD13	1:C:144:LEU:HD11	2.02	0.41
1:C:254:GLY:HA2	1:D:161:ARG:HG2	2.03	0.41
1:A:259:THR:HG22	1:A:263:MET:HE2	2.02	0.41
1:C:241:CYS:HB3	1:C:267:VAL:HB	2.03	0.41
1:C:272:TYR:HA	1:C:276:TYR:O	2.21	0.41
1:C:284:THR:HA	1:C:285:PRO:HD3	1.83	0.41
1:D:90:ILE:HG23	1:D:95:PHE:HB2	2.02	0.40
1:B:201:ARG:CG	1:B:201:ARG:NH2	2.75	0.40
1:C:37:TYR:N	1:C:37:TYR:CD1	2.89	0.40
1:B:37:TYR:O	1:B:344:PHE:HB3	2.21	0.40
1:B:339:ILE:HD12	1:B:341:TYR:CD1	2.56	0.40
1:C:158:HIS:HD2	1:C:208:ARG:HH21	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/357 (99%)	340 (97%)	10 (3%)	2 (1%)	21	51
1	B	352/357 (99%)	336 (96%)	14 (4%)	2 (1%)	21	51
1	C	352/357 (99%)	334 (95%)	16 (4%)	2 (1%)	21	51
1	D	352/357 (99%)	335 (95%)	15 (4%)	2 (1%)	21	51
All	All	1408/1428 (99%)	1345 (96%)	55 (4%)	8 (1%)	21	51

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	183	ASP
1	A	183	ASP
1	C	183	ASP
1	D	183	ASP
1	A	287	PHE
1	B	287	PHE
1	C	287	PHE
1	D	287	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/282 (100%)	270 (96%)	11 (4%)	28	64
1	B	281/282 (100%)	271 (96%)	10 (4%)	31	66
1	C	281/282 (100%)	269 (96%)	12 (4%)	26	60
1	D	281/282 (100%)	270 (96%)	11 (4%)	28	64
All	All	1124/1128 (100%)	1080 (96%)	44 (4%)	28	64

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	24	THR
1	A	27	LYS
1	A	61	GLU
1	A	72	ARG
1	A	92	GLU
1	A	123	LEU
1	A	166	ASP
1	A	178	LEU
1	A	185	THR
1	A	201	ARG
1	B	11	GLN
1	B	24	THR
1	B	27	LYS
1	B	61	GLU
1	B	72	ARG
1	B	123	LEU
1	B	166	ASP
1	B	178	LEU
1	B	201	ARG
1	B	223	GLN
1	C	11	GLN
1	C	24	THR
1	C	27	LYS
1	C	37	TYR
1	C	61	GLU
1	C	72	ARG
1	C	92	GLU
1	C	113	LEU
1	C	123	LEU
1	C	166	ASP
1	C	201	ARG
1	C	223	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	11	GLN
1	D	24	THR
1	D	27	LYS
1	D	61	GLU
1	D	72	ARG
1	D	92	GLU
1	D	123	LEU
1	D	166	ASP
1	D	201	ARG
1	D	223	GLN
1	D	313	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	ASN
1	A	158	HIS
1	A	223	GLN
1	A	294	GLN
1	B	158	HIS
1	B	223	GLN
1	B	294	GLN
1	C	145	GLN
1	C	149	ASN
1	C	158	HIS
1	C	223	GLN
1	D	158	HIS
1	D	223	GLN
1	D	294	GLN
1	D	313	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	122	1	10,11,12	0.90	0	6,12,14	1.06	1 (16%)
1	KCX	C	122	1	10,11,12	0.91	0	6,12,14	1.10	1 (16%)
1	KCX	D	122	1	10,11,12	0.89	0	6,12,14	1.08	1 (16%)
1	LLP	C	33	1	23,24,25	2.64	4 (17%)	25,32,34	1.50	3 (12%)
1	LLP	A	33	1	23,24,25	2.64	4 (17%)	25,32,34	1.52	3 (12%)
1	LLP	D	33	1	23,24,25	2.63	4 (17%)	25,32,34	1.52	3 (12%)
1	LLP	B	33	1	23,24,25	2.66	4 (17%)	25,32,34	1.52	3 (12%)
1	KCX	B	122	1	10,11,12	0.94	0	6,12,14	1.10	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	122	1	-	1/9/10/12	-
1	KCX	C	122	1	-	1/9/10/12	-
1	KCX	D	122	1	-	1/9/10/12	-
1	LLP	C	33	1	-	4/16/17/19	0/1/1/1
1	LLP	A	33	1	-	5/16/17/19	0/1/1/1
1	LLP	D	33	1	-	4/16/17/19	0/1/1/1
1	LLP	B	33	1	-	5/16/17/19	0/1/1/1
1	KCX	B	122	1	-	0/9/10/12	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	33	LLP	C3-C2	8.05	1.49	1.41
1	B	33	LLP	C3-C2	8.03	1.49	1.41
1	C	33	LLP	C3-C2	7.98	1.49	1.41
1	D	33	LLP	C3-C2	7.91	1.49	1.41
1	D	33	LLP	O-C	6.32	1.44	1.20
1	B	33	LLP	O-C	6.32	1.44	1.20
1	C	33	LLP	O-C	6.27	1.43	1.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	33	LLP	O-C	6.27	1.43	1.20
1	B	33	LLP	C4-C5	5.41	1.49	1.42
1	A	33	LLP	C4-C5	5.34	1.49	1.42
1	C	33	LLP	C4-C5	5.28	1.49	1.42
1	D	33	LLP	C4-C5	5.25	1.49	1.42
1	C	33	LLP	C4-C3	4.73	1.48	1.41
1	B	33	LLP	C4-C3	4.65	1.48	1.41
1	D	33	LLP	C4-C3	4.61	1.48	1.41
1	A	33	LLP	C4-C3	4.58	1.48	1.41

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	33	LLP	C4-C3-C2	-5.13	117.26	120.14
1	C	33	LLP	C4-C3-C2	-5.11	117.26	120.14
1	D	33	LLP	C4-C3-C2	-4.82	117.43	120.14
1	A	33	LLP	C4-C3-C2	-4.64	117.53	120.14
1	A	33	LLP	CD-CE-NZ	-2.93	103.09	110.83
1	D	33	LLP	CD-CE-NZ	-2.70	103.69	110.83
1	B	33	LLP	CD-CE-NZ	-2.62	103.89	110.83
1	B	122	KCX	OQ1-CX-NZ	-2.53	121.08	124.92
1	C	122	KCX	OQ1-CX-NZ	-2.53	121.08	124.92
1	B	33	LLP	C6-N1-C2	2.49	123.71	119.20
1	D	122	KCX	OQ1-CX-NZ	-2.44	121.22	124.92
1	D	33	LLP	C6-N1-C2	2.42	123.59	119.20
1	C	33	LLP	CD-CE-NZ	-2.41	104.44	110.83
1	A	122	KCX	OQ1-CX-NZ	-2.39	121.30	124.92
1	A	33	LLP	C6-N1-C2	2.38	123.52	119.20
1	C	33	LLP	C6-N1-C2	2.36	123.48	119.20

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	33	LLP	CG-CD-CE-NZ
1	C	33	LLP	CG-CD-CE-NZ
1	D	33	LLP	C5-C4-C4'-NZ
1	D	33	LLP	CG-CD-CE-NZ
1	A	33	LLP	CG-CD-CE-NZ
1	D	33	LLP	C3-C4-C4'-NZ
1	B	33	LLP	C3-C4-C4'-NZ
1	A	33	LLP	C3-C4-C4'-NZ

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	C	33	LLP	C3-C4-C4'-NZ
1	B	33	LLP	CD-CE-NZ-C4'
1	D	33	LLP	CD-CE-NZ-C4'
1	B	33	LLP	C5-C4-C4'-NZ
1	A	33	LLP	C4-C5-C5'-OP4
1	C	33	LLP	CD-CE-NZ-C4'
1	B	33	LLP	C5'-OP4-P-OP1
1	A	33	LLP	CD-CE-NZ-C4'
1	A	122	KCX	CG-CD-CE-NZ
1	A	33	LLP	C5-C4-C4'-NZ
1	C	33	LLP	C5-C4-C4'-NZ
1	D	122	KCX	CG-CD-CE-NZ
1	C	122	KCX	CG-CD-CE-NZ

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	33	LLP	1	0
1	A	33	LLP	1	0
1	D	33	LLP	1	0
1	B	33	LLP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	C	3400	-	3,3,3	0.81	0	3,3,3	1.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	B	400	-	3,3,3	0.79	0	3,3,3	1.37	0
2	ACT	A	1400	-	3,3,3	0.80	0	3,3,3	1.33	0
2	ACT	D	2400	-	3,3,3	0.82	0	3,3,3	1.32	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3400	ACT	1	0
2	A	1400	ACT	1	0
2	D	2400	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/357 (99%)	0.01	5 (1%) 73 64	40, 49, 60, 64	0
1	B	354/357 (99%)	0.18	9 (2%) 58 48	43, 51, 63, 68	0
1	C	354/357 (99%)	0.06	3 (0%) 82 75	44, 54, 67, 70	0
1	D	354/357 (99%)	0.15	6 (1%) 69 60	44, 55, 69, 72	0
All	All	1416/1428 (99%)	0.10	23 (1%) 70 61	40, 52, 65, 72	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	356	GLY	7.2
1	D	313	GLN	3.6
1	A	286	VAL	3.5
1	D	355	SER	3.5
1	A	356	GLY	3.4
1	B	355	SER	3.4
1	B	166	ASP	3.2
1	B	286	VAL	3.1
1	D	286	VAL	3.1
1	C	286	VAL	3.0
1	B	11	GLN	2.8
1	D	148	ALA	2.8
1	C	355	SER	2.7
1	D	287	PHE	2.7
1	D	166	ASP	2.4
1	A	147	SER	2.4
1	B	287	PHE	2.4
1	B	24	THR	2.4
1	A	287	PHE	2.3
1	C	185	THR	2.3
1	A	25	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	93	HIS	2.0
1	B	43	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	D	122	12/13	0.90	0.11	57,57,57,58	0
1	KCX	A	122	12/13	0.92	0.10	44,46,47,47	0
1	LLP	D	33	24/25	0.93	0.10	45,45,48,48	0
1	KCX	B	122	12/13	0.94	0.10	49,50,50,50	0
1	KCX	C	122	12/13	0.95	0.08	54,55,55,55	0
1	LLP	B	33	24/25	0.96	0.08	39,40,44,45	0
1	LLP	A	33	24/25	0.96	0.08	34,35,39,39	0
1	LLP	C	33	24/25	0.96	0.08	42,43,45,45	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	C	3400	4/4	0.94	0.13	55,55,55,55	0
2	ACT	D	2400	4/4	0.94	0.10	52,52,52,52	0
2	ACT	A	1400	4/4	0.95	0.12	49,49,49,49	0
2	ACT	B	400	4/4	0.96	0.08	45,45,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.