



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:58 AM UTC

PDB ID : 2ODU / pdb_00002odu
Title : Crystal structure of a fragment of the plakin domain of plectin
Authors : de Pereda, J.M.
Deposited on : 2006-12-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	10% 70% 20% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plectin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	5	0
			1842	1162	340	332	8			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	296	GLY	-	cloning artifact	UNP Q6S383
A	297	SER	-	cloning artifact	UNP Q6S383
A	298	HIS	-	cloning artifact	UNP Q6S383
A	299	MET	-	cloning artifact	UNP Q6S383
A	506	LYS	GLU	engineered mutation	UNP Q6S383


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	16	Total	O	0	0
			16	16		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plectin 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	154.15Å 26.34Å 58.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.47 – 2.30 46.47 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.47-2.30) 99.8 (46.47-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.38 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.216 , 0.269 0.213 , 0.268	Depositor DCC
R_{free} test set	533 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1858	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	1/1874 (0.1%)	1.10	2/2517 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	424	ILE	C-O	-5.28	1.17	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	394	GLU	N-CA-C	5.91	117.39	111.07
1	A	348	PHE	N-CA-C	5.04	116.46	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1842	0	1853	40	0
2	A	16	0	0	1	0
All	All	1858	0	1853	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389[A]:HIS:CD2	1:A:391:LEU:H	1.72	1.07
1:A:389[A]:HIS:HD2	1:A:391:LEU:H	1.17	0.90
1:A:389[A]:HIS:CD2	1:A:391:LEU:HB3	2.10	0.86
1:A:389[A]:HIS:HD2	1:A:391:LEU:HB3	1.37	0.85
1:A:428:LEU:HD12	1:A:479:LEU:CD2	2.12	0.79
1:A:428:LEU:HD12	1:A:479:LEU:HD22	1.67	0.75
1:A:453:LEU:HD21	1:A:459:PRO:HD3	1.67	0.75
1:A:467:ARG:HD2	1:A:471:LYS:HE2	1.69	0.74
1:A:343:ILE:O	1:A:347:GLN:HG3	1.88	0.74
1:A:389[A]:HIS:HD2	1:A:391:LEU:N	1.87	0.73
1:A:304:LEU:HD21	1:A:382:LEU:HD21	1.74	0.70
1:A:389[A]:HIS:HD2	1:A:391:LEU:CB	2.05	0.69
1:A:319:LEU:O	1:A:323:ARG:HG3	1.95	0.66
1:A:389[A]:HIS:CD2	1:A:391:LEU:N	2.55	0.65
1:A:438:GLN:HG2	1:A:475:MET:HE1	1.81	0.63
1:A:472:ALA:HA	1:A:475:MET:HE3	1.83	0.60
1:A:304:LEU:C	1:A:304:LEU:HD23	2.28	0.59
1:A:364:ASN:HA	1:A:367:LYS:HD2	1.83	0.58
1:A:428:LEU:HD12	1:A:479:LEU:HD23	1.85	0.57
1:A:467:ARG:HH11	1:A:471:LYS:NZ	2.03	0.56
1:A:337:SER:HB2	2:A:14:HOH:O	2.05	0.55
1:A:467:ARG:CD	1:A:471:LYS:HE2	2.36	0.55
1:A:421:LEU:HD21	1:A:489:GLY:HA3	1.91	0.53
1:A:389[A]:HIS:CD2	1:A:391:LEU:CB	2.86	0.52
1:A:391:LEU:C	1:A:391:LEU:HD23	2.34	0.52
1:A:468:ASP:HA	1:A:471:LYS:HE3	1.92	0.52
1:A:318:LEU:HD22	1:A:322:MET:HE2	1.94	0.50
1:A:448:SER:HA	1:A:451:ARG:HG2	1.94	0.50
1:A:438:GLN:HG2	1:A:475:MET:CE	2.43	0.49
1:A:428:LEU:HD13	1:A:482:ASP:HB2	1.98	0.46
1:A:450:VAL:HG22	1:A:518:ARG:HG2	1.97	0.46
1:A:353:GLU:O	1:A:357:PRO:HG2	2.16	0.46
1:A:304:LEU:C	1:A:304:LEU:CD2	2.90	0.44
1:A:311:TYR:CZ	1:A:315:VAL:HG21	2.54	0.42
1:A:329:PHE:C	1:A:331:GLU:N	2.77	0.42
1:A:389[A]:HIS:HD2	1:A:391:LEU:CA	2.32	0.42
1:A:421:LEU:O	1:A:425:VAL:HG23	2.20	0.42
1:A:329:PHE:C	1:A:331:GLU:H	2.29	0.41
1:A:467:ARG:HH11	1:A:471:LYS:HZ3	1.66	0.40
1:A:304:LEU:HD23	1:A:304:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/235 (93%)	217 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/203 (96%)	179 (92%)	16 (8%)	10	14

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	303	GLU
1	A	304	LEU
1	A	312	ARG
1	A	318	LEU
1	A	331	GLU
1	A	350	LYS
1	A	374	GLU
1	A	383	LYS
1	A	384	VAL
1	A	391	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	407	ARG
1	A	422	GLN
1	A	458	VAL
1	A	490	ARG
1	A	496	GLN
1	A	512	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	GLN
1	A	401	HIS
1	A	493	GLN
1	A	496	GLN
1	A	516	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	217/235 (92%)	0.97	23 (10%)	11 12	17, 41, 50, 59	5 (2%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	494	GLY	4.6
1	A	334	PHE	4.2
1	A	386	PRO	4.1
1	A	377	VAL	3.5
1	A	306	LEU	3.3
1	A	497	MET	3.2
1	A	378	GLN	3.0
1	A	451	ARG	2.8
1	A	391	LEU	2.8
1	A	333	ARG	2.7
1	A	392	ASP	2.5
1	A	493	GLN	2.5
1	A	500	ARG	2.5
1	A	308	TRP	2.4
1	A	423[A]	ARG	2.4
1	A	498	TYR	2.4
1	A	436	GLU	2.4
1	A	455	ALA	2.3
1	A	501	VAL	2.3
1	A	387	GLY	2.2
1	A	373	LEU	2.2
1	A	379	ALA	2.2
1	A	336	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.