



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:13 AM UTC

PDB ID : 7ODC / pdb_00007odc
Title : CRYSTAL STRUCTURE ORNITHINE DECARBOXYLASE FROM
MOUSE, TRUNCATED 37 RESIDUES FROM THE C-TERMINUS, TO 1.6
ANGSTROM RESOLUTION
Authors : Kern, A.D.; Oliveira, M.A.; Coffino, P.; Hackert, M.L.
Deposited on : 1999-03-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	10% 77% 10% 9%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ORNITHINE DECARBOXYLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			3030	1944	507	558	21			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	421	Total	O	0	0
			421	421		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (ORNITHINE DECARBOXYLASE)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.50Å 74.00Å 45.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.90 – 1.60 27.90 – 1.60	Depositor EDS
% Data completeness (in resolution range)	91.3 (27.90-1.60) 91.3 (27.90-1.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.60Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.199 , 0.232 0.197 , 0.227	Depositor DCC
R_{free} test set	1995 reflections (4.06%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	1.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3466	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/3097	1.43	21/4189 (0.5%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	15	ASP	CA-CB-CG	-8.68	103.92	112.60
1	A	280	VAL	N-CA-C	7.20	120.74	113.47
1	A	9	PHE	CA-CB-CG	6.92	120.72	113.80
1	A	233	ASP	CA-CB-CG	6.85	119.45	112.60
1	A	206	ASP	CA-CB-CG	-6.34	106.26	112.60
1	A	402	ARG	NE-CZ-NH2	6.27	124.84	119.20
1	A	69	LYS	CB-CA-C	-6.17	99.22	110.63
1	A	183	ARG	NE-CZ-NH2	6.15	124.73	119.20
1	A	61	ARG	CD-NE-CZ	5.97	132.76	124.40
1	A	320	ASP	CA-CB-CG	5.62	118.22	112.60
1	A	335	HIS	CA-CB-CG	-5.61	108.19	113.80
1	A	69	LYS	N-CA-CB	5.56	118.79	110.22
1	A	15	ASP	N-CA-C	5.53	122.58	110.80
1	A	19	THR	N-CA-CB	-5.52	102.47	110.70
1	A	4	PHE	CA-CB-CG	5.47	119.27	113.80
1	A	348	GLU	CB-CA-C	5.29	119.20	109.46
1	A	87	PHE	CA-CB-CG	-5.29	108.52	113.80
1	A	81	ALA	O-C-N	5.28	127.50	122.07
1	A	145	ALA	N-CA-C	5.23	119.81	113.38
1	A	270	ARG	NE-CZ-NH2	-5.17	114.55	119.20
1	A	270	ARG	CD-NE-CZ	5.01	131.42	124.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3030	0	3007	43	0
2	A	15	0	6	1	0
3	A	421	0	0	13	0
All	All	3466	0	3013	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:340:LEU:HD23	1:A:342:LYS:HE2	1.62	0.79
1:A:48:ILE:HD11	1:A:76:ILE:HD12	1.71	0.71
1:A:57:LYS:O	1:A:57:LYS:HE2	1.94	0.68
1:A:183:ARG:HD3	3:A:756:HOH:O	1.96	0.65
1:A:340:LEU:CD2	1:A:342:LYS:HE2	2.29	0.63
1:A:413:MET:HA	1:A:413:MET:HE2	1.80	0.62
1:A:19:THR:HG21	1:A:21:LYS:NZ	2.14	0.62
1:A:205:PRO:HG3	1:A:246:LEU:HD11	1.85	0.58
1:A:19:THR:HG23	1:A:21:LYS:H	1.69	0.57
1:A:15:ASP:O	1:A:16:GLU:O	2.21	0.57
1:A:402:ARG:HD3	3:A:916:HOH:O	2.05	0.57
1:A:250:GLU:HG3	3:A:844:HOH:O	2.05	0.56
1:A:411:ARG:HB3	1:A:412:PRO:HD3	1.88	0.55
1:A:231:LEU:HD23	1:A:270:ARG:HD3	1.89	0.55
1:A:230:HIS:HD2	3:A:646:HOH:O	1.89	0.55
1:A:148:LYS:HA	1:A:148:LYS:HE2	1.88	0.55
1:A:49:LEU:HD11	1:A:416:LEU:HD12	1.89	0.54
1:A:340:LEU:HD11	1:A:372:LEU:HD22	1.90	0.53
1:A:343:ARG:HG3	1:A:343:ARG:HH11	1.76	0.50
1:A:297:TRP:HA	1:A:312:GLN:HB2	1.94	0.48
1:A:2:SER:HA	3:A:796:HOH:O	2.14	0.47
1:A:369:ARG:HB2	1:A:369:ARG:NH1	2.30	0.47
1:A:247:LYS:HG3	3:A:772:HOH:O	2.13	0.47
1:A:342:LYS:HD2	1:A:343:ARG:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:GLU:OE1	1:A:243:ASP:N	2.49	0.45
1:A:19:THR:HG22	1:A:22:ASP:H	1.82	0.45
1:A:274:GLU:O	2:A:425:PLP:H6	2.17	0.44
1:A:22:ASP:O	1:A:26:GLN:HG3	2.18	0.44
1:A:342:LYS:O	1:A:342:LYS:HE3	2.17	0.44
1:A:69:LYS:HE3	3:A:688:HOH:O	2.17	0.44
1:A:21:LYS:HG3	3:A:790:HOH:O	2.17	0.44
1:A:136:GLU:HG2	3:A:774:HOH:O	2.18	0.43
1:A:203:THR:HG22	3:A:838:HOH:O	2.19	0.42
1:A:402:ARG:NH1	3:A:897:HOH:O	2.52	0.42
1:A:180:LEU:HD23	1:A:183:ARG:HH21	1.85	0.42
1:A:230:HIS:HE1	3:A:664:HOH:O	2.03	0.42
1:A:295:THR:CG2	1:A:311:GLU:N	2.83	0.42
1:A:204:ASP:HA	1:A:205:PRO:HD3	1.91	0.41
1:A:180:LEU:CD2	1:A:183:ARG:HH21	2.33	0.41
1:A:15:ASP:HB2	3:A:792:HOH:O	2.21	0.41
1:A:19:THR:HG23	1:A:21:LYS:N	2.33	0.41
1:A:231:LEU:CD2	1:A:270:ARG:HD3	2.51	0.41
1:A:19:THR:HG21	1:A:21:LYS:HZ2	1.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	379/424 (89%)	369 (97%)	9 (2%)	1 (0%)	36 20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	326/362 (90%)	304 (93%)	22 (7%)	15 3

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	THR
1	A	21	LYS
1	A	49	LEU
1	A	57	LYS
1	A	69	LYS
1	A	183	ARG
1	A	202	CYS
1	A	203	THR
1	A	206	ASP
1	A	241	SER
1	A	242	GLU
1	A	244	THR
1	A	247	LYS
1	A	266	ASP
1	A	311	GLU
1	A	312	GLN
1	A	342	LYS
1	A	343	ARG
1	A	345	LYS
1	A	347	ASP
1	A	363	LEU
1	A	398	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	116	GLN
1	A	129	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	GLN
1	A	230	HIS
1	A	371	ASN
1	A	376	HIS
1	A	385	ASN
1	A	415	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	425	1	15,15,16	1.25	2 (13%)	21,22,23	1.69	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	425	1	-	0/6/6/8	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	425	PLP	C4A-C4	-2.21	1.47	1.51
2	A	425	PLP	P-O3P	-2.13	1.46	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	425	PLP	C6-C5-C4	4.85	122.07	118.10
2	A	425	PLP	C5-C6-N1	-2.68	119.47	123.83
2	A	425	PLP	C5A-C5-C4	-2.53	117.64	122.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	425	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/424 (91%)	0.48	44 (11%) 10 9	11, 18, 39, 50	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	TRP	5.1
1	A	266	ASP	4.8
1	A	4	PHE	4.6
1	A	204	ASP	4.2
1	A	5	THR	4.2
1	A	243	ASP	4.2
1	A	203	THR	3.8
1	A	202	CYS	3.7
1	A	414	TRP	3.5
1	A	18	PHE	3.2
1	A	360	CYS	3.2
1	A	16	GLU	3.1
1	A	417	MET	3.0
1	A	29	ASN	2.9
1	A	415	GLN	2.9
1	A	7	ASP	2.9
1	A	250	GLU	2.8
1	A	418	LYS	2.8
1	A	206	ASP	2.8
1	A	416	LEU	2.7
1	A	157	THR	2.7
1	A	311	GLU	2.6
1	A	174	LEU	2.6
1	A	339	LEU	2.6
1	A	217	CYS	2.5
1	A	6	LYS	2.5
1	A	15	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	3	SER	2.5
1	A	183	ARG	2.4
1	A	245	LYS	2.4
1	A	242	GLU	2.3
1	A	26	GLN	2.3
1	A	209	VAL	2.3
1	A	201	GLY	2.3
1	A	369	ARG	2.2
1	A	246	LEU	2.2
1	A	244	THR	2.2
1	A	342	LYS	2.2
1	A	144	ARG	2.2
1	A	335	HIS	2.2
1	A	102	GLY	2.1
1	A	205	PRO	2.1
1	A	9	PHE	2.1
1	A	247	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLP	A	425	15/16	0.96	0.09	13,18,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.