



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:52 PM UTC

PDB ID : 7ODD / pdb_00007odd
Title : Crystal structure of bovine Hsc70(aa1-554)E213A/D214A in complex with
tricine
Authors : Zehe, M.; Grimm, C.; Sotriffer, C.
Deposited on : 2021-04-29
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	554	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock cognate 71 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	539	4161	2605	725	819	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	ALA	GLU	engineered mutation	UNP P19120
A	214	ALA	ASP	engineered mutation	UNP P19120

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0

- Molecule 3 is N-[1,3-dihydroxy-2-(hydroxymethyl)propan-2-yl]glycine (CCD ID: VLS) (formula: C₆H₁₃NO₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	24	12	2	10	0	1

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	15	8	2	4	1	0	0

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	K		
5	A	2	2	2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	146	146	146	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Heat shock cognate 71 kDa protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.06Å 49.07Å 87.27Å 90.00° 99.69° 90.00°	Depositor
Resolution (Å)	42.63 – 1.98 42.63 – 1.98	Depositor EDS
% Data completeness (in resolution range)	98.1 (42.63-1.98) 98.2 (42.63-1.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.98Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.225 , 0.245 0.228 , 0.248	Depositor DCC
R_{free} test set	1893 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	43.6	Xtrriage
Anisotropy	0.413	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4366	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VLS, GOL, EPE, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/4221	1.19	1/5701 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	481	ASP	CA-C-O	-6.31	114.68	121.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4161	0	4183	22	0
2	A	18	0	24	1	0
3	A	24	0	0	0	0
4	A	15	0	17	1	0
5	A	2	0	0	0	0
6	A	146	0	0	2	0
All	All	4366	0	4224	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:THR:HG22	1:A:418:THR:H	1.63	0.64
1:A:72:ARG:HA	2:A:603:GOL:H2	1.87	0.57
1:A:176:PRO:HB2	1:A:210:LEU:HD11	1.88	0.56
1:A:475:GLU:HB2	1:A:493:LYS:HD3	1.87	0.56
1:A:397:THR:CG2	1:A:418:THR:H	2.21	0.53
1:A:404:GLU:HB2	1:A:450:THR:HG21	1.90	0.52
1:A:191:ALA:HA	1:A:417:ASN:HD21	1.76	0.50
1:A:237:MET:HE1	1:A:271:LYS:HB2	1.94	0.48
1:A:342:ARG:NH1	6:A:702:HOH:O	2.22	0.48
1:A:43:ALA:HB3	1:A:50:LEU:HB2	1.95	0.48
1:A:552:THR:HG22	1:A:553:VAL:HG23	1.96	0.47
1:A:175:GLU:N	1:A:176:PRO:HD2	2.30	0.46
4:A:605:EPE:H31	4:A:605:EPE:H82	1.57	0.46
1:A:28:ILE:HG13	1:A:367:GLU:HB3	1.97	0.45
1:A:395:ASP:O	1:A:396:VAL:HG22	2.17	0.45
1:A:539:LYS:HA	1:A:539:LYS:HD2	1.77	0.44
1:A:103:VAL:O	1:A:113:SER:HA	2.18	0.43
1:A:10:ASP:OD2	6:A:701:HOH:O	2.22	0.42
1:A:336:LEU:HD12	1:A:361:LYS:HB3	2.01	0.41
1:A:213:ALA:HB1	1:A:417:ASN:CG	2.45	0.41
1:A:42:VAL:HG11	1:A:122:MET:SD	2.61	0.41
1:A:481:ASP:OD2	1:A:485:ILE:HB	2.21	0.40
1:A:415:LYS:HG2	1:A:416:ARG:N	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	535/554 (97%)	530 (99%)	4 (1%)	1 (0%)	43 35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	396	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	451/466 (97%)	444 (98%)	7 (2%)	55 50

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	THR
1	A	235	ASN
1	A	396	VAL
1	A	480	ILE
1	A	507	LYS
1	A	520	GLN
1	A	553	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	A	33	GLN
1	A	249	HIS
1	A	417	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	601	-	5,5,5	0.10	0	5,5,5	0.24	0
2	GOL	A	602	-	5,5,5	0.11	0	5,5,5	0.24	0
3	VLS	A	604[A]	-	11,11,11	0.82	0	13,14,14	1.43	2 (15%)
2	GOL	A	603	-	5,5,5	0.11	0	5,5,5	0.32	0
3	VLS	A	604[B]	-	11,11,11	0.82	0	13,14,14	1.57	3 (23%)
4	EPE	A	605	-	15,15,15	2.09	1 (6%)	19,20,20	1.14	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	0/4/4/4	-
2	GOL	A	602	-	-	2/4/4/4	-
3	VLS	A	604[A]	-	-	6/15/15/15	-
2	GOL	A	603	-	-	1/4/4/4	-
3	VLS	A	604[B]	-	-	8/15/15/15	-
4	EPE	A	605	-	-	4/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	605	EPE	C10-S	-7.93	1.66	1.77

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	604[B]	VLS	O07-C06-C03	-3.05	105.48	111.68
3	A	604[A]	VLS	O07-C06-C03	-2.91	105.77	111.68
3	A	604[B]	VLS	O05-C04-C03	-2.83	105.92	111.68
3	A	604[A]	VLS	O05-C04-C03	-2.45	106.70	111.68
4	A	605	EPE	O2S-S-C10	2.29	110.19	106.73
3	A	604[B]	VLS	C06-C03-C04	-2.26	105.07	110.02

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	602	GOL	O1-C1-C2-C3
3	A	604[B]	VLS	C02-C03-C04-O05
3	A	604[B]	VLS	C06-C03-C04-O05
3	A	604[B]	VLS	N08-C03-C04-O05
3	A	604[B]	VLS	C04-C03-C06-O07
4	A	605	EPE	C8-C7-N4-C3
2	A	603	GOL	C1-C2-C3-O3
4	A	605	EPE	N4-C7-C8-O8
2	A	602	GOL	O1-C1-C2-O2
3	A	604[A]	VLS	C04-C03-C06-O07
3	A	604[A]	VLS	C02-C03-N08-C09
3	A	604[B]	VLS	C02-C03-C06-O07
4	A	605	EPE	C8-C7-N4-C5
3	A	604[A]	VLS	C02-C03-C06-O07
3	A	604[A]	VLS	C10-C09-N08-C03
3	A	604[B]	VLS	C10-C09-N08-C03
4	A	605	EPE	S-C10-C9-N1
3	A	604[B]	VLS	C02-C03-N08-C09
3	A	604[A]	VLS	N08-C03-C06-O07
3	A	604[B]	VLS	N08-C03-C06-O07
3	A	604[A]	VLS	C06-C03-C04-O05


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	GOL	1	0
4	A	605	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	539/554 (97%)	0.69	44 (8%) 17 24	34, 56, 100, 132	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	GLU	4.3
1	A	189	VAL	4.2
1	A	191	ALA	3.7
1	A	536	VAL	3.5
1	A	553	VAL	3.4
1	A	396	VAL	3.4
1	A	552	THR	3.4
1	A	89	HIS	3.3
1	A	95	VAL	3.2
1	A	190	GLY	2.9
1	A	540	ASN	2.8
1	A	94	VAL	2.8
1	A	105	VAL	2.8
1	A	60	ALA	2.7
1	A	98	ALA	2.7
1	A	510	LEU	2.7
1	A	395	ASP	2.6
1	A	539	LYS	2.6
1	A	210	LEU	2.6
1	A	79	ASP	2.5
1	A	518	MET	2.5
1	A	75	GLY	2.5
1	A	92	PHE	2.4
1	A	138	THR	2.4
1	A	5	PRO	2.4
1	A	61	MET	2.4
1	A	114	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	482	ALA	2.3
1	A	115	TYR	2.3
1	A	254	SER	2.3
1	A	463	GLY	2.2
1	A	538	SER	2.2
1	A	551	ALA	2.2
1	A	88	LYS	2.2
1	A	417	ASN	2.2
1	A	483	ASN	2.2
1	A	139	VAL	2.2
1	A	480	ILE	2.1
1	A	62	ASN	2.1
1	A	248	LYS	2.1
1	A	481	ASP	2.1
1	A	170	LEU	2.0
1	A	99	GLY	2.0
1	A	247	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	VLS	A	604[A]	12/12	0.67	0.18	57,78,78,80	12
3	VLS	A	604[B]	12/12	0.67	0.18	61,78,79,79	12
2	GOL	A	603	6/6	0.84	0.17	68,85,93,96	0
2	GOL	A	602	6/6	0.85	0.14	58,68,71,76	0
5	K	A	606	1/1	0.86	0.27	135,135,135,135	0
5	K	A	607	1/1	0.86	0.21	132,132,132,132	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	601	6/6	0.89	0.11	75,79,86,95	0
4	EPE	A	605	15/15	0.91	0.12	51,66,76,77	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.