



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:28 AM UTC

PDB ID : 7ODM / pdb_00007odm
Title : AmGSTF1 Y118S variant
Authors : Pohl, E.; Eno, R.F.M.
Deposited on : 2021-04-30
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	219	5% 73% 23% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3424 atoms, of which 1681 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutathione transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	215	3362	1096	1665	281	313	7	52	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	118	SER	TYR	engineered mutation	UNP Q9ZS17

- Molecule 2 is GLUTATHIONE (CCD ID: GSH) (formula: C₁₀H₁₇N₃O₆S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	S		
2	AAA	1	36	10	16	3	6	1	1	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	26	Total	O	0	0
			26	26		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutathione transferase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	112.61Å 112.61Å 103.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.34 – 2.60 45.34 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.34-2.60) 100.0 (45.34-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.188 , 0.257 0.185 , 0.254	Depositor DCC
R_{free} test set	549 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å ²)	66.9	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 63.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3424	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GSH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.29	3/1744 (0.2%)	1.67	19/2378 (0.8%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	171	ASN	C-O	6.59	1.32	1.24
1	AAA	162	GLY	C-O	5.20	1.29	1.23
1	AAA	158	SER	C-O	5.01	1.30	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	19	THR	CA-CB-OG1	7.88	121.42	109.60
1	AAA	17	ARG	CA-C-N	7.63	130.18	120.56
1	AAA	17	ARG	C-N-CA	7.63	130.18	120.56
1	AAA	151	GLU	CB-CG-CD	6.44	123.54	112.60
1	AAA	151	GLU	CB-CA-C	-6.14	100.98	110.81
1	AAA	62	ASP	CA-CB-CG	6.05	118.65	112.60
1	AAA	45	GLU	N-CA-C	-5.94	104.72	111.07
1	AAA	24	GLU	CB-CG-CD	5.84	122.53	112.60
1	AAA	152	ALA	N-CA-C	-5.79	104.66	110.97
1	AAA	122	PHE	CA-CB-CG	5.72	119.52	113.80
1	AAA	25	VAL	N-CA-C	-5.49	107.45	113.43
1	AAA	25	VAL	N-CA-CB	-5.26	106.42	112.21
1	AAA	146	VAL	N-CA-C	-5.16	105.51	110.72
1	AAA	102	THR	CA-C-N	5.14	127.16	120.28
1	AAA	102	THR	C-N-CA	5.14	127.16	120.28
1	AAA	25	VAL	CB-CA-C	5.09	117.20	110.84
1	AAA	18	VAL	CA-C-O	-5.09	115.78	121.17
1	AAA	71	ILE	CA-C-N	5.07	127.03	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	71	ILE	C-N-CA	5.07	127.03	120.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1697	1665	1643	26	0
2	AAA	20	16	15	2	0
3	AAA	26	0	0	1	0
All	All	1743	1681	1658	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:180:MET:HE1	1:AAA:188:PHE:CD2	2.30	0.67
1:AAA:88:ARG:NH1	1:AAA:94:GLU:OE2	2.30	0.64
1:AAA:180:MET:HE1	1:AAA:188:PHE:CG	2.34	0.62
1:AAA:180:MET:CE	1:AAA:188:PHE:CD2	2.85	0.59
1:AAA:105:ASP:O	1:AAA:110:ASN:HB2	2.04	0.57
1:AAA:14:ASN:OD1	1:AAA:17:ARG:NH1	2.38	0.57
1:AAA:22:LEU:HD13	1:AAA:29:TYR:CG	2.41	0.55
1:AAA:182:THR:HB	1:AAA:183:PRO:HD2	1.88	0.55
1:AAA:20:LEU:HA	1:AAA:210:ILE:HG13	1.91	0.53
1:AAA:114:SER:OG	1:AAA:115:PRO:HD3	2.12	0.50
1:AAA:5:LYS:O	1:AAA:58:PHE:HA	2.13	0.49
1:AAA:182:THR:HB	1:AAA:183:PRO:CD	2.43	0.48
1:AAA:54:GLN:HG2	2:AAA:301:GSH:O32	2.14	0.48
1:AAA:105:ASP:HB2	1:AAA:172:HIS:CE1	2.49	0.48
1:AAA:73:LYS:O	1:AAA:77:ARG:HG3	2.15	0.46
1:AAA:59:GLN:HB3	1:AAA:64:LEU:HD23	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:142:LYS:HE3	1:AAA:142:LYS:HB2	1.83	0.45
1:AAA:17:ARG:NH2	3:AAA:405:HOH:O	2.51	0.44
1:AAA:118:SER:HA	1:AAA:122:PHE:HB2	2.00	0.44
1:AAA:60:ASP:OD2	1:AAA:78:LYS:NZ	2.38	0.43
1:AAA:64:LEU:HD23	1:AAA:64:LEU:HA	1.82	0.43
1:AAA:59:GLN:HB3	1:AAA:64:LEU:CD2	2.48	0.43
1:AAA:115:PRO:O	1:AAA:119:GLN:HB2	2.20	0.42
1:AAA:34:ILE:CD1	1:AAA:46:HIS:ND1	2.84	0.41
1:AAA:180:MET:HA	1:AAA:185:ALA:HB2	2.02	0.41
1:AAA:55:ILE:HG13	2:AAA:301:GSH:HB22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	213/219 (97%)	191 (90%)	20 (9%)	2 (1%)	14	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	119	GLN
1	AAA	123	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	179/189 (95%)	170 (95%)	9 (5%)	22	46

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	30	GLU
1	AAA	37	ASN
1	AAA	103	GLU
1	AAA	134	LYS
1	AAA	139	SER
1	AAA	141	GLU
1	AAA	151	GLU
1	AAA	158	SER
1	AAA	195	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GSH	AAA	301	-	18,19,19	0.76	0	21,24,24	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSH	AAA	301	-	-	11/24/24/24	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	301	GSH	N1-CA1-CB1-CG1
2	AAA	301	GSH	C2-CA2-N2-CD1
2	AAA	301	GSH	OE1-CD1-N2-CA2
2	AAA	301	GSH	N2-CA2-CB2-SG2
2	AAA	301	GSH	C1-CA1-CB1-CG1
2	AAA	301	GSH	O31-C3-CA3-N3
2	AAA	301	GSH	O32-C3-CA3-N3
2	AAA	301	GSH	CG1-CD1-N2-CA2
2	AAA	301	GSH	N3-C2-CA2-N2
2	AAA	301	GSH	CB2-CA2-N2-CD1
2	AAA	301	GSH	O2-C2-CA2-N2


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	301	GSH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	215/219 (98%)	-0.12	11 (5%) 33 28	53, 76, 134, 171	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	122	PHE	4.3
1	AAA	127	ARG	3.8
1	AAA	126	MET	3.5
1	AAA	125	MET	3.3
1	AAA	216	PRO	2.8
1	AAA	2	ALA	2.8
1	AAA	120	CYS	2.5
1	AAA	36	PHE	2.4
1	AAA	130	PRO	2.3
1	AAA	121	LEU	2.1
1	AAA	129	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GSH	AAA	301	20/20	0.93	0.13	36,69,81,95	36

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.