



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 11:32 PM UTC

PDB ID : 1OE5 / pdb_00001oe5
Title : Xenopus SMUG1, an anti-mutator uracil-DNA Glycosylase
Authors : Wibley, J.E.A.; Pearl, L.H.
Deposited on : 2003-03-19
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	% 67% 26% • •
1	B	247	2% 63% 29% 5% • •
2	E	12	25% 50% 25%
3	F	13	23% 54% 23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	URA	A	1282	-	X	-	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SINGLE-STRAND SELECTIVE MONOFUNCTIONAL URACIL DNA GLYCOSYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	2	1
			1934	1241	338	342	13			
1	B	245	Total	C	N	O	S	0	0	0
			1939	1244	339	344	12			


- Molecule 2 is a DNA chain called 5'-D(*CP*CP*CP*GP*TP*GP*AP*GP*TP*CP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	12	Total	C	N	O	P	0	0	0
			241	115	44	71	11			

- Molecule 3 is a DNA chain called 5'-D(*CP*3DRP*GP*GP*AP*CP*TP*3DRP*AP*CP*GP*GP*GP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	13	Total	C	N	O	P	0	0	0
			248	117	46	73	12			

- Molecule 4 is URACIL (CCD ID: URA) (formula: C₄H₄N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	3	1		
6	B	1	Total	C	O	0	0
			4	3	1		

- Molecule 7 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 8 is 2'-DEOXYURIDINE (CCD ID: DUR) (formula: $C_9H_{12}N_2O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	F	1	Total	C	N	O	0	0
			16	9	2	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	123	Total	O	0	0
			123	123		
9	B	82	Total	O	0	0
			82	82		
9	E	12	Total	O	0	0
			12	12		
9	F	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SINGLE-STRAND SELECTIVE MONOFUNCTIONAL URACIL DNA GLYCOSYLASE




• Molecule 1: SINGLE-STRAND SELECTIVE MONOFUNCTIONAL URACIL DNA GLYCOSYLASE



• Molecule 2: 5'-D(*CP*CP*CP*GP*TP*GP*AP*GP*TP*CP*CP*G)-3'



• Molecule 3: 5'-D(*CP*3DRP*GP*GP*AP*CP*TP*3DRP*AP*CP*GP*GP*GP)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.60Å 85.65Å 78.47Å 90.00° 118.56° 90.00°	Depositor
Resolution (Å)	69.01 – 2.30 68.92 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.8 (69.01-2.30) 95.8 (68.92-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.97 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.163 , 0.229 0.164 , 0.224	Depositor DCC
R_{free} test set	1512 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4657	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, 3DR, DUR, URA, GOL, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.78	27/1993 (1.4%)	1.51	12/2701 (0.4%)
1	B	1.69	21/1990 (1.1%)	1.43	14/2698 (0.5%)
2	E	1.25	2/269 (0.7%)	2.31	16/413 (3.9%)
3	F	1.18	0/252	2.06	12/383 (3.1%)
All	All	1.68	50/4504 (1.1%)	1.58	54/6195 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	212	ALA	CA-CB	10.21	1.69	1.53
1	B	249	MET	SD-CE	9.35	2.02	1.79
1	B	237	MET	SD-CE	9.11	2.02	1.79
1	B	212	ALA	CA-CB	8.68	1.67	1.53
1	B	67	ASN	N-CA	-8.41	1.38	1.46

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	291	DC	C5'-C4'-C3'	-13.50	94.65	114.90
1	A	215	VAL	N-CA-C	8.83	119.63	110.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	402	DG	P-O3'-C3'	7.98	132.17	120.20
1	A	259	ASN	N-CA-C	7.53	122.31	113.20
3	F	400	DA	P-O3'-C3'	7.08	130.83	120.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	259	ASN	Peptide
1	A	59	GLN	Peptide
1	B	59	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1934	0	1940	40	0
1	B	1939	0	1942	48	0
2	E	241	0	136	5	0
3	F	248	0	140	6	0
4	A	8	0	3	1	0
5	A	6	0	8	2	0
6	A	4	0	8	0	0
6	B	4	0	8	0	0
7	A	15	0	18	0	0
7	B	15	0	18	0	0
8	F	16	0	12	4	0
9	A	123	0	0	8	0
9	B	82	0	0	12	0
9	E	12	0	0	1	0
9	F	10	0	0	2	0
All	All	4657	0	4233	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ASP:HB3	1:A:61:PRO:CD	1.18	1.48
1:B:249:MET:CE	1:B:249:MET:SD	2.02	1.47
1:B:237:MET:SD	1:B:237:MET:CE	2.02	1.46
1:A:60:ASP:CB	1:A:61:PRO:CD	2.13	1.26
1:A:77:GLU:HG2	9:A:2019:HOH:O	1.33	1.20

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/247 (99%)	235 (96%)	7 (3%)	2 (1%)	16	20
1	B	243/247 (98%)	227 (93%)	12 (5%)	4 (2%)	7	7
All	All	487/494 (99%)	462 (95%)	19 (4%)	6 (1%)	10	12

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	ASP
1	B	60	ASP
1	B	196	ALA
1	A	277	SER
1	B	61	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/215 (100%)	193 (90%)	21 (10%)	7	10
1	B	214/215 (100%)	188 (88%)	26 (12%)	5	5
All	All	428/430 (100%)	381 (89%)	47 (11%)	6	7

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	60	ASP
1	B	155	LEU
1	B	77	GLU
1	B	89	GLU
1	B	179	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	78	ASN
1	B	96	ASN
1	B	257	GLN
1	B	169	HIS
1	A	173	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	3DR	F	393	3	8,11,12	0.76	0	7,14,17	0.76	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	3DR	F	399	3	8,11,12	1.71	2 (25%)	7,14,17	2.33	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3DR	F	393	3	-	2/3/15/16	0/1/1/1
3	3DR	F	399	3	-	1/3/15/16	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	399	3DR	O3'-C3'	4.09	1.51	1.43
3	F	399	3DR	O4'-C4'	2.23	1.48	1.44

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	399	3DR	C1'-C2'-C3'	-4.81	98.10	103.26
3	F	399	3DR	O3'-C3'-C4'	2.77	120.56	110.07
3	F	399	3DR	O3'-C3'-C2'	2.02	116.18	111.43

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	393	3DR	C3'-C4'-C5'-O5'
3	F	393	3DR	O4'-C4'-C5'-O5'
3	F	399	3DR	C4'-C5'-O5'-P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	393	3DR	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	DUR	F	1405	-	17,17,17	0.66	0	24,24,24	0.94	1 (4%)
7	EPE	A	1285	-	15,15,15	1.54	1 (6%)	19,20,20	2.75	11 (57%)
7	EPE	B	1282	-	15,15,15	1.77	1 (6%)	19,20,20	2.59	10 (52%)
6	IPA	A	1284	-	3,3,3	0.81	0	3,3,3	0.41	0
5	GOL	A	1283	-	5,5,5	0.57	0	5,5,5	1.05	0
4	URA	A	1282	-	8,8,8	4.74	5 (62%)	10,10,10	3.42	5 (50%)
6	IPA	B	1281	-	3,3,3	1.06	0	3,3,3	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	DUR	F	1405	-	-	2/6/18/18	0/2/2/2
7	EPE	A	1285	-	-	4/9/19/19	0/1/1/1
7	EPE	B	1282	-	-	5/9/19/19	0/1/1/1
5	GOL	A	1283	-	-	4/4/4/4	-
4	URA	A	1282	-	-	-	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1282	URA	C4-N3	6.87	1.50	1.38
4	A	1282	URA	C6-N1	6.20	1.45	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1282	URA	C2-N1	6.00	1.44	1.36
7	B	1282	EPE	C10-S	5.92	1.85	1.77
4	A	1282	URA	C2-N3	5.33	1.46	1.37

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1285	EPE	C5-N4-C3	6.13	122.04	108.84
4	A	1282	URA	O2-C2-N1	-5.90	116.70	122.79
4	A	1282	URA	O4-C4-N3	5.75	127.61	119.27
7	B	1282	EPE	C5-N4-C3	5.42	120.51	108.84
7	B	1282	EPE	O2S-S-C10	5.24	114.64	106.73

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1283	GOL	O1-C1-C2-C3
7	A	1285	EPE	C10-C9-N1-C6
7	B	1282	EPE	C9-C10-S-O1S
7	B	1282	EPE	C9-C10-S-O2S
7	B	1282	EPE	C9-C10-S-O3S

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	F	1405	DUR	4	0
5	A	1283	GOL	2	0
4	A	1282	URA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/247 (98%)	-0.54	3 (1%) 76 77	13, 29, 47, 58	2 (0%)
1	B	245/247 (99%)	-0.13	4 (1%) 70 72	19, 37, 62, 74	2 (0%)
2	E	12/12 (100%)	0.11	0 100 100	30, 55, 70, 71	0
3	F	11/13 (84%)	0.24	0 100 100	36, 52, 83, 84	0
All	All	512/519 (98%)	-0.31	7 (1%) 73 75	13, 34, 61, 84	4 (0%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	SER	3.1
1	B	60	ASP	3.1
1	B	237	MET	2.5
1	A	60	ASP	2.4
1	B	187	ASN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	3DR	F	393	11/12	0.79	0.19	66,83,93,93	0
3	3DR	F	399	11/12	0.94	0.10	55,63,66,67	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	1283	6/6	0.85	0.21	77,83,84,84	0
7	EPE	B	1282	15/15	0.87	0.19	46,86,97,97	0
8	DUR	F	1405	16/16	0.88	0.39	73,76,81,81	16
4	URA	A	1282	8/8	0.93	0.15	23,36,42,45	0
7	EPE	A	1285	15/15	0.95	0.14	37,69,74,74	0
6	IPA	B	1281	4/4	0.96	0.07	18,19,24,26	0
6	IPA	A	1284	4/4	0.98	0.08	20,21,28,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.