



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:18 AM UTC

PDB ID : 2OEG / pdb_00002oeg
Title : Open and Closed Structures of the UDP-Glucose Pyrophosphorylase from Leishmania major
Authors : Steiner, T.
Deposited on : 2006-12-29
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UTP-glucose-1-phosphate uridylyltransferase 2, putative.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	482	Total	C	N	O	S	Se	0	0	0
			3721	2345	634	716	10	16			

There are 29 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q4QDU3
A	5	MSE	MET	modified residue	UNP Q4QDU3
A	19	MSE	MET	modified residue	UNP Q4QDU3
A	38	MSE	MET	modified residue	UNP Q4QDU3
A	53	MSE	MET	modified residue	UNP Q4QDU3
A	89	MSE	MET	modified residue	UNP Q4QDU3
A	128	MSE	MET	modified residue	UNP Q4QDU3
A	130	MSE	MET	modified residue	UNP Q4QDU3
A	161	MSE	MET	modified residue	UNP Q4QDU3
A	215	MSE	MET	modified residue	UNP Q4QDU3
A	235	MSE	MET	modified residue	UNP Q4QDU3
A	244	MSE	MET	modified residue	UNP Q4QDU3
A	293	MSE	MET	modified residue	UNP Q4QDU3
A	321	MSE	MET	modified residue	UNP Q4QDU3
A	356	MSE	MET	modified residue	UNP Q4QDU3
A	362	MSE	MET	modified residue	UNP Q4QDU3
A	423	MSE	MET	modified residue	UNP Q4QDU3
A	424	MSE	MET	modified residue	UNP Q4QDU3
A	495	MSE	-	cloning artifact	UNP Q4QDU3
A	496	ARG	-	cloning artifact	UNP Q4QDU3
A	497	PRO	-	cloning artifact	UNP Q4QDU3
A	498	LEU	-	cloning artifact	UNP Q4QDU3
A	499	GLU	-	cloning artifact	UNP Q4QDU3
A	500	HIS	-	cloning artifact	UNP Q4QDU3
A	501	HIS	-	cloning artifact	UNP Q4QDU3
A	502	HIS	-	cloning artifact	UNP Q4QDU3
A	503	HIS	-	cloning artifact	UNP Q4QDU3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	504	HIS	-	cloning artifact	UNP Q4QDU3
A	505	HIS	-	cloning artifact	UNP Q4QDU3

- Molecule 2 is URIDINE-5'-DIPHOSPHATE-GLUCOSE (CCD ID: UPG) (formula: $C_{15}H_{24}N_2O_{17}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			36	15	2	17	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	225	Total	O	0	0
			225	225		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UTP-glucose-1-phosphate uridylyltransferase 2, putative



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	80.25Å 89.93Å 137.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 2.30 19.96 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.3 (19.96-2.30) 97.3 (19.96-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	18.48 (at 2.30Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.210 , 0.255 0.210 , 0.254	Depositor DCC
R_{free} test set	1065 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtriage
Anisotropy	0.188	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3982	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UPG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/3775	0.96	14/5088 (0.3%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	399	PHE	N-CA-C	7.60	121.96	112.24
1	A	158	VAL	N-CA-C	7.31	118.25	111.45
1	A	132	SER	N-CA-C	-6.51	100.57	110.14
1	A	362	MSE	N-CA-C	6.47	121.18	113.16
1	A	225	ALA	N-CA-C	6.28	118.97	108.73
1	A	159	GLU	N-CA-C	6.15	119.54	109.76
1	A	453	GLY	N-CA-C	-6.08	104.24	112.57
1	A	345	ALA	N-CA-C	-5.99	106.01	113.38
1	A	262	GLN	N-CA-C	-5.91	100.34	109.14
1	A	136	SER	N-CA-C	5.49	116.96	110.97
1	A	462	VAL	N-CA-C	5.36	115.61	108.11
1	A	475	ILE	CA-C-N	5.26	125.54	119.92
1	A	475	ILE	C-N-CA	5.26	125.54	119.92
1	A	156	SER	N-CA-C	5.09	116.91	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3721	0	3707	103	0
2	A	36	0	22	12	0
3	A	225	0	0	12	0
All	All	3982	0	3729	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:ASN:ND2	2:A:5206:UPG:H2'	1.49	1.28
1:A:89:MSE:HE2	1:A:424:MSE:HE1	1.25	1.14
1:A:248:ARG:HH11	1:A:248:ARG:HG2	1.03	1.11
1:A:308:ASN:ND2	2:A:5206:UPG:C2'	2.27	0.97
1:A:310:LEU:HD22	1:A:356:MSE:HE3	1.45	0.96
1:A:74:GLN:HE22	1:A:122:SER:HB2	1.31	0.96
1:A:308:ASN:OD1	1:A:356:MSE:HE2	1.66	0.95
1:A:248:ARG:HG2	1:A:248:ARG:NH1	1.77	0.95
1:A:426:GLY:O	1:A:429:LYS:HD3	1.66	0.95
1:A:308:ASN:HD22	2:A:5206:UPG:H2'	1.33	0.92
1:A:316:VAL:O	1:A:320:THR:HG23	1.84	0.77
1:A:74:GLN:NE2	1:A:122:SER:HB2	2.00	0.77
1:A:189:PRO:HG2	1:A:193:ASP:HB2	1.66	0.76
1:A:308:ASN:HD22	2:A:5206:UPG:C1'	1.99	0.75
1:A:308:ASN:HD21	2:A:5206:UPG:H2'	1.55	0.72
1:A:308:ASN:HD22	2:A:5206:UPG:C2'	1.92	0.72
1:A:469:SER:HA	3:A:5324:HOH:O	1.91	0.69
1:A:89:MSE:HE2	1:A:424:MSE:CE	2.15	0.68
1:A:191:HIS:CD2	2:A:5206:UPG:H5C1	2.30	0.67
1:A:248:ARG:HH11	1:A:248:ARG:CG	1.93	0.66
1:A:310:LEU:HD22	1:A:356:MSE:CE	2.25	0.63
1:A:464:ILE:HD13	1:A:475:ILE:HD11	1.80	0.63
1:A:468:ASP:O	1:A:469:SER:HB2	2.00	0.62
1:A:323:GLU:C	1:A:323:GLU:OE2	2.43	0.61
1:A:268:GLY:O	1:A:269:LYS:HB2	2.00	0.61
1:A:466:ASN:CG	1:A:468:ASP:OD1	2.45	0.59
1:A:245:GLU:HB2	1:A:309:ASN:HB2	1.85	0.58
1:A:189:PRO:HG2	1:A:193:ASP:CB	2.33	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:487:SER:HB2	1:A:488:PRO:HD2	1.86	0.57
1:A:450:VAL:HG12	1:A:451:GLN:N	2.20	0.57
1:A:248:ARG:NH1	3:A:5273:HOH:O	2.38	0.56
1:A:439:VAL:HG13	1:A:440:GLU:HG2	1.86	0.56
1:A:153:VAL:HB	1:A:157:GLU:HG2	1.88	0.56
1:A:86:GLY:HA3	1:A:91:LEU:HD12	1.89	0.55
1:A:248:ARG:NH1	1:A:248:ARG:CG	2.57	0.55
1:A:278:ARG:HD2	3:A:5249:HOH:O	2.06	0.55
1:A:393:TYR:CZ	1:A:403:LEU:HD13	2.41	0.55
1:A:308:ASN:ND2	2:A:5206:UPG:C1'	2.63	0.55
1:A:400:ARG:HH11	1:A:400:ARG:HG3	1.71	0.55
1:A:310:LEU:CD2	1:A:356:MSE:HG2	2.36	0.54
1:A:380:LYS:HE3	2:A:5206:UPG:O6'	2.07	0.54
1:A:20:ARG:HH11	1:A:20:ARG:HG3	1.73	0.54
1:A:464:ILE:HD13	1:A:475:ILE:CD1	2.38	0.53
1:A:91:LEU:HD21	1:A:424:MSE:HE3	1.91	0.53
1:A:100:VAL:HG13	1:A:388:LEU:HD13	1.89	0.53
1:A:16:VAL:HG23	1:A:33:ILE:HD11	1.90	0.53
1:A:404:ASP:OD1	1:A:406:ARG:HG3	2.10	0.52
1:A:73:LEU:HD13	1:A:73:LEU:C	2.35	0.52
1:A:73:LEU:HD13	1:A:73:LEU:O	2.10	0.51
1:A:205:GLN:O	1:A:209:GLU:HG3	2.11	0.51
1:A:294:GLU:HB3	3:A:5317:HOH:O	2.11	0.51
1:A:482:ASN:O	1:A:483:ASP:C	2.53	0.51
1:A:400:ARG:NE	3:A:5240:HOH:O	2.42	0.50
1:A:96:THR:HG23	3:A:5259:HOH:O	2.11	0.50
1:A:461:THR:O	1:A:483:ASP:HA	2.11	0.50
1:A:38:MSE:HE3	3:A:5262:HOH:O	2.13	0.49
1:A:429:LYS:HE3	1:A:449:LEU:HD11	1.94	0.49
1:A:450:VAL:CG1	1:A:451:GLN:N	2.75	0.49
1:A:53:MSE:HE3	1:A:53:MSE:HA	1.94	0.49
1:A:310:LEU:HD23	1:A:356:MSE:HG2	1.94	0.49
1:A:444:VAL:HA	1:A:462:VAL:O	2.13	0.49
1:A:320:THR:HA	1:A:323:GLU:HG3	1.94	0.48
1:A:468:ASP:OD1	1:A:468:ASP:N	2.43	0.48
1:A:85:LEU:HB3	1:A:87:THR:HG23	1.94	0.48
1:A:400:ARG:NH2	3:A:5240:HOH:O	2.44	0.48
1:A:240:ILE:HG23	1:A:365:SER:O	2.14	0.48
1:A:316:VAL:HA	1:A:319:GLU:HG2	1.96	0.48
2:A:5206:UPG:H5C2	2:A:5206:UPG:H6	1.96	0.47
1:A:53:MSE:HE3	3:A:5417:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:MSE:HG3	1:A:54:PRO:HD2	1.97	0.47
1:A:53:MSE:HG3	1:A:54:PRO:CD	2.45	0.47
1:A:313:ARG:HG2	1:A:316:VAL:HG23	1.97	0.47
1:A:289:PRO:HG2	1:A:292:ASP:HB2	1.97	0.46
1:A:282:LEU:HD22	1:A:283:ARG:N	2.30	0.46
1:A:107:LEU:HD23	1:A:142:PHE:CD2	2.52	0.45
1:A:147:TYR:HD2	1:A:149:TRP:CZ3	2.35	0.45
1:A:480:LYS:C	1:A:481:LEU:HD23	2.42	0.45
1:A:458:LEU:HD23	1:A:458:LEU:N	2.31	0.45
1:A:172:THR:OG1	1:A:174:GLU:HG3	2.17	0.44
1:A:191:HIS:HD2	2:A:5206:UPG:H5C1	1.79	0.44
1:A:261:ARG:CZ	1:A:278:ARG:HG3	2.47	0.44
1:A:231:VAL:HG11	1:A:311:TRP:CE2	2.52	0.44
1:A:147:TYR:HD2	1:A:149:TRP:CH2	2.36	0.44
1:A:74:GLN:HE22	1:A:122:SER:CB	2.16	0.44
1:A:439:VAL:HG13	1:A:440:GLU:N	2.33	0.43
1:A:457:VAL:C	1:A:458:LEU:HD23	2.43	0.43
1:A:89:MSE:CE	1:A:424:MSE:HE1	2.19	0.43
1:A:308:ASN:HD21	2:A:5206:UPG:C2'	2.17	0.43
1:A:282:LEU:CD2	1:A:283:ARG:N	2.83	0.42
1:A:475:ILE:HA	1:A:476:PRO:HD3	1.87	0.42
1:A:110:THR:O	1:A:113:GLN:HB2	2.20	0.42
1:A:217:VAL:HG11	1:A:356:MSE:SE	2.69	0.42
1:A:136:SER:HB3	3:A:5412:HOH:O	2.21	0.41
1:A:288:CYS:O	1:A:289:PRO:C	2.63	0.41
1:A:248:ARG:HA	1:A:248:ARG:HD3	1.71	0.41
1:A:308:ASN:OD1	1:A:356:MSE:CE	2.54	0.41
1:A:398:ASP:OD1	1:A:400:ARG:NH1	2.53	0.41
1:A:172:THR:O	1:A:173:LEU:HB2	2.21	0.41
1:A:428:GLU:HG3	3:A:5269:HOH:O	2.20	0.41
1:A:429:LYS:HE3	1:A:449:LEU:CD1	2.51	0.41
1:A:107:LEU:HD12	1:A:107:LEU:HA	1.93	0.41
1:A:373:ARG:HD3	3:A:5365:HOH:O	2.21	0.40
1:A:191:HIS:C	1:A:193:ASP:N	2.79	0.40
1:A:468:ASP:O	1:A:469:SER:CB	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	480/505 (95%)	454 (95%)	24 (5%)	2 (0%)	30	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	483	ASP
1	A	469	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/413 (99%)	382 (93%)	27 (7%)	15	21

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	13	GLN
1	A	46	SER
1	A	49	ASP
1	A	53	MSE
1	A	61	LEU
1	A	138	SER
1	A	152	GLN
1	A	164	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	248	ARG
1	A	266	VAL
1	A	267	LYS
1	A	270	ASP
1	A	278	ARG
1	A	282	LEU
1	A	290	LYS
1	A	310	LEU
1	A	313	ARG
1	A	320	THR
1	A	323	GLU
1	A	339	VAL
1	A	362	MSE
1	A	386	LEU
1	A	388	LEU
1	A	428	GLU
1	A	429	LYS
1	A	438	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	74	GLN
1	A	113	GLN
1	A	124	HIS
1	A	164	GLN
1	A	308	ASN
1	A	322	GLN
1	A	451	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UPG	A	5206	-	37,38,38	1.40	5 (13%)	55,58,58	1.11	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPG	A	5206	-	-	3/23/59/59	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5206	UPG	PA-O3A	-4.07	1.55	1.59
2	A	5206	UPG	C2-N1	3.45	1.43	1.38
2	A	5206	UPG	C5-C4	2.50	1.49	1.43
2	A	5206	UPG	O5C-C5C	-2.45	1.35	1.44
2	A	5206	UPG	PA-O2A	-2.03	1.45	1.55

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5206	UPG	O6'-C6'-C5'	-2.47	102.92	111.33
2	A	5206	UPG	PB-O3B-C1'	2.21	130.22	121.21
2	A	5206	UPG	O4-C4-N3	2.03	122.22	119.27
2	A	5206	UPG	O2A-PA-O5C	2.02	116.71	107.57

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	5206	UPG	C5C-O5C-PA-O1A
2	A	5206	UPG	C5C-O5C-PA-O2A
2	A	5206	UPG	C5C-O5C-PA-O3A

There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5206	UPG	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	466/505 (92%)	0.07	16 (3%)	48 50	9, 26, 50, 74	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	268	GLY	3.5
1	A	92	CYS	3.3
1	A	265	TYR	3.2
1	A	466	ASN	3.1
1	A	266	VAL	2.8
1	A	467	THR	2.8
1	A	468	ASP	2.8
1	A	275	ALA	2.6
1	A	133	PHE	2.6
1	A	274	ASP	2.5
1	A	470	ALA	2.5
1	A	134	ASN	2.3
1	A	324	HIS	2.3
1	A	471	SER	2.2
1	A	269	LYS	2.2
1	A	267	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UPG	A	5206	36/36	0.95	0.07	11,16,23,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.