



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 02:58 AM UTC

PDB ID : 4OE1 / pdb_00004oe1
Title : Crystal structure of the pentatricopeptide repeat protein PPR10 (C256S/C430S/C449S) in complex with an 18-nt PSAJ rna element
Authors : Li, Q.; Yan, C.; Wu, J.; Yin, P.; Yan, N.
Deposited on : 2014-01-11
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)
RNA backbone	3983	1114 (3.00-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	718	7% 48% 39% 8% • •
1	B	718	5% 53% 35% 7% 5%
2	C	18	6% 44% 11% 22% 22%
2	D	18	6% 11% 44% 44%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chloroplast pentatricopeptide repeat protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	687	Total	C	N	O	S	16	0	0
			5202	3284	890	985	43			
1	B	683	Total	C	N	O	S	14	0	0
			5179	3273	891	973	42			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	256	SER	CYS	engineered mutation	UNP B8Y6I0
A	430	SER	CYS	engineered mutation	UNP B8Y6I0
A	449	SER	CYS	engineered mutation	UNP B8Y6I0
B	256	SER	CYS	engineered mutation	UNP B8Y6I0
B	430	SER	CYS	engineered mutation	UNP B8Y6I0
B	449	SER	CYS	engineered mutation	UNP B8Y6I0

- Molecule 2 is a RNA chain called psaJ RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	18	Total	C	N	O	P	3	0	0
			368	167	53	131	17			
2	C	14	Total	C	N	O	P	4	0	0
			279	126	41	99	13			

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	5	0
			5	4	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chloroplast pentatricopeptide repeat protein 10





• Molecule 1: Chloroplast pentatricopeptide repeat protein 10



• Molecule 2: psaJ RNA



• Molecule 2: psaJ RNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	83.32Å 83.32Å 225.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.26 – 2.80 34.26 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.2 (34.26-2.80) 98.1 (34.26-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.263 , 0.278 0.260 , 0.270	Depositor DCC
R_{free} test set	1855 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	67.0	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 62.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.054 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	11033	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.94	2/5294 (0.0%)	0.88	12/7177 (0.2%)
1	B	1.07	4/5269 (0.1%)	0.94	13/7136 (0.2%)
2	C	0.29	0/308	0.49	0/474
2	D	0.40	0/408	0.66	0/631
All	All	0.98	6/11279 (0.1%)	0.89	25/15418 (0.2%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	548	GLU	C-N	-12.99	1.17	1.33
1	B	748	CYS	C-N	7.84	1.44	1.33
1	B	743	VAL	CA-CB	-6.14	1.47	1.54
1	A	260	LEU	CA-C	-5.64	1.47	1.53
1	A	594	PRO	C-N	-5.37	1.25	1.33

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	242	GLU	CA-C-N	8.75	128.79	119.78
1	B	242	GLU	C-N-CA	8.75	128.79	119.78
1	B	743	VAL	CB-CA-C	-6.98	103.53	111.55
1	B	313	PRO	N-CA-CB	6.85	110.54	103.00
1	A	593	ASN	CA-C-N	6.61	128.11	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5202	0	5106	391	2
1	B	5179	0	5068	291	1
2	C	279	0	146	10	0
2	D	368	0	189	57	0
3	B	5	0	0	0	0
All	All	11033	0	10509	715	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 715 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:706:HIS:CE1	1:A:738:LEU:HD22	1.19	1.62
1:A:296:TYR:CE1	1:A:300:LEU:HD11	1.38	1.59
1:A:489:CYS:SG	1:A:491:THR:HG22	1.50	1.51
1:A:79:LEU:HD13	1:A:98:ARG:NH2	1.32	1.43
1:A:515:LYS:NZ	1:A:519:LYS:HE3	1.21	1.43

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:PRO:O	1:A:447:ARG:NH1[1_565]	1.96	0.24
1:A:413:ARG:NE	1:B:518:THR:O[3_744]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	677/718 (94%)	599 (88%)	65 (10%)	13 (2%)	6	22
1	B	671/718 (94%)	605 (90%)	50 (8%)	16 (2%)	4	17
All	All	1348/1436 (94%)	1204 (89%)	115 (8%)	29 (2%)	5	19

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	168	PRO
1	B	315	ALA
1	B	665	PRO
1	A	511	SER
1	A	698	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/587 (92%)	449 (83%)	90 (17%)	2	8
1	B	530/587 (90%)	467 (88%)	63 (12%)	5	17
All	All	1069/1174 (91%)	916 (86%)	153 (14%)	3	11

5 of 153 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	363	VAL
1	B	656	GLN
1	B	390	LEU
1	B	550	GLU
1	B	747	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	153	GLN
1	B	284	ASN
1	B	661	GLN
1	B	213	ASN
1	B	473	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	12/18 (66%)	5 (41%)	2 (16%)
2	D	17/18 (94%)	9 (52%)	3 (17%)
All	All	29/36 (80%)	14 (48%)	5 (17%)

5 of 14 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	6	C
2	D	7	U
2	D	9	U
2	D	10	A
2	D	11	A

All (5) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	5	U
2	D	9	U
2	D	14	A
2	C	12	U
2	C	13	U

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	801	-	4,4,4	0.96	0	6,6,6	0.47	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	548:GLU	C	549:ASN	N	1.17

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	687/718 (95%)	0.49	52 (7%)	20 14	30, 72, 128, 229	103 (14%)
1	B	683/718 (95%)	0.43	33 (4%)	35 28	26, 72, 123, 237	86 (12%)
2	C	14/18 (77%)	0.38	1 (7%)	22 16	56, 72, 86, 120	4 (28%)
2	D	18/18 (100%)	0.92	1 (5%)	30 23	41, 79, 105, 158	6 (33%)
All	All	1402/1472 (95%)	0.47	87 (6%)	26 20	26, 72, 127, 237	199 (14%)

The worst 5 of 87 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	THR	7.5
2	D	13	U	5.2
1	A	748	CYS	5.2
1	A	292	LYS	5.0
1	A	293	ALA	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	B	801	5/5	-	-	78,79,79,79	5

6.5 Other polymers [i](#)

There are no such residues in this entry.