



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:18 PM UTC

PDB ID : 5OE6 / pdb_00005oe6
Title : Crystal structure of the N-terminal domain of PqsA in complex with 6-fluoro anthraniloyl-AMP (crystal form 1)
Authors : Witzgall, F.; Ewert, W.; Blankenfeldt, W.
Deposited on : 2017-07-07
Resolution : 1.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1054 (1.68-1.68)
Clashscore	190562	1078 (1.68-1.68)
Ramachandran outliers	187476	1068 (1.68-1.68)
Sidechain outliers	187428	1067 (1.68-1.68)
RSRZ outliers	180081	1055 (1.68-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	84% 13% .
1	B	407	90% 6% .
1	C	407	% 82% 14% .
1	D	407	90% 6% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 25754 atoms, of which 12038 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anthranilate–CoA ligase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	394	Total	C	H	N	O	S	0	8	0
			6073	1947	3017	539	557	13			
1	B	392	Total	C	H	N	O	S	0	6	0
			6054	1937	3012	543	549	13			
1	C	393	Total	C	H	N	O	S	0	5	0
			5985	1923	2964	535	550	13			
1	D	391	Total	C	H	N	O	S	0	3	0
			5970	1919	2959	532	548	12			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	400	LEU	-	expression tag	UNP Q9I4X3
A	401	GLU	-	expression tag	UNP Q9I4X3
A	402	HIS	-	expression tag	UNP Q9I4X3
A	403	HIS	-	expression tag	UNP Q9I4X3
A	404	HIS	-	expression tag	UNP Q9I4X3
A	405	HIS	-	expression tag	UNP Q9I4X3
A	406	HIS	-	expression tag	UNP Q9I4X3
A	407	HIS	-	expression tag	UNP Q9I4X3
B	400	LEU	-	expression tag	UNP Q9I4X3
B	401	GLU	-	expression tag	UNP Q9I4X3
B	402	HIS	-	expression tag	UNP Q9I4X3
B	403	HIS	-	expression tag	UNP Q9I4X3
B	404	HIS	-	expression tag	UNP Q9I4X3
B	405	HIS	-	expression tag	UNP Q9I4X3
B	406	HIS	-	expression tag	UNP Q9I4X3
B	407	HIS	-	expression tag	UNP Q9I4X3
C	400	LEU	-	expression tag	UNP Q9I4X3
C	401	GLU	-	expression tag	UNP Q9I4X3
C	402	HIS	-	expression tag	UNP Q9I4X3
C	403	HIS	-	expression tag	UNP Q9I4X3
C	404	HIS	-	expression tag	UNP Q9I4X3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	405	HIS	-	expression tag	UNP Q9I4X3
C	406	HIS	-	expression tag	UNP Q9I4X3
C	407	HIS	-	expression tag	UNP Q9I4X3
D	400	LEU	-	expression tag	UNP Q9I4X3
D	401	GLU	-	expression tag	UNP Q9I4X3
D	402	HIS	-	expression tag	UNP Q9I4X3
D	403	HIS	-	expression tag	UNP Q9I4X3
D	404	HIS	-	expression tag	UNP Q9I4X3
D	405	HIS	-	expression tag	UNP Q9I4X3
D	406	HIS	-	expression tag	UNP Q9I4X3
D	407	HIS	-	expression tag	UNP Q9I4X3

- Molecule 2 is 6-fluoroanthraniloyl-AMP (CCD ID: 9SN) (formula: C₁₇H₁₈FN₆O₈P).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	F	H	N	O	P	0	0
			51	17	1	18	6	8	1		
2	B	1	Total	C	F	H	N	O	P	0	0
			51	17	1	18	6	8	1		
2	C	1	Total	C	F	H	N	O	P	0	0
			51	17	1	18	6	8	1		
2	D	1	Total	C	F	H	N	O	P	0	0
			51	17	1	18	6	8	1		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	360	Total	O	0	2
			360	360		
4	B	388	Total	O	0	1
			388	388		
4	C	352	Total	O	0	0
			352	352		
4	D	344	Total	O	0	0
			344	344		

- Molecule 1: Anthranilate–CoA ligase





Chain D:

90%

6%

 .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.56Å 84.13Å 138.21Å 90.00° 92.59° 90.00°	Depositor
Resolution (Å)	46.02 – 1.67 46.02 – 1.67	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.02-1.67) 87.2 (46.02-1.67)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.67Å)	Xtriage
Refinement program	PHENIX 1.12rc2_2821	Depositor
R, R_{free}	0.270 , 0.313 0.269 , 0.314	Depositor DCC
R_{free} test set	9232 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	9.6	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.119 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	25754	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1411e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, 9SN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/3156	0.41	0/4292
1	B	0.26	0/3136	0.49	0/4262
1	C	0.18	0/3112	0.40	0/4233
1	D	0.26	0/3096	0.49	0/4210
All	All	0.23	0/12500	0.45	0/16997

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3056	3017	3016	48	1
1	B	3042	3012	3012	27	0
1	C	3021	2964	2964	64	1
1	D	3011	2959	2958	23	0
2	A	33	18	0	1	0
2	B	33	18	0	0	0
2	C	33	18	0	1	0
2	D	33	18	0	1	0
3	C	10	14	14	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	360	0	0	30	8
4	B	388	0	0	18	6
4	C	352	0	0	45	8
4	D	344	0	0	16	6
All	All	13716	12038	11964	166	19

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 166 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:309:VAL:O	4:D:1002:HOH:O	1.85	0.94
1:A:89:GLU:OE1	4:A:1001:HOH:O	1.85	0.94
1:D:367:GLU:O	4:D:1001:HOH:O	1.85	0.93
1:A:314:ARG:NH1	4:A:1005:HOH:O	2.04	0.91
1:D:385:GLU:OE2	1:D:393:ARG:NH1	2.05	0.89

The worst 5 of 19 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1028:HOH:O	4:D:1138:HOH:O[1_545]	1.75	0.45
4:B:1292:HOH:O	4:D:1214:HOH:O[2_746]	1.88	0.32
4:C:1221:HOH:O	4:C:1303:HOH:O[2_545]	1.91	0.29
4:C:1010:HOH:O	4:C:1191:HOH:O[2_545]	1.95	0.25
4:B:1268:HOH:O	4:B:1321:HOH:O[2_646]	1.96	0.24

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/407 (98%)	391 (98%)	7 (2%)	0	100	100
1	B	394/407 (97%)	388 (98%)	6 (2%)	0	100	100
1	C	394/407 (97%)	388 (98%)	6 (2%)	0	100	100
1	D	390/407 (96%)	383 (98%)	7 (2%)	0	100	100
All	All	1576/1628 (97%)	1550 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/320 (98%)	313 (99%)	2 (1%)	78	67
1	B	312/320 (98%)	310 (99%)	2 (1%)	78	67
1	C	308/320 (96%)	308 (100%)	0	100	100
1	D	307/320 (96%)	307 (100%)	0	100	100
All	All	1242/1280 (97%)	1238 (100%)	4 (0%)	86	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	240	LEU
1	A	337	ARG
1	B	66	LEU
1	B	117	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	340	HIS
1	B	317	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	42	GLN
1	C	264	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	9SN	C	900	-	36,36,36	0.48	0	54,54,54	0.68	2 (3%)
2	9SN	B	900	-	36,36,36	0.48	1 (2%)	54,54,54	0.47	0
2	9SN	D	900	-	36,36,36	0.42	1 (2%)	54,54,54	0.51	0
3	PGE	C	901	-	9,9,9	0.28	0	8,8,8	0.46	0
2	9SN	A	900	-	36,36,36	0.49	1 (2%)	54,54,54	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9SN	C	900	-	-	4/19/35/35	0/4/4/4
2	9SN	B	900	-	-	4/19/35/35	0/4/4/4
2	9SN	D	900	-	-	2/19/35/35	0/4/4/4
3	PGE	C	901	-	-	4/7/7/7	-
2	9SN	A	900	-	-	4/19/35/35	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	900	9SN	P12-O14	2.25	1.64	1.60
2	A	900	9SN	P12-O14	2.25	1.64	1.60
2	D	900	9SN	P12-O14	2.11	1.64	1.60

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	900	9SN	O14-C24-O26	2.97	124.52	122.13
2	C	900	9SN	P12-O14-C24	2.15	127.69	123.01

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	901	PGE	O2-C3-C4-O3
3	C	901	PGE	O1-C1-C2-O2
3	C	901	PGE	O3-C5-C6-O4
2	A	900	9SN	O14-C24-C25-C31
3	C	901	PGE	C3-C4-O3-C5


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	900	9SN	1	0
2	D	900	9SN	1	0
3	C	901	PGE	3	0
2	A	900	9SN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 9SN B 900




Bond lengths



Bond angles




Torsions




Rings


Ligand 9SN D 900




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/407 (96%)	-0.42	2 (0%) 87 92	12, 27, 42, 60	8 (2%)
1	B	392/407 (96%)	-1.04	0 100 100	3, 12, 27, 45	6 (1%)
1	C	393/407 (96%)	-0.29	3 (0%) 82 90	10, 29, 48, 63	5 (1%)
1	D	391/407 (96%)	-1.00	0 100 100	3, 14, 29, 35	3 (0%)
All	All	1570/1628 (96%)	-0.69	5 (0%) 90 95	3, 23, 40, 63	22 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	269	LEU	3.5
1	A	118	ALA	3.1
1	C	115	GLY	2.1
1	A	145	LEU	2.1
1	C	244	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PGE	C	901	10/10	0.98	0.06	23,35,42,50	0
2	9SN	B	900	33/33	0.99	0.03	8,8,10,13	0
2	9SN	C	900	33/33	0.99	0.05	5,15,24,34	0
2	9SN	D	900	33/33	0.99	0.03	8,8,11,23	0
2	9SN	A	900	33/33	0.99	0.03	8,19,25,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 9SN B 900:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9SN C 900:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9SN D 900:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9SN A 900:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.