



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:31 AM UTC

PDB ID : 8OEG / pdb_00008oeg
Title : PDE4B bound to MAPI compound 92a
Authors : Rizzi, A.; Armani, E.
Deposited on : 2023-03-10
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2881 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2714	1711	460	524	19	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	245	MET	-	initiating methionine	UNP Q07343
A	295	LYS	SER	conflict	UNP Q07343
A	296	VAL	PRO	conflict	UNP Q07343
A	298	ALA	GLN	conflict	UNP Q07343
A	299	GLU	LYS	conflict	UNP Q07343
A	300	GLU	ASP	conflict	UNP Q07343
A	?	-	ARG	deletion	UNP Q07343
A	?	-	GLU	deletion	UNP Q07343
A	?	-	LYS	deletion	UNP Q07343
A	?	-	LYS	deletion	UNP Q07343
A	?	-	LYS	deletion	UNP Q07343
A	301	ALA	LYS	conflict	UNP Q07343
A	302	PRO	GLN	conflict	UNP Q07343
A	304	PRO	LEU	conflict	UNP Q07343
A	660	HIS	-	expression tag	UNP Q07343
A	661	HIS	-	expression tag	UNP Q07343
A	662	HIS	-	expression tag	UNP Q07343
A	663	HIS	-	expression tag	UNP Q07343
A	664	HIS	-	expression tag	UNP Q07343
A	665	HIS	-	expression tag	UNP Q07343


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is [(1 {S})-2-[3,5-bis(chloranyl)-1-oxidanyl-pyridin-4-yl]-1-(3,4-dimethoxyphenyl)ethyl] 5-[[[(1 {R})-2-[[[(3 {R})-1-azabicyclo[2.2.2]octan-3-yl]oxy]-2-oxidanylidene-1-phenyl-ethyl]amino]methyl]thiophene-2-carboxylate (CCD ID: VL9) (formula: C₃₆H₃₈Cl₂N₃O₇S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	S	0	0
			49	36	2	3	7	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	116	Total	O	0	0
			116	116		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4B



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.76Å 95.76Å 85.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	83.07 – 1.89 82.93 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.5 (83.07-1.89) 99.5 (82.93-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.158 , 0.195 0.191 , 0.227	Depositor DCC
R_{free} test set	1771 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 23.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2881	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VL9, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.76	31/2770 (1.1%)	1.43	18/3757 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	558	ASN	C-O	-8.19	1.14	1.24
1	A	346	ASN	N-CA	-8.19	1.36	1.46
1	A	534	GLU	C-O	-7.62	1.14	1.24
1	A	576	ARG	CD-NE	-7.62	1.35	1.46
1	A	329	GLY	N-CA	7.61	1.56	1.45
1	A	329	GLY	CA-C	7.33	1.62	1.51
1	A	557	ARG	CD-NE	-6.66	1.36	1.46
1	A	333	GLU	N-CA	6.59	1.54	1.46
1	A	372	PHE	N-CA	-6.36	1.38	1.46
1	A	402	ASP	C-O	-6.33	1.15	1.24
1	A	393	MET	SD-CE	6.24	1.95	1.79
1	A	607	HIS	C-O	-6.24	1.15	1.24
1	A	464	GLU	CA-C	-6.13	1.44	1.52
1	A	623	VAL	C-O	6.11	1.30	1.24
1	A	555	VAL	CA-CB	6.09	1.61	1.54
1	A	545	LEU	N-CA	6.08	1.54	1.46
1	A	511	ILE	C-O	6.07	1.31	1.24
1	A	429	ALA	CA-C	-5.98	1.43	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	561	HIS	N-CA	5.87	1.53	1.46
1	A	524	SER	CB-OG	-5.83	1.30	1.42
1	A	653	GLN	CG-CD	-5.66	1.38	1.52
1	A	535	THR	N-CA	-5.48	1.39	1.46
1	A	630	TRP	C-O	5.47	1.30	1.24
1	A	610	SER	CA-C	-5.45	1.46	1.52
1	A	557	ARG	NE-CZ	5.42	1.39	1.33
1	A	455	ASN	N-CA	5.41	1.52	1.46
1	A	403	VAL	N-CA	5.31	1.53	1.46
1	A	631	ALA	N-CA	5.15	1.52	1.46
1	A	535	THR	CA-C	5.07	1.59	1.52
1	A	601	SER	CA-C	5.03	1.59	1.53
1	A	614	SER	N-CA	-5.03	1.40	1.46

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	PHE	CA-C-N	-9.93	101.95	121.41
1	A	328	PHE	C-N-CA	-9.93	101.95	121.41
1	A	543	VAL	N-CA-CB	-8.73	102.39	112.34
1	A	524	SER	CB-CA-C	-7.11	99.67	110.90
1	A	576	ARG	NE-CZ-NH1	5.92	127.42	121.50
1	A	559	MET	N-CA-C	5.81	117.61	111.28
1	A	657	PRO	CA-C-N	5.69	131.95	121.70
1	A	657	PRO	C-N-CA	5.69	131.95	121.70
1	A	332	THR	CB-CA-C	-5.60	99.27	110.42
1	A	614	SER	N-CA-CB	5.55	118.38	110.16
1	A	328	PHE	O-C-N	-5.40	115.14	122.81
1	A	576	ARG	NE-CZ-NH2	-5.30	114.42	119.20
1	A	536	LYS	CB-CG-CD	-5.30	99.11	111.30
1	A	329	GLY	N-CA-C	5.29	125.71	113.18
1	A	393	MET	CG-SD-CE	-5.17	89.52	100.90
1	A	330	VAL	CB-CA-C	-5.05	103.01	111.29
1	A	520	SER	CA-CB-OG	-5.05	101.01	111.10
1	A	532	MET	CB-CG-SD	-5.01	97.67	112.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	329	GLY	Peptide
1	A	332	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2714	0	2639	22	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	49	0	0	1	0
5	A	116	0	0	4	0
All	All	2881	0	2639	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:332:THR:HG23	1:A:358:TYR:O	1.46	1.14
1:A:539:THR:HG22	1:A:543:VAL:H	1.42	0.82
1:A:610:SER:HB3	5:A:860:HOH:O	1.81	0.80
1:A:610:SER:CB	5:A:860:HOH:O	2.31	0.79
1:A:539:THR:HG23	1:A:541:SER:H	1.56	0.70
1:A:332:THR:CG2	1:A:358:TYR:O	2.34	0.69
1:A:328:PHE:O	1:A:329:GLY:C	2.41	0.64
1:A:539:THR:HG23	1:A:541:SER:N	2.15	0.61
1:A:508:LYS:HD2	5:A:843:HOH:O	2.11	0.50
1:A:610:SER:HB2	5:A:860:HOH:O	2.03	0.50
1:A:539:THR:CG2	1:A:541:SER:OG	2.62	0.47
1:A:539:THR:HG21	1:A:541:SER:OG	2.14	0.47
1:A:554:GLN:NE2	1:A:557:ARG:NH1	2.64	0.46
1:A:530:LYS:O	1:A:534:GLU:HG3	2.15	0.46
1:A:332:THR:HG22	1:A:334:ASN:HB2	1.99	0.45
1:A:519:MET:HG2	4:A:703:VL9:C11	2.48	0.44
1:A:442:ALA:O	1:A:446:HIS:HB3	2.19	0.41
1:A:469:TYR:CE1	1:A:477:ASN:HB3	2.56	0.41
1:A:332:THR:C	1:A:334:ASN:H	2.28	0.41
1:A:340:LYS:HD3	1:A:340:LYS:O	2.21	0.41
1:A:430:VAL:HG11	1:A:546:LEU:HD12	2.03	0.41
1:A:332:THR:HG22	1:A:334:ASN:H	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	334/421 (79%)	328 (98%)	5 (2%)	1 (0%)	36 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	305/385 (79%)	290 (95%)	15 (5%)	22 14

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	326	SER
1	A	332	THR
1	A	333	GLU
1	A	340	LYS
1	A	385	SER
1	A	465	LEU
1	A	489	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	501	LYS
1	A	532	MET
1	A	539	THR
1	A	543	VAL
1	A	573	GLU
1	A	610	SER
1	A	653	GLN
1	A	658	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	334	ASN
1	A	548	ASN
1	A	653	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	VL9	A	703	-	48,54,54	1.98	12 (25%)	63,76,76	2.09	16 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VL9	A	703	-	-	4/37/71/71	0/7/6/6

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	703	VL9	O2-C18	5.74	1.47	1.34
4	A	703	VL9	C20-N2	5.18	1.55	1.46
4	A	703	VL9	C10-C9	4.17	1.42	1.35
4	A	703	VL9	C21-N2	3.43	1.56	1.46
4	A	703	VL9	C16-N1	-3.33	1.42	1.46
4	A	703	VL9	C26-C17	-3.22	1.47	1.52
4	A	703	VL9	C24-C23	2.97	1.60	1.53
4	A	703	VL9	C23-C19	2.67	1.58	1.52
4	A	703	VL9	C7-C6	2.44	1.59	1.51
4	A	703	VL9	C32-C33	2.44	1.47	1.39
4	A	703	VL9	C25-N2	2.15	1.52	1.46
4	A	703	VL9	C24-C25	2.03	1.57	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	VL9	C19-O2-C18	6.94	128.34	117.85
4	A	703	VL9	C35-O6-C	5.12	125.02	117.51
4	A	703	VL9	C7-C8-C9	4.43	124.19	117.92
4	A	703	VL9	O2-C18-C17	4.15	121.12	110.61
4	A	703	VL9	C25-N2-C20	3.80	115.86	109.53
4	A	703	VL9	C16-C15-C32	3.68	138.49	127.56
4	A	703	VL9	C7-C6-C4	3.27	118.73	113.01
4	A	703	VL9	C13-C14-S	-3.25	112.90	119.98
4	A	703	VL9	C18-C17-N1	-3.15	102.84	109.64
4	A	703	VL9	O2-C18-O3	-3.10	118.35	123.95
4	A	703	VL9	C16-C15-S	-3.06	113.84	120.70
4	A	703	VL9	C15-C16-N1	-2.79	104.72	112.06
4	A	703	VL9	C33-C14-C13	2.42	135.13	127.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	VL9	C26-C17-C18	-2.36	102.50	109.78
4	A	703	VL9	C21-N2-C20	-2.30	105.70	109.53
4	A	703	VL9	O1-C6-C7	-2.24	102.46	106.66

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	703	VL9	C32-C15-C16-N1
4	A	703	VL9	S-C15-C16-N1
4	A	703	VL9	C23-C19-O2-C18
4	A	703	VL9	C20-C19-O2-C18

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	703	VL9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	336/421 (79%)	0.14	11 (3%)	49 52	13, 20, 35, 54	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	329	GLY	4.3
1	A	325	ILE	3.6
1	A	323	THR	3.4
1	A	491	HIS	2.8
1	A	331	ASN	2.8
1	A	332	THR	2.8
1	A	428	ASP	2.7
1	A	658	GLN	2.7
1	A	326	SER	2.2
1	A	490	GLU	2.1
1	A	324	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	VL9	A	703	49/49	0.96	0.10	17,25,48,52	0
3	MG	A	702	1/1	0.99	0.10	11,11,11,11	0
2	ZN	A	701	1/1	1.00	0.06	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around VL9 A 703:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around MG A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.