



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:26 PM UTC

PDB ID : 8OEK / pdb_00008oek
Title : Crystal structure of the HormR-GAIN domains of adhesion GPCR ADGRB2 (BAI2) in the uncleaved state
Authors : Pohl, F.; Strater, N.
Deposited on : 2023-03-10
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	8% 68% 8% 23%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5446 atoms, of which 2692 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adhesion G protein-coupled receptor B2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	344	5395	1722	2678	474	505	16	0	1	0

There are 55 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	510	GLU	-	expression tag	UNP O60241
A	511	THR	-	expression tag	UNP O60241
A	512	GLY	-	expression tag	UNP O60241
A	513	TYR	-	expression tag	UNP O60241
A	514	PRO	-	expression tag	UNP O60241
A	515	TYR	-	expression tag	UNP O60241
A	516	ASP	-	expression tag	UNP O60241
A	517	VAL	-	expression tag	UNP O60241
A	518	PRO	-	expression tag	UNP O60241
A	519	ASP	-	expression tag	UNP O60241
A	520	TYR	-	expression tag	UNP O60241
A	521	ALA	-	expression tag	UNP O60241
A	522	THR	-	expression tag	UNP O60241
A	523	GLY	-	expression tag	UNP O60241
A	524	PRO	-	expression tag	UNP O60241
A	525	ARG	-	expression tag	UNP O60241
A	526	GLU	-	expression tag	UNP O60241
A	527	LEU	-	expression tag	UNP O60241
A	922	GLY	-	expression tag	UNP O60241
A	923	THR	-	expression tag	UNP O60241
A	924	ASP	-	expression tag	UNP O60241
A	925	ASP	-	expression tag	UNP O60241
A	926	ASP	-	expression tag	UNP O60241
A	927	ASP	-	expression tag	UNP O60241
A	928	LYS	-	expression tag	UNP O60241
A	929	SER	-	expression tag	UNP O60241
A	930	ALA	-	expression tag	UNP O60241

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	931	TRP	-	expression tag	UNP O60241
A	932	SER	-	expression tag	UNP O60241
A	933	HIS	-	expression tag	UNP O60241
A	934	PRO	-	expression tag	UNP O60241
A	935	GLN	-	expression tag	UNP O60241
A	936	PHE	-	expression tag	UNP O60241
A	937	GLU	-	expression tag	UNP O60241
A	938	LYS	-	expression tag	UNP O60241
A	939	GLY	-	expression tag	UNP O60241
A	940	GLY	-	expression tag	UNP O60241
A	941	GLY	-	expression tag	UNP O60241
A	942	SER	-	expression tag	UNP O60241
A	943	GLY	-	expression tag	UNP O60241
A	944	GLY	-	expression tag	UNP O60241
A	945	GLY	-	expression tag	UNP O60241
A	946	SER	-	expression tag	UNP O60241
A	947	GLY	-	expression tag	UNP O60241
A	948	GLY	-	expression tag	UNP O60241
A	949	SER	-	expression tag	UNP O60241
A	950	ALA	-	expression tag	UNP O60241
A	951	TRP	-	expression tag	UNP O60241
A	952	SER	-	expression tag	UNP O60241
A	953	HIS	-	expression tag	UNP O60241
A	954	PRO	-	expression tag	UNP O60241
A	955	GLN	-	expression tag	UNP O60241
A	956	PHE	-	expression tag	UNP O60241
A	957	GLU	-	expression tag	UNP O60241
A	958	LYS	-	expression tag	UNP O60241

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 3 is CACODYLATE ION (CCD ID: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	As	C	O	0	0
			5	1	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	18	Total	O	0	0
			18	18		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adhesion G protein-coupled receptor B2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.86Å 95.18Å 49.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.34 – 2.22 63.34 – 2.22	Depositor EDS
% Data completeness (in resolution range)	63.8 (63.34-2.22) 64.2 (63.34-2.22)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.229 , 0.287 0.230 , 0.286	Depositor DCC
R_{free} test set	626 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5446	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2784	0.36	0/3789

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2717	2678	2677	24	0
2	A	14	14	13	0	0
3	A	5	0	0	0	0
4	A	18	0	0	0	0
All	All	2754	2692	2690	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:723:VAL:HG21	1:A:752:VAL:HG21	1.68	0.75
1:A:750:ASP:OD1	1:A:754:HIS:ND1	2.26	0.67
1:A:844:VAL:CG1	1:A:911:LEU:HD22	2.32	0.60
1:A:844:VAL:HG11	1:A:911:LEU:HD22	1.89	0.55
1:A:820:LEU:HD13	1:A:842:MET:HE3	1.89	0.54
1:A:912:SER:OG	1:A:913:THR:N	2.40	0.54
1:A:644:ARG:CZ	1:A:697:VAL:HG12	2.41	0.51
1:A:631:TYR:HB2	1:A:635:ASP:HB2	1.94	0.49
1:A:615:MET:HA	1:A:615:MET:HE2	1.96	0.48
1:A:748:MET:HE2	1:A:752:VAL:CG1	2.43	0.48
1:A:858:ILE:HD13	1:A:911:LEU:HD23	1.99	0.44
1:A:833:PRO:N	1:A:834:PRO:HD2	2.31	0.44
1:A:583:SER:OG	1:A:683:ASP:OD1	2.17	0.44
1:A:748:MET:HE2	1:A:752:VAL:HG11	2.00	0.43
1:A:644:ARG:NE	1:A:697:VAL:HG12	2.33	0.43
1:A:723:VAL:HG23	1:A:751:TRP:CZ2	2.53	0.43
1:A:628:ARG:O	1:A:630:THR:HG23	2.19	0.42
1:A:548:ALA:O	1:A:551:GLU:HG2	2.19	0.42
1:A:615:MET:HE2	1:A:615:MET:CA	2.50	0.42
1:A:722:LEU:N	1:A:722:LEU:HD12	2.35	0.41
1:A:716:LEU:HD21	1:A:748:MET:SD	2.61	0.40
1:A:755:SER:OG	1:A:756:GLU:N	2.54	0.40
1:A:760:PHE:HB3	1:A:859:THR:HB	2.02	0.40
1:A:643:LEU:HD12	1:A:672:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/449 (76%)	325 (96%)	13 (4%)	1 (0%)	36 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	912	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	301/377 (80%)	295 (98%)	6 (2%)	48 63

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	552	ILE
1	A	633	SER
1	A	714	SER
1	A	722	LEU
1	A	725	SER
1	A	857	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	602	HIS
1	A	909	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CAC	A	1002	-	2,4,4	2.75	2 (100%)	4,6,6	1.76	1 (25%)
2	NAG	A	1001	1	14,14,15	0.33	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1001	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1002	CAC	AS-C2	2.79	1.96	1.90
3	A	1002	CAC	AS-C1	2.71	1.96	1.90

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002	CAC	O1-AS-C2	-2.13	108.85	111.50

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	NAG	C4-C5-C6-O6
2	A	1001	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/449 (76%)	0.96	35 (10%) 12 10	24, 40, 56, 72	1 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	769	SER	5.6
1	A	541	LEU	5.4
1	A	746	ARG	4.8
1	A	594	TYR	3.8
1	A	833	PRO	3.8
1	A	601	GLU	3.6
1	A	575	GLY	3.5
1	A	539	VAL	3.4
1	A	747	GLY	3.3
1	A	630	THR	3.3
1	A	811	SER	3.0
1	A	629	ARG	2.9
1	A	540	MET	2.9
1	A	543	THR	2.8
1	A	595	LEU	2.8
1	A	535	ARG	2.8
1	A	676	GLU	2.7
1	A	892	GLU	2.7
1	A	545	LYS	2.6
1	A	573	ALA	2.6
1	A	753	ARG	2.5
1	A	671	PHE	2.5
1	A	574	GLN	2.4
1	A	854	ALA	2.3
1	A	590	HIS	2.3
1	A	631	TYR	2.3
1	A	745	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	600	ARG	2.2
1	A	851	GLN	2.2
1	A	533	MET	2.2
1	A	865	ILE	2.1
1	A	571	LEU	2.1
1	A	610	LEU	2.1
1	A	628	ARG	2.1
1	A	852	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1001	14/15	0.79	0.14	44,51,60,62	0
3	CAC	A	1002	5/5	0.87	0.17	36,40,53,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.