



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:11 PM UTC

PDB ID : 1OFR / pdb_00001ofr
Title : CRYSTAL STRUCTURE OF THE TYROSINE-REGULATED 3-DEOXY-D-ARABINO-HEPTULOSONATE-7-PHOSPHATE SYNTHASE FROM SACCCHAROMYCES CEREVISIAE COMPLEXED WITH PHENYLALANINE AND MANGANESE
Authors : Koenig, V.; Pfeil, A.; Heinrich, G.; Braus, G.; Schneider, T.R.
Deposited on : 2003-04-18
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	370	
1	B	370	
1	C	370	
1	D	370	
1	E	370	
1	F	370	
1	G	370	
1	H	370	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	0	1
			2631	1636	474	511	10			
1	B	348	Total	C	N	O	S	0	0	1
			2618	1629	469	510	10			
1	C	349	Total	C	N	O	S	0	0	1
			2626	1634	471	511	10			
1	D	351	Total	C	N	O	S	0	0	1
			2623	1630	470	513	10			
1	E	339	Total	C	N	O	S	0	0	1
			2560	1591	460	499	10			
1	F	347	Total	C	N	O	S	0	0	1
			2604	1620	467	507	10			
1	G	344	Total	C	N	O	S	0	0	1
			2581	1605	462	504	10			
1	H	347	Total	C	N	O	S	0	0	1
			2599	1614	466	509	10			

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		
2	B	1	Total	Mn	0	0
			1	1		
2	C	1	Total	Mn	0	0
			1	1		
2	D	1	Total	Mn	0	0
			1	1		
2	E	1	Total	Mn	0	0
			1	1		
2	F	1	Total	Mn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Mn	0	0
			1	1		
2	H	1	Total	Mn	0	0
			1	1		

- Molecule 3 is PHENYLALANINE (CCD ID: PHE) (formula: $C_9H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			12	9	1	2		
3	D	1	Total	C	N	O	0	0
			12	9	1	2		
3	G	1	Total	C	N	O	0	0
			12	9	1	2		
3	H	1	Total	C	N	O	0	0
			12	9	1	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	O	0	0
			6	6		
4	B	4	Total	O	0	0
			4	4		
4	C	5	Total	O	0	0
			5	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	5	Total 5	O 5	0	0
4	E	2	Total 2	O 2	0	0
4	F	4	Total 4	O 4	0	0
4	G	3	Total 3	O 3	0	0
4	H	3	Total 3	O 3	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE




• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE




• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain C: 




• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE


Chain D: 



• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE


Chain E: 






• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain F: 59% 31% 6%



• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain G: 50% 36% 7% 7%



• Molecule 1: PHOSPHO-2-DEHYDRO-3-DEOXYHEPTONATE ALDOLASE

Chain H: 50% 39% 6%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	82.29Å 94.03Å 105.12Å 65.14° 85.68° 75.73°	Depositor
Resolution (Å)	37.20 – 2.70	Depositor
% Data completeness (in resolution range)	85.9 (37.20-2.70)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.232 , 0.274	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	20930	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	1/2669 (0.0%)	0.93	5/3613 (0.1%)
1	B	0.55	1/2654 (0.0%)	0.95	6/3590 (0.2%)
1	C	0.53	1/2664 (0.0%)	0.98	12/3605 (0.3%)
1	D	0.54	1/2661 (0.0%)	0.97	4/3605 (0.1%)
1	E	0.49	1/2596 (0.0%)	0.95	7/3513 (0.2%)
1	F	0.51	1/2641 (0.0%)	0.97	3/3575 (0.1%)
1	G	0.51	1/2617 (0.0%)	0.94	8/3544 (0.2%)
1	H	0.50	1/2637 (0.0%)	0.93	4/3577 (0.1%)
All	All	0.52	8/21139 (0.0%)	0.95	49/28622 (0.2%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	369	LYS	C-N	-5.68	1.25	1.33
1	E	369	LYS	C-N	-5.52	1.25	1.33
1	F	369	LYS	C-N	-5.52	1.25	1.33
1	A	369	LYS	C-N	-5.50	1.25	1.33
1	H	368	ASN	C-N	-5.39	1.25	1.33

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	309	ALA	N-CA-C	-8.09	103.61	113.97
1	G	303	ILE	N-CA-C	-7.48	103.57	110.82
1	C	125	ASP	CA-C-N	7.47	126.74	118.97
1	C	125	ASP	C-N-CA	7.47	126.74	118.97
1	D	104	ILE	N-CA-C	7.44	119.33	109.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2631	0	2634	130	0
1	B	2618	0	2618	136	0
1	C	2626	0	2631	130	0
1	D	2623	0	2605	124	0
1	E	2560	0	2562	127	0
1	F	2604	0	2596	104	0
1	G	2581	0	2564	154	0
1	H	2599	0	2576	139	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	C	12	0	8	0	0
3	D	12	0	8	0	0
3	G	12	0	8	1	0
3	H	12	0	8	1	0
4	A	6	0	0	2	0
4	B	4	0	0	0	0
4	C	5	0	0	1	0
4	D	5	0	0	1	0
4	E	2	0	0	0	0
4	F	4	0	0	1	0
4	G	3	0	0	0	0
4	H	3	0	0	0	0
All	All	20930	0	20818	994	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 994 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:ILE:HD12	1:C:134:ILE:H	1.19	1.05
1:B:134:ILE:H	1:B:134:ILE:HD12	1.19	1.05
1:E:324:GLN:HE21	1:E:338:VAL:HB	1.18	1.01
1:D:247:ILE:HD13	1:D:314:MET:HE1	1.45	0.96
1:C:247:ILE:HD13	1:C:314:MET:HE1	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/370 (94%)	298 (86%)	41 (12%)	7 (2%)	6	16
1	B	340/370 (92%)	296 (87%)	35 (10%)	9 (3%)	4	11
1	C	345/370 (93%)	302 (88%)	35 (10%)	8 (2%)	5	13
1	D	347/370 (94%)	299 (86%)	38 (11%)	10 (3%)	3	9
1	E	333/370 (90%)	294 (88%)	35 (10%)	4 (1%)	10	27
1	F	341/370 (92%)	307 (90%)	32 (9%)	2 (1%)	21	44
1	G	336/370 (91%)	287 (85%)	38 (11%)	11 (3%)	3	7
1	H	343/370 (93%)	298 (87%)	34 (10%)	11 (3%)	3	7
All	All	2731/2960 (92%)	2381 (87%)	288 (10%)	62 (2%)	5	13

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	111	GLU
1	B	22	ASP
1	B	286	ASN
1	B	324	GLN
1	C	323	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/298 (94%)	263 (94%)	17 (6%)	17	40
1	B	279/298 (94%)	264 (95%)	15 (5%)	20	45
1	C	280/298 (94%)	265 (95%)	15 (5%)	20	45
1	D	277/298 (93%)	265 (96%)	12 (4%)	26	54
1	E	275/298 (92%)	257 (94%)	18 (6%)	15	37
1	F	277/298 (93%)	259 (94%)	18 (6%)	15	37
1	G	274/298 (92%)	254 (93%)	20 (7%)	13	32
1	H	276/298 (93%)	264 (96%)	12 (4%)	26	54
All	All	2218/2384 (93%)	2091 (94%)	127 (6%)	18	43

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	341	THR
1	G	317	SER
1	E	286	ASN
1	G	307	GLU
1	H	116	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 80 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	320	ASN
1	H	147	ASN
1	G	124	ASN
1	G	286	ASN
1	H	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PHE	G	1002	-	11,12,12	0.74	0	11,15,15	0.31	0
3	PHE	H	1002	-	11,12,12	0.68	0	11,15,15	0.32	0
3	PHE	C	1002	-	11,12,12	0.75	0	11,15,15	0.34	0
3	PHE	D	1002	-	11,12,12	0.84	0	11,15,15	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PHE	G	1002	-	-	4/8/8/8	0/1/1/1
3	PHE	H	1002	-	-	2/8/8/8	0/1/1/1
3	PHE	C	1002	-	-	3/8/8/8	0/1/1/1
3	PHE	D	1002	-	-	2/8/8/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	1002	PHE	OXT-C-CA-N
3	D	1002	PHE	OXT-C-CA-N
3	G	1002	PHE	O-C-CA-CB
3	H	1002	PHE	OXT-C-CA-N
3	G	1002	PHE	OXT-C-CA-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1002	PHE	1	0
3	H	1002	PHE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.