



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:02 PM UTC

PDB ID : 2OF7 / pdb_00002of7
Title : Structural Genomics, the crystal structure of a tetR-family transcriptional regulator from Streptomyces coelicolor A3
Authors : Tan, K.; Xu, X.; Zheng, H.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2007-01-02
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	5% 32% 25% 12% • 27%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative tetR-family transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	Se	0	0	0
			1489	920	276	287	6			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MSE	-	cloning artifact	UNP Q9EWH2
A	-21	THR	-	cloning artifact	UNP Q9EWH2
A	-20	MSE	-	cloning artifact	UNP Q9EWH2
A	-19	GLY	-	cloning artifact	UNP Q9EWH2
A	-18	SER	-	cloning artifact	UNP Q9EWH2
A	-17	SER	-	cloning artifact	UNP Q9EWH2
A	-16	HIS	-	cloning artifact	UNP Q9EWH2
A	-15	HIS	-	cloning artifact	UNP Q9EWH2
A	-14	HIS	-	cloning artifact	UNP Q9EWH2
A	-13	HIS	-	cloning artifact	UNP Q9EWH2
A	-12	HIS	-	cloning artifact	UNP Q9EWH2
A	-11	HIS	-	cloning artifact	UNP Q9EWH2
A	-10	SER	-	cloning artifact	UNP Q9EWH2
A	-9	SER	-	cloning artifact	UNP Q9EWH2
A	-8	GLY	-	cloning artifact	UNP Q9EWH2
A	-7	ARG	-	cloning artifact	UNP Q9EWH2
A	-6	GLU	-	cloning artifact	UNP Q9EWH2
A	-5	ASN	-	cloning artifact	UNP Q9EWH2
A	-4	LEU	-	cloning artifact	UNP Q9EWH2
A	-3	TYR	-	cloning artifact	UNP Q9EWH2
A	-2	PHE	-	cloning artifact	UNP Q9EWH2
A	-1	GLN	-	cloning artifact	UNP Q9EWH2
A	0	GLY	-	cloning artifact	UNP Q9EWH2
A	1	HIS	-	cloning artifact	UNP Q9EWH2
A	2	MSE	-	cloning artifact	UNP Q9EWH2
A	82	MSE	MET	modified residue	UNP Q9EWH2
A	135	MSE	MET	modified residue	UNP Q9EWH2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	MSE	MET	modified residue	UNP Q9EWH2
A	145	MSE	MET	modified residue	UNP Q9EWH2
A	168	MSE	MET	modified residue	UNP Q9EWH2
A	175	MSE	MET	modified residue	UNP Q9EWH2
A	236	GLY	-	cloning artifact	UNP Q9EWH2
A	237	SER	-	cloning artifact	UNP Q9EWH2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	53	Total O 53 53	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative tetR-family transcriptional regulator



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	75.10Å 73.24Å 48.08Å 90.00° 107.27° 90.00°	Depositor
Resolution (Å)	36.61 – 2.30 36.61 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (36.61-2.30) 98.5 (36.61-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.19 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.231 , 0.293 0.229 , 0.294	Depositor DCC
R_{free} test set	526 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	40.7	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1542	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.61	15/1503 (1.0%)	1.66	28/2023 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	193	LEU	C-N	-17.53	1.11	1.34
1	A	192	GLU	C-N	-17.51	1.11	1.33
1	A	170	LEU	CA-C	-6.43	1.44	1.53
1	A	102	VAL	CA-CB	6.22	1.63	1.54
1	A	106	ALA	CA-CB	6.13	1.63	1.53
1	A	175	MSE	CA-C	6.08	1.60	1.52
1	A	142	THR	CA-CB	6.07	1.63	1.53
1	A	166	VAL	CA-CB	5.91	1.60	1.54
1	A	157	ASP	CA-C	5.87	1.60	1.52
1	A	167	SER	CA-C	5.75	1.60	1.52
1	A	142	THR	CA-C	5.58	1.60	1.52
1	A	81	VAL	CA-CB	5.13	1.61	1.54
1	A	188	GLU	N-CA	-5.04	1.39	1.46
1	A	163	VAL	N-CA	-5.02	1.39	1.46
1	A	198	LEU	C-O	-5.01	1.18	1.24

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	LEU	CA-C-N	10.65	152.56	127.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	LEU	C-N-CA	10.65	152.56	127.00
1	A	157	ASP	CA-C-N	10.52	130.85	119.28
1	A	157	ASP	C-N-CA	10.52	130.85	119.28
1	A	193	LEU	O-C-N	-8.92	112.89	122.07
1	A	205	LEU	C-N-CD	-8.11	102.75	120.60
1	A	18	LEU	N-CA-C	-8.07	100.49	111.55
1	A	142	THR	CB-CA-C	7.64	125.63	110.42
1	A	206	PRO	N-CA-C	7.38	131.29	112.10
1	A	77	GLU	N-CA-C	7.20	122.07	113.28
1	A	128	VAL	CA-C-N	-6.50	112.30	119.19
1	A	128	VAL	C-N-CA	-6.50	112.30	119.19
1	A	169	SER	CB-CA-C	6.13	122.81	109.99
1	A	154	THR	N-CA-C	-6.12	106.41	114.31
1	A	58	SER	CA-C-N	5.94	127.27	119.84
1	A	58	SER	C-N-CA	5.94	127.27	119.84
1	A	175	MSE	CG-SE-CE	-5.82	86.13	98.92
1	A	123	ARG	NE-CZ-NH2	5.75	124.37	119.20
1	A	22	LYS	N-CA-C	-5.74	105.11	111.36
1	A	113	GLU	N-CA-C	-5.68	106.01	113.17
1	A	124	LEU	N-CA-C	5.67	117.14	111.07
1	A	206	PRO	CA-N-CD	-5.61	103.64	111.50
1	A	107	LEU	N-CA-C	5.58	120.15	113.23
1	A	166	VAL	CB-CA-C	-5.49	104.85	111.88
1	A	157	ASP	CB-CA-C	5.29	118.36	109.69
1	A	120	LEU	N-CA-C	-5.23	105.25	111.69
1	A	153	ARG	N-CA-C	5.03	119.14	113.15
1	A	190	LEU	N-CA-C	-5.02	105.81	111.28

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	141	ASP	Peptide
1	A	89	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1489	0	1496	95	0
2	A	53	0	0	9	0
All	All	1542	0	1496	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:THR:CG2	1:A:143:GLY:HA2	1.57	1.34
1:A:142:THR:HG23	1:A:143:GLY:CA	1.64	1.27
1:A:100:ARG:HG2	1:A:100:ARG:HH11	1.10	1.10
1:A:141:ASP:O	1:A:142:THR:HB	1.55	1.06
1:A:100:ARG:HH11	1:A:100:ARG:CG	1.68	1.02
1:A:105:LYS:HD2	2:A:267:HOH:O	1.69	0.92
1:A:129:PRO:O	1:A:133:ALA:N	2.05	0.89
1:A:91:ALA:O	1:A:93:GLU:N	2.06	0.89
1:A:31:ARG:HH22	1:A:117:LEU:HD21	1.38	0.88
1:A:98:SER:O	1:A:102:VAL:HG13	1.77	0.84
1:A:92:GLY:HA2	2:A:281:HOH:O	1.78	0.83
1:A:123:ARG:O	1:A:127:GLU:HG2	1.82	0.79
1:A:22:LYS:O	1:A:26:THR:HG22	1.83	0.79
1:A:23:LYS:HE3	1:A:65:TYR:HE1	1.49	0.78
1:A:139:MSE:CE	1:A:168:MSE:SE	2.82	0.77
1:A:100:ARG:CG	1:A:100:ARG:NH1	2.41	0.76
1:A:85:GLU:OE2	1:A:101:HIS:NE2	2.19	0.76
1:A:166:VAL:O	1:A:170:LEU:HG	1.85	0.75
1:A:31:ARG:HH22	1:A:117:LEU:CD2	2.00	0.74
1:A:91:ALA:C	1:A:93:GLU:H	1.95	0.74
1:A:139:MSE:HE3	1:A:168:MSE:SE	2.39	0.72
1:A:89:ARG:C	1:A:91:ALA:H	1.99	0.71
1:A:100:ARG:HG2	1:A:100:ARG:NH1	1.96	0.71
1:A:205:LEU:HD12	2:A:268:HOH:O	1.89	0.70
1:A:140:SER:O	1:A:142:THR:N	2.25	0.69
1:A:128:VAL:HG12	1:A:131:VAL:H	1.60	0.67
1:A:135:MSE:O	1:A:139:MSE:HB2	1.96	0.66
1:A:186:HIS:HD2	2:A:238:HOH:O	1.80	0.65
1:A:142:THR:HG22	2:A:254:HOH:O	1.95	0.64
1:A:91:ALA:C	1:A:93:GLU:N	2.53	0.64
1:A:20:GLU:O	1:A:24:THR:HG23	1.98	0.64
1:A:68:THR:O	1:A:71:ASP:HB2	1.98	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:HIS:HD2	1:A:187:GLU:OE2	1.82	0.63
1:A:136:LEU:HD12	1:A:139:MSE:SE	2.49	0.62
1:A:139:MSE:HE2	1:A:168:MSE:SE	2.48	0.62
1:A:142:THR:HG23	1:A:143:GLY:HA2	0.75	0.62
1:A:48:VAL:HG12	2:A:277:HOH:O	2.00	0.60
1:A:52:ALA:HB1	1:A:57:VAL:O	2.01	0.59
1:A:100:ARG:NH1	1:A:104:ARG:HD2	2.18	0.59
1:A:151:ALA:O	1:A:155:GLY:N	2.35	0.59
1:A:188:GLU:CD	1:A:196:ARG:HH21	2.10	0.58
1:A:31:ARG:NH2	1:A:117:LEU:HD21	2.15	0.57
1:A:186:HIS:H	1:A:186:HIS:CD2	2.22	0.57
1:A:147:ALA:HB1	1:A:163:VAL:HG13	1.86	0.57
1:A:137:GLU:O	1:A:138:ASN:C	2.48	0.55
1:A:23:LYS:HE2	1:A:27:ARG:HH22	1.70	0.55
1:A:141:ASP:O	1:A:142:THR:CB	2.44	0.53
1:A:23:LYS:HE3	1:A:65:TYR:CE1	2.37	0.53
1:A:77:GLU:O	1:A:78:TYR:HB2	2.08	0.53
1:A:36:GLY:HA2	1:A:39:ARG:NH2	2.24	0.53
1:A:41:GLN:HE22	1:A:50:GLN:HE22	1.56	0.52
1:A:140:SER:C	1:A:142:THR:N	2.68	0.52
1:A:139:MSE:O	1:A:139:MSE:HG2	2.10	0.52
1:A:40:GLN:HB2	1:A:41:GLN:HG2	1.91	0.51
1:A:41:GLN:O	1:A:45:ALA:HB3	2.11	0.51
1:A:144:ARG:HH11	1:A:144:ARG:HG3	1.76	0.51
1:A:144:ARG:HH11	1:A:144:ARG:CG	2.25	0.50
1:A:89:ARG:HB2	1:A:153:ARG:NH1	2.27	0.50
1:A:75:THR:O	1:A:77:GLU:HG3	2.12	0.49
1:A:136:LEU:CD1	1:A:139:MSE:SE	3.09	0.49
1:A:165:ILE:O	1:A:169:SER:HB3	2.12	0.49
1:A:142:THR:HG23	1:A:143:GLY:C	2.34	0.49
1:A:40:GLN:HB2	1:A:41:GLN:CG	2.43	0.49
1:A:148:ARG:NH1	2:A:282:HOH:O	2.45	0.48
1:A:142:THR:CB	1:A:143:GLY:HA2	2.39	0.48
1:A:151:ALA:O	1:A:152:ASP:C	2.55	0.48
1:A:89:ARG:C	1:A:91:ALA:N	2.67	0.48
1:A:140:SER:C	1:A:142:THR:H	2.21	0.48
1:A:142:THR:CG2	2:A:254:HOH:O	2.56	0.47
1:A:122:THR:HG23	1:A:178:SER:HB2	1.97	0.46
1:A:19:ARG:HH22	1:A:23:LYS:HD2	1.80	0.46
1:A:89:ARG:HD2	1:A:89:ARG:HA	1.69	0.46
1:A:99:LEU:HD11	1:A:170:LEU:HD13	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:MSE:O	1:A:139:MSE:CG	2.64	0.46
1:A:136:LEU:O	1:A:139:MSE:HB3	2.15	0.46
1:A:142:THR:CG2	1:A:143:GLY:CA	2.52	0.46
1:A:143:GLY:C	1:A:145:MSE:H	2.24	0.45
1:A:100:ARG:O	1:A:101:HIS:C	2.56	0.45
1:A:105:LYS:CE	2:A:271:HOH:O	2.65	0.44
1:A:143:GLY:O	1:A:144:ARG:HB2	2.16	0.44
1:A:94:PRO:HG2	1:A:97:ASP:OD2	2.17	0.44
1:A:157:ASP:HA	1:A:158:PRO:HD2	1.44	0.44
1:A:78:TYR:O	1:A:82:MSE:HG3	2.18	0.43
1:A:184:HIS:CD2	1:A:187:GLU:OE2	2.67	0.43
1:A:25:ARG:HD3	1:A:25:ARG:HA	1.60	0.43
1:A:129:PRO:HA	1:A:132:ARG:HB3	2.01	0.43
1:A:132:ARG:O	1:A:136:LEU:HB2	2.19	0.42
1:A:203:ASN:O	1:A:206:PRO:HG2	2.19	0.42
1:A:103:LEU:HD11	1:A:174:LEU:HB2	2.02	0.42
1:A:156:LEU:O	1:A:158:PRO:HD3	2.20	0.42
1:A:64:ARG:HH11	1:A:64:ARG:HD3	1.73	0.41
1:A:174:LEU:HD12	1:A:174:LEU:C	2.45	0.41
1:A:176:GLU:OE2	1:A:176:GLU:HA	2.20	0.41
1:A:137:GLU:C	1:A:139:MSE:N	2.76	0.41
1:A:78:TYR:CA	1:A:81:VAL:HG22	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/260 (73%)	167 (88%)	15 (8%)	7 (4%)	2 1

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	78	TYR
1	A	92	GLY
1	A	141	ASP
1	A	142	THR
1	A	206	PRO
1	A	88	ALA
1	A	94	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	153/199 (77%)	118 (77%)	35 (23%)	1 1

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	20	GLU
1	A	22	LYS
1	A	26	THR
1	A	31	ARG
1	A	39	ARG
1	A	41	GLN
1	A	48	VAL
1	A	57	VAL
1	A	58	SER
1	A	63	LEU
1	A	74	LEU
1	A	81	VAL
1	A	100	ARG
1	A	102	VAL
1	A	105	LYS
1	A	117	LEU
1	A	120	LEU
1	A	127	GLU
1	A	134	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	136	LEU
1	A	137	GLU
1	A	140	SER
1	A	142	THR
1	A	144	ARG
1	A	146	LEU
1	A	153	ARG
1	A	156	LEU
1	A	163	VAL
1	A	169	SER
1	A	174	LEU
1	A	196	ARG
1	A	199	ASP
1	A	205	LEU
1	A	206	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	A	184	HIS
1	A	186	HIS
1	A	203	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	192:GLU	C	193:LEU	N	1.11
1	A	193:LEU	C	194:VAL	N	1.11

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/260 (71%)	0.68	13 (7%) 22 24	23, 48, 71, 92	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	LEU	4.3
1	A	207	ALA	3.3
1	A	206	PRO	3.2
1	A	140	SER	3.2
1	A	205	LEU	2.9
1	A	76	ASP	2.6
1	A	90	PRO	2.5
1	A	112	GLY	2.5
1	A	170	LEU	2.4
1	A	91	ALA	2.4
1	A	95	TRP	2.2
1	A	78	TYR	2.2
1	A	136	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.