



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 12:32 PM UTC

PDB ID : 3OFS / pdb_00003ofs
Title : Dynamic conformations of the CD38-mediated NAD cyclization captured using multi-copy crystallography
Authors : Zhang, H.; Lee, H.C.; Hao, Q.
Deposited on : 2010-08-16
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	 4% 79% 12% 9%
1	B	257	 5% 81% 10% 9%
1	C	257	 2% 81% 10% 8%
1	D	257	 2% 77% 13% 9%
1	E	257	 5% 80% 11% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	257	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '6%', a large green segment labeled '76%', a yellow segment labeled '14%', and a small grey segment at the end labeled '9%'.</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosyl cyclase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	1893	1198	326	355	14	0	0	0
1	B	234	1889	1196	326	353	14	0	0	0
1	C	236	1908	1206	332	356	14	0	0	0
1	D	233	1883	1191	328	350	14	0	0	0
1	E	236	1898	1201	328	355	14	0	0	0
1	F	233	1875	1187	322	352	14	0	0	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLU	-	expression tag	UNP P28907
A	45	PHE	-	expression tag	UNP P28907
A	49	THR	GLN	engineered mutation	UNP P28907
A	100	ASP	ASN	engineered mutation	UNP P28907
A	146	ALA	GLU	engineered mutation	UNP P28907
A	164	ASP	ASN	engineered mutation	UNP P28907
A	209	ASP	ASN	engineered mutation	UNP P28907
A	219	ASP	ASN	engineered mutation	UNP P28907
B	44	GLU	-	expression tag	UNP P28907
B	45	PHE	-	expression tag	UNP P28907
B	49	THR	GLN	engineered mutation	UNP P28907
B	100	ASP	ASN	engineered mutation	UNP P28907
B	146	ALA	GLU	engineered mutation	UNP P28907
B	164	ASP	ASN	engineered mutation	UNP P28907
B	209	ASP	ASN	engineered mutation	UNP P28907
B	219	ASP	ASN	engineered mutation	UNP P28907
C	44	GLU	-	expression tag	UNP P28907

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	45	PHE	-	expression tag	UNP P28907
C	49	THR	GLN	engineered mutation	UNP P28907
C	100	ASP	ASN	engineered mutation	UNP P28907
C	146	ALA	GLU	engineered mutation	UNP P28907
C	164	ASP	ASN	engineered mutation	UNP P28907
C	209	ASP	ASN	engineered mutation	UNP P28907
C	219	ASP	ASN	engineered mutation	UNP P28907
D	44	GLU	-	expression tag	UNP P28907
D	45	PHE	-	expression tag	UNP P28907
D	49	THR	GLN	engineered mutation	UNP P28907
D	100	ASP	ASN	engineered mutation	UNP P28907
D	146	ALA	GLU	engineered mutation	UNP P28907
D	164	ASP	ASN	engineered mutation	UNP P28907
D	209	ASP	ASN	engineered mutation	UNP P28907
D	219	ASP	ASN	engineered mutation	UNP P28907
E	44	GLU	-	expression tag	UNP P28907
E	45	PHE	-	expression tag	UNP P28907
E	49	THR	GLN	engineered mutation	UNP P28907
E	100	ASP	ASN	engineered mutation	UNP P28907
E	146	ALA	GLU	engineered mutation	UNP P28907
E	164	ASP	ASN	engineered mutation	UNP P28907
E	209	ASP	ASN	engineered mutation	UNP P28907
E	219	ASP	ASN	engineered mutation	UNP P28907
F	44	GLU	-	expression tag	UNP P28907
F	45	PHE	-	expression tag	UNP P28907
F	49	THR	GLN	engineered mutation	UNP P28907
F	100	ASP	ASN	engineered mutation	UNP P28907
F	146	ALA	GLU	engineered mutation	UNP P28907
F	164	ASP	ASN	engineered mutation	UNP P28907
F	209	ASP	ASN	engineered mutation	UNP P28907
F	219	ASP	ASN	engineered mutation	UNP P28907


- Molecule 2 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methyl [(2R,3R,4R)-4-fluoro-3-hydroxytetrahydrofuran-2-yl]methyl dihydrogen diphosphate (CCD ID: AVU) (formula: C₁₅H₂₂FN₅O₁₂P₂).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			P
2	A	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0
2	B	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0
2	C	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0
2	D	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0
2	E	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0
2	F	1	Total 35	C 15	F 1	N 5	O 12	P 2	0	0

- Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total 58	O 58	0	0
3	B	47	Total 47	O 47	0	0
3	C	77	Total 77	O 77	0	0
3	D	80	Total 80	O 80	0	0
3	E	71	Total 71	O 71	0	0
3	F	59	Total 59	O 59	0	0



• Molecule 1: ADP-ribosyl cyclase 1



• Molecule 1: ADP-ribosyl cyclase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.66Å 146.66Å 191.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.73 – 2.20 48.73 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.2 (48.73-2.20) 97.2 (48.73-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.10 (at 2.20Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.203 , 0.250 0.223 , 0.264	Depositor DCC
R_{free} test set	3969 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtrriage
Anisotropy	0.058	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11948	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0668e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AVU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	1/1941 (0.1%)	0.78	0/2633
1	B	0.57	0/1937	0.75	1/2628 (0.0%)
1	C	0.61	0/1957	0.78	0/2655
1	D	0.62	0/1930	0.76	1/2616 (0.0%)
1	E	0.61	0/1947	0.78	0/2643
1	F	0.59	0/1922	0.74	1/2607 (0.0%)
All	All	0.60	1/11634 (0.0%)	0.77	3/15782 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	231	GLN	CD-OE1	5.81	1.34	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	225	VAL	N-CA-C	5.90	116.64	110.62
1	F	225	VAL	N-CA-C	5.58	116.36	110.72
1	D	225	VAL	N-CA-C	5.30	115.51	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1893	0	1820	21	0
1	B	1889	0	1816	19	0
1	C	1908	0	1839	20	0
1	D	1883	0	1817	26	0
1	E	1898	0	1822	32	0
1	F	1875	0	1799	27	0
2	A	35	0	19	2	0
2	B	35	0	19	2	0
2	C	35	0	19	0	0
2	D	35	0	19	3	0
2	E	35	0	19	2	0
2	F	35	0	19	3	0
3	A	58	0	0	0	0
3	B	47	0	0	0	0
3	C	77	0	0	1	0
3	D	80	0	0	2	0
3	E	71	0	0	1	0
3	F	59	0	0	0	0
All	All	11948	0	11027	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:165:THR:HG23	1:C:167:LYS:H	1.31	0.95
1:A:162:GLU:CG	1:A:165:THR:HG22	2.02	0.88
1:A:165:THR:HG23	1:A:167:LYS:H	1.44	0.82
1:A:162:GLU:HG3	1:A:165:THR:HG22	1.62	0.81
1:B:241:TRP:HB3	1:B:278:ILE:HD12	1.66	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/257 (90%)	222 (96%)	6 (3%)	2 (1%)	14	14
1	B	230/257 (90%)	221 (96%)	8 (4%)	1 (0%)	30	34
1	C	234/257 (91%)	228 (97%)	5 (2%)	1 (0%)	30	34
1	D	229/257 (89%)	223 (97%)	6 (3%)	0	100	100
1	E	234/257 (91%)	221 (94%)	11 (5%)	2 (1%)	14	14
1	F	229/257 (89%)	223 (97%)	5 (2%)	1 (0%)	30	34
All	All	1386/1542 (90%)	1338 (96%)	41 (3%)	7 (0%)	24	27

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	GLU
1	E	233	GLU
1	E	128	ILE
1	A	128	ILE
1	F	128	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/237 (90%)	210 (99%)	3 (1%)	59	75
1	B	212/237 (90%)	209 (99%)	3 (1%)	59	75
1	C	214/237 (90%)	210 (98%)	4 (2%)	50	66
1	D	211/237 (89%)	206 (98%)	5 (2%)	43	58
1	E	212/237 (90%)	209 (99%)	3 (1%)	59	75
1	F	210/237 (89%)	207 (99%)	3 (1%)	59	75
All	All	1272/1422 (90%)	1251 (98%)	21 (2%)	53	69

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	231	GLN
1	E	233	GLU
1	F	227	VAL
1	F	107	GLN
1	E	231	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	270	ASN
1	F	255	GLN
1	D	74	HIS
1	D	83	GLN
1	E	83	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AVU	B	301	1	38,38,38	1.37	4 (10%)	54,58,58	1.56	9 (16%)
2	AVU	E	301	1	38,38,38	1.24	5 (13%)	54,58,58	1.73	11 (20%)
2	AVU	F	301	1	38,38,38	1.52	6 (15%)	54,58,58	1.72	9 (16%)
2	AVU	C	301	1	38,38,38	1.18	5 (13%)	54,58,58	1.57	7 (12%)
2	AVU	D	301	1	38,38,38	1.21	3 (7%)	54,58,58	1.85	14 (25%)
2	AVU	A	301	1	38,38,38	1.29	4 (10%)	54,58,58	1.64	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AVU	B	301	1	-	6/22/51/51	0/4/4/4
2	AVU	E	301	1	-	8/22/51/51	0/4/4/4
2	AVU	F	301	1	-	5/22/51/51	0/4/4/4
2	AVU	C	301	1	-	2/22/51/51	0/4/4/4
2	AVU	D	301	1	-	3/22/51/51	0/4/4/4
2	AVU	A	301	1	-	4/22/51/51	0/4/4/4

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	AVU	PA-O3A	4.57	1.64	1.59
2	F	301	AVU	PA-O3A	4.28	1.64	1.59
2	F	301	AVU	PB-O3A	4.26	1.64	1.59
2	B	301	AVU	PA-O3A	4.07	1.63	1.59
2	B	301	AVU	PB-O3A	3.66	1.63	1.59

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	301	AVU	C5-C4-N3	-5.26	119.48	126.72
2	D	301	AVU	N3-C2-N1	-5.16	120.77	128.58
2	F	301	AVU	N3-C2-N1	-5.15	120.78	128.58
2	C	301	AVU	N3-C2-N1	-5.12	120.83	128.58
2	F	301	AVU	C5-C4-N3	-5.10	119.69	126.72

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	AVU	C5R-O5R-PB-O2B
2	B	301	AVU	C5'-O5'-PA-O1A
2	B	301	AVU	C5R-O5R-PB-O2B
2	B	301	AVU	C5R-O5R-PB-O3A
2	D	301	AVU	O4R-C4R-C5R-O5R


There are no ring outliers.


5 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	AVU	2	0
2	E	301	AVU	2	0
2	F	301	AVU	3	0
2	D	301	AVU	3	0
2	A	301	AVU	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/257 (91%)	0.37	10 (4%) 40 36	29, 45, 74, 91	0
1	B	234/257 (91%)	0.48	13 (5%) 30 27	28, 48, 84, 98	0
1	C	236/257 (91%)	0.25	6 (2%) 58 55	28, 42, 66, 85	0
1	D	233/257 (90%)	0.26	5 (2%) 63 60	25, 44, 67, 81	0
1	E	236/257 (91%)	0.40	12 (5%) 33 30	25, 45, 87, 108	0
1	F	233/257 (90%)	0.53	16 (6%) 23 20	28, 48, 89, 127	0
All	All	1406/1542 (91%)	0.38	62 (4%) 39 36	25, 45, 78, 127	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	138	VAL	4.7
1	B	245	GLY	4.2
1	E	114	THR	4.0
1	B	248	GLU	3.6
1	E	120	ASN	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	AVU	A	301	35/35	0.89	0.15	34,52,68,69	0
2	AVU	C	301	35/35	0.89	0.12	31,46,63,64	0
2	AVU	D	301	35/35	0.89	0.12	33,48,55,56	0
2	AVU	B	301	35/35	0.92	0.11	33,49,66,67	0
2	AVU	F	301	35/35	0.92	0.11	31,45,55,57	0
2	AVU	E	301	35/35	0.94	0.10	33,44,53,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AVU A 301:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around AVU D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AVU B 301:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AVU F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.